

LARGE WEIGHT DOES NOT YIELD AN IRREDUCIBLE BASE

SAHARON SHELAH¹

Institute of Mathematics, Hebrew University

Jerusalem 91904, Israel

E-mail: shelah@math.huji.ac.il

(Received October 20, 2010; Accepted March 5, 2012)

[Communicated by Lajos Soukup]

Abstract

Answering a question of Juhász, Soukup and Szentmiklóssy, we show that it is consistent that some first countable space of uncountable weight does not contain an uncountable subspace which has an irreducible base.

1. Introduction

For a topological space X , $w(X)$ is the minimal cardinality of a base for X , $\chi(p, X) = \min\{|u| : u \text{ is a neighbourhood base of } p\}$, and $\chi(X) = \sup\{\chi(p, X) : p \in X\}$.

In [1] the following problem was investigated: What makes a space have weight larger than its character? The notion of *irreducible base* was introduced, and it was proved [1, Lemma 2.6] that *if a topological space X has an irreducible base then $w(X) = |X| \cdot \chi(X)$* . The following question was formulated:

PROBLEM 1. *Does every first countable space of uncountable weight contain an uncountable subspace which has an irreducible base?*

We show that the answer is consistently NO. We thank Lajos Soukup for actually writing the paper.

DEFINITION 1.1. Let X be a topological space. A base \mathcal{U} of X is called *irreducible* if it has an *irreducible decomposition* $\mathcal{U} = \bigcup\{\mathcal{U}_x : x \in X\}$, i.e. (i) and (ii) below hold:

- (i) \mathcal{U}_x is a neighbourhood base of x in X for each $x \in X$;
- (ii) for each $x \in X$ the family $\mathcal{U}_x^- = \bigcup_{y \neq x} \mathcal{U}_y$ is not a base of X .

Mathematics subject classification number: 54E25.

Key words and phrases: general topology, weight, irreducible base, forcing.

¹ Research supported by the United States–Israel Binational Science Foundation. Paper no. 588 on the author’s publication list.

THEOREM 1.2. *There is a c.c.c poset $P = \langle P, \leq \rangle$ of size ω_1 such that in V^P there is a first countable space $X = \langle \omega_1, \tau \rangle$ of uncountable weight which does not contain an uncountable subspace which has an irreducible base.*

PROOF. The elements of the poset P will be finite ‘‘approximations’’ of a base $\{U(\alpha, n) : \alpha < \omega_1, n < \omega\}$ of X .

We define the poset $P = \langle P, \leq \rangle$ as follows. The underlying set of P consists of the triples $\langle A, n, U \rangle$ satisfying (P1)–(P3) below:

- (P1) $A \in [\omega_1]^{<\omega}$, $n \in \omega$ and U is a function, $U: A \times n \rightarrow \mathcal{P}(A)$,
- (P2) $\alpha \in U(\alpha, j)$ and $U(\alpha, i) \subset U(\alpha, i-1)$ for each $\alpha \in A$ and $j < n$, $0 < i < n$,
- (P3) if $\beta \in U(\alpha, i) \subset U(\beta, 0)$ for some $i < n$, then $\beta \leq \alpha$.

For $p \in P$ write $p = \langle A_p, n_p, U_p \rangle$. Let us remark that property (P3) will guarantee that $w(X) = \omega_1$.

Define the order \leq on P as follows. For $p, q \in P$ we put $q \leq p$ if

- (a) $A_p \subset A_q$,
- (b) $n_p \leq n_q$,
- (c) $U_p(\alpha, i) = U_q(\alpha, i) \cap A_p$ for each $\langle \alpha, i \rangle \in A_p \times n_p$,
- (d) for each $\langle \alpha, i \rangle, \langle \beta, j \rangle \in A_p \times n_p$,

$$\text{if } U_p(\alpha, i) \cap U_p(\beta, j) = \emptyset \text{ then } U_q(\alpha, i) \cap U_q(\beta, j) = \emptyset, \quad (\text{d1})$$

$$\text{if } U_p(\alpha, i) \subset U_p(\beta, j) \text{ then } U_q(\alpha, i) \subset U_q(\beta, j). \quad (\text{d2})$$

We say that the conditions $p_0 = \langle A_0, n_0, U_0 \rangle$ and $p_1 = \langle A_1, n_1, U_1 \rangle$ are *twins* iff $n_0 = n_1$, $|A_0| = |A_1|$ and denoting by σ the unique $<_{\mathcal{O}_n}$ -preserving bijection between A_0 and A_1 we have

- (I1) $\sigma|_{A_0 \cap A_1} = \text{id}_{A_0 \cap A_1}$,
- (I2) σ is an isomorphism between p_0 and p_1 , i.e. for each $\alpha \in A_0$ and $i < n_0$ we have $U_1(\sigma(\alpha), i) = \sigma''U_0(\alpha, i)$.

We say that σ is the *twin function* between p_0 and p_1 . Define the *smashing function* $\underline{\sigma}$ of p_0 and p_1 as follows: $\underline{\sigma} = \sigma^{-1} \cup \text{id}_{A_0}$. The function σ^* defined by the formula $\sigma^* = \sigma \cup \sigma^{-1}$ is called the *exchange function* of p_0 and p_1 .

The burden of the proof is to verify the next lemma. □

LEMMA 1.3. (Amalgamation lemma) *Assume that $p_0 = \langle A_0, n_0, U_0 \rangle$ and $p_1 = \langle A_1, n_1, U_1 \rangle$ are twins, $A_0 \cap A_1 < A_0 \setminus A_1 < A_1 \setminus A_0$, $\xi_0 \in A_0 \setminus A_1$, $\xi_1 = \sigma(\xi_0)$, where σ is the twin function between p_0 and p_1 , and let $k < m < n_0$. Then p_0 and p_1 have a common extension $p = \langle A, n, U \rangle$ in P such that*

$$\xi_0 \in U(\xi_1, m) \subset U(\xi_1, k) \subset U(\xi_0, k). \quad (*)$$

PROOF. Write $n = n_0 = n_1$, $D = A_0 \cap A_1$ and $A^* = A_0 \cup A_1$. Unfortunately we can not assume that $A = A^*$ because in this case we can not guarantee (P3) for p . So we need to add further elements to A^* to get a large enough A as follows. Choose a set $B \subset \omega_1 \setminus A^*$ of cardinality $|A^* \times n|$ and fix a bijection ρ between $A^* \times n$ and B . We will take $A = A^* \cup B$. To simplify the notation we will write $\langle \alpha, i \rangle$ for $\rho(\alpha, i)$, for all $\alpha \in A^*$ and $i < n$, i.e. we identify the elements of B and of $A^* \times n$.

The idea of the proof is the following: for each $\langle \alpha, i \rangle \in A^* \times n$ we put the element $\langle \alpha, i \rangle$ into $U(\alpha, i)$. On the other hand, we try to keep $U(\alpha, i)$ small, so we put $\langle \beta, j \rangle$ into $U(\alpha, i)$ if and only if we can “derive” from the property (d2) that $U(\beta, j) \subset U(\alpha, i)$ should hold in any condition $p = \langle A, n, U \rangle$ which is a common extension of p_0 and p_1 and which satisfies (*).

The condition p will be constructed in two steps. First we construct a condition $p' = \langle A, n, U' \rangle$ extending both p_0 and p_1 . This p' can be considered as the minimal amalgamation of p_0 and p_1 . Then, in the second step, we carry out small modifications on the function U' , namely we increase its value on certain places to guarantee (*).

Now we carry out our construction. For $\varepsilon < 2$ and $\langle \beta, j \rangle \in A_\varepsilon \times n$ let

$$V_\varepsilon(\beta, j) = \{\langle \alpha, i \rangle \in A_\varepsilon \times n : U_\varepsilon(\alpha, i) \subset U_\varepsilon(\beta, j)\} \quad (1)$$

and

$$W_\varepsilon(\beta, j) = \{\langle \alpha, i \rangle \in A_{1-\varepsilon} \times n : \exists \langle \gamma, l \rangle \in D \times n \\ U_{1-\varepsilon}(\alpha, i) \subset U_{1-\varepsilon}(\gamma, l) \wedge U_\varepsilon(\gamma, l) \subset U_\varepsilon(\beta, j)\}. \quad (2)$$

If we want to define p' in such a way that $p' \leq p_0, p_1$, then (d2) implies that $U'(\alpha, i) \subset U'(\beta, j)$ should hold whenever $\langle \alpha, i \rangle \in V(\beta, j) \cup W(\beta, j)$.

Now we are ready to define the function U' . For $\varepsilon < 2$, $\beta \in A_\varepsilon$ and $j < n$ let

$$U'(\beta, j) = U_\varepsilon(\beta, j) \cup U_{1-\varepsilon}(\sigma^*(\beta), j) \cup V_\varepsilon(\beta, j) \cup W_\varepsilon(\beta, j). \quad (3)$$

For $\langle \alpha, i \rangle \in A^* \times n$ and $j < n$ let

$$U'(\langle \alpha, i \rangle, j) = \{\langle \alpha, i \rangle\}. \quad (4)$$

Let us remark that $U'(\delta, j)$ is well defined even for $\delta \in A_0 \cap A_1$. Indeed, in this case $\sigma^*(\delta) = \delta$ and $V_\varepsilon(\delta, j) = W_{1-\varepsilon}(\delta, j)$, and so

$$U'(\delta, j) = U_0(\delta, j) \cup U_1(\delta, j) \cup V_0(\delta, j) \cup V_1(\delta, j).$$

Now put $p' = \langle A, n, U' \rangle$.

CLAIM 1.4. *If $\alpha \in U'(\beta, j)$ so $\alpha \in A^*$, then $\underline{\alpha} \in U_0(\underline{\alpha}(\beta), j)$.*

Indeed, if $\beta \in A_\varepsilon$, $j < n$, then $U'(\beta, j) \cap A^* = U_\varepsilon(\beta, j) \cup U_{1-\varepsilon}(\sigma^*(\beta), j)$.

CLAIM 1.5. *If $\langle \alpha, i \rangle \in U'(\beta, j)$ then $\underline{\alpha} \in U_0(\underline{\alpha}(\beta), j)$.*

PROOF OF CLAIM 1.5. Assume that $\beta \in A_\varepsilon$. If $\langle \alpha, i \rangle \in V_\varepsilon(\beta, j)$ then $\alpha \in U_\varepsilon(\alpha, i) \subset U_\varepsilon(\beta, j)$ and $U_\varepsilon(\beta, j) \subset U'(\beta, j)$. So we have $\alpha \in U'(\beta, j)$ which implies $\underline{\alpha} \in U_0(\underline{\alpha}(\beta), j)$ by Claim 1.4.

If $\langle \alpha, i \rangle \in W_{1-\varepsilon}(\beta, j)$ then for some $\langle \gamma, l \rangle \in D \times n$ we have $U_{1-\varepsilon}(\alpha, i) \subset U_{1-\varepsilon}(\gamma, l) \wedge U_\varepsilon(\gamma, l) \subset U_\varepsilon(\beta, j)$. Thus $\alpha \in U_\varepsilon(\beta, j) \subset U'(\beta, j)$, which implies $\underline{\alpha} \in U_0(\underline{\alpha}(\beta), j)$ by Claim 1.4. \square

CLAIM 1.6. $p' \in P$.

PROOF OF CLAIM 1.6. (P1) and (P2) clearly hold, so we need to check only (P3).

Assume on the contrary that (P3) fails for p' . Since $U'(\langle \nu, s \rangle, j) = \{\langle \nu, s \rangle\}$ by (4) for each $\langle \nu, s \rangle \in B$ and $j < n$, we can assume that some $\alpha < \beta \in A^*$ and $i < n$ witness that (P3) fails, i.e. $\beta \in U'(\alpha, i) \subset U'(\beta, 0)$. Then $\underline{\alpha}(\beta) \in U_0(\underline{\alpha}(\alpha), i) \subset U'(\underline{\alpha}(\beta), 0)$ by Claim 1.4. Since p_0 satisfies (P3) it follows that $\underline{\alpha}(\beta) \leq \underline{\alpha}(\alpha)$, and so $\alpha \in A_0 \setminus A_1$ and $\beta \in A_1 \setminus A_0$. Consider the element $u = \langle \alpha, i \rangle \in A \setminus A^*$. Then $u \in U'(\alpha, i)$ and so $u \in U'(\beta, 0)$ as well. By the definition of $U'(\beta, 0)$ this means that $\langle \alpha, i \rangle \in W_1(\beta, 0)$, that is, there is $\langle \gamma, l \rangle \in D \times n$ such that $U_0(\alpha, i) \subset U_0(\gamma, l)$ and $U_1(\gamma, l) \subset U_1(\beta, j)$. Thus

$$\underline{\alpha}(\beta) \in U_0(\alpha, i) \subset U_0(\gamma, l) \subset U_0(\underline{\alpha}(\beta), 0)$$

by Claim 1.4. Thus $\underline{\alpha}(\beta) \in U_0(\gamma, l) \subset U_0(\underline{\alpha}(\beta), 0)$ and so $\underline{\alpha}(\beta) \leq \gamma$ because p_0 satisfies (P3). But this is a contradiction because $\gamma \in D = A_0 \cap A_1$, $\underline{\alpha}(\beta) \in A_0 \setminus A_1$ and we assumed that $(A_0 \cap A_1) < (A_0 \setminus A_1)$. \square

CLAIM 1.7. $p' \leq p_0, p_1$.

PROOF OF CLAIM 1.7. Conditions (a) and (b) are clear.

To check (c) assume that $\alpha \in A_\varepsilon$ and $i \in n$. By (3),

$$\begin{aligned} U'(\alpha, i) \cap A_\varepsilon &= (U_\varepsilon(\alpha, i) \cup U_{1-\varepsilon}(\sigma^*(\alpha), i)) \cap A_\varepsilon \\ &= U_\varepsilon(\alpha, i) \cup (U_{1-\varepsilon}(\sigma^*(\alpha), i) \cap A_\varepsilon) = U_\varepsilon(\alpha, i) \end{aligned}$$

because $U_{1-\varepsilon}(\sigma^*(\alpha), i) = \sigma^*[U_\varepsilon(\alpha, i)]$.

To check (d1) assume that $\beta, \gamma \in A_\varepsilon$ and $j, k < n$ are such that $U'(\beta, j) \cap U'(\gamma, k) \neq \emptyset$. Fix $x \in U'(\beta, j) \cap U'(\gamma, k)$. Then

$$\underline{\alpha}(x) \in U_0(\underline{\alpha}(\beta), j) \cap U_0(\underline{\alpha}(\gamma), k)$$

by Claim 1.4 if $x = \alpha \in A^*$, and by Claim 1.5 if $x = \langle \alpha, i \rangle \in A \setminus A^*$.

If $\varepsilon = 0$ then $\underline{\alpha}(\beta) = \beta$ and $\underline{\alpha}(\gamma) = \gamma$, so $\underline{\alpha}(x) \in U_\varepsilon(\beta, j) \cap U_\varepsilon(\gamma, k)$.

If $\varepsilon = 1$ then $\underline{\alpha}(\beta) = \sigma^*(\beta)$ and $\underline{\alpha}(\gamma) = \sigma^*(\gamma)$, and so $\sigma^*(\underline{\alpha}(\alpha)) \in U_\varepsilon(\beta, j) \cap U_\varepsilon(\gamma, k)$.

Finally to check (d2) assume that $\beta, \gamma \in A_\varepsilon$ and $j, k < n$ are such that $U_\varepsilon(\beta, j) \subset U_\varepsilon(\gamma, k)$. Then clearly

$$U_{1-\varepsilon}(\sigma^*(\beta), j) = \sigma^*[U_\varepsilon(\beta, j)] \subset \sigma^*[U_\varepsilon(\gamma, k)] = U_{1-\varepsilon}(\sigma^*(\gamma), k),$$

moreover $V_\varepsilon(\beta, j) \subset V_\varepsilon(\gamma, k)$ by (1), and $W_\varepsilon(\beta, j) \subset W_\varepsilon(\gamma, k)$ by (2), and so $U'(\beta, j) \subset U'_\varepsilon(\gamma, k)$ by (3). \square

Now carry out the promised modification of U' to obtain U as follows. If $z \in A$ and $j < n$, let

$$U(z, j) = \begin{cases} U'(z, j) \cup U'(\xi_1, k), & \text{if } U_0(\xi_0, k) \subset U_0(z, j) \text{ so } \xi \in A_0, \\ U'(z, j), & \text{otherwise.} \end{cases}$$

Put

$$p = \langle A, n, U \rangle.$$

If $U_0(\xi_0, k) \subset U_0(z, j)$ then $U_1(\xi_1, k) \subset U_1(\sigma^*(z), j) \subset U'(z, j)$ and $W_1(\xi_1, k) \subset V_0(\xi_0, k) \subset U'(z, j)$. So

$$U(z, j) \setminus U'(z, j) \subset V_1(\xi_1, k). \quad (5)$$

Moreover

$$U(z, j) = \begin{cases} U'(z, j) \cup V_1(\xi_1, k), & \text{if } U_0(\xi_0, k) \subset U_0(z, j), \\ U'(z, j), & \text{otherwise.} \end{cases} \quad (6)$$

CLAIM 1.8. *If $\langle \alpha, i \rangle \in U(\beta, j)$ then $\underline{\alpha}(\alpha) \in U_0(\underline{\alpha}(\beta), j)$.*

Indeed, if $\langle \alpha, i \rangle \in U(\beta, j)$ then $\langle \alpha, i \rangle \in U'(\beta, j)$ or $\langle \alpha, i \rangle \in U'(\sigma^*(\beta), j)$, and now apply Claim 1.5.

CLAIM 1.9. $p \in P$.

PROOF OF CLAIM 1.9. (P1) and (P2) clearly hold, so we need to check (P3) only.

Assume on the contrary that (P3) fails for p . Since $U(\langle \nu, s \rangle, j) = \{\langle \nu, s \rangle\}$ for each $\langle \nu, s \rangle \in A \setminus A^*$ and $j < n$, we can assume that there are $\alpha < \beta \in A^*$ and $i < n$ witnessing that (P3) fails, i.e.

$$\beta \in U(\alpha, i) \subset U(\beta, 0). \quad (7)$$

Then $\underline{\alpha}(\beta) \in U_0(\underline{\alpha}(\alpha), i) \subset U(\underline{\alpha}(\beta), 0)$. But p_0 satisfies (P3) so $\underline{\alpha}(\beta) \leq \underline{\alpha}(\alpha)$, and so $\alpha \in A_0 \setminus A_1$ and $\beta \in A_1 \setminus A_0$.

Thus $U_0(\beta, j)$ is undefined, and so

$$U'(\beta, 0) = U(\beta, 0) \text{ and } U(\alpha, i) \setminus U'(\alpha, i) \subset A \setminus A^*$$

by (6). So (7) yields

$$\beta \in U'(\alpha, i) \subset U'(\beta, 0).$$

However this is a contradiction because p' satisfies (P3). □

CLAIM 1.10. $p \leq p_0, p_1$.

PROOF OF CLAIM 1.10. (a) and (b) are trivial. (c) also holds because $p' \leq p_\varepsilon$ and $(U(\alpha, i) \setminus U'(\alpha, i)) \cap A_\varepsilon = \emptyset$ by (5).

To check (d1) assume that $\beta, \gamma \in A_\varepsilon$ and $j, k < n$ are such that $U(\beta, j) \cap U(\gamma, k) \neq \emptyset$. Pick $x \in U(\beta, j) \cap U(\gamma, k)$. Then

$$\underline{\sigma}(x) \in U_0(\underline{\sigma}(\beta), j) \cap U_0(\underline{\sigma}(\gamma), k)$$

by Claim 1.4 if $x = \alpha \in A^*$, and by Claim 1.8 if $x = \langle \alpha, i \rangle \in A \setminus A^*$.

If $\varepsilon = 0$ then $\underline{\sigma}(\beta) = \beta$ and $\underline{\sigma}(\gamma) = \gamma$, so $\underline{\sigma}(x) \in U_\varepsilon(\beta, j) \cap U_\varepsilon(\gamma, k)$.

If $\varepsilon = 1$ then $\underline{\sigma}(\beta) = \sigma^*(\beta)$ and $\underline{\sigma}(\gamma) = \sigma^*(\gamma)$, and so $\sigma^*(\underline{\sigma}(x)) \in U_\varepsilon(\beta, j) \cap U_\varepsilon(\gamma, k)$.

Finally to check (d2) assume that $\beta, \gamma \in A_\varepsilon$ and $i, j < n$ are such that $U_\varepsilon(\beta, i) \subset U_\varepsilon(\gamma, j)$. Since $p' \leq p_\varepsilon$, we have $U'(\beta, i) \subset U'(\gamma, j)$. If $U(\beta, i) = U'(\beta, i)$, we are done. So we can assume that $U(\beta, i) = U'(\beta, i) \cup V(\xi_1, k)$. Then $\varepsilon = 0$ and $U_0(\xi_0, k) \subset U_0(\beta, i)$. But, if $\gamma \in A_0$, w.l.o.g. $\varepsilon = 0$ and then $U_0(\xi_0, k) \subset U_0(\gamma, j)$ and so $U(\gamma, j) = U'(\gamma, j) \cup V(\xi_1, k)$, and so $U(\beta, i) \subset U(\gamma, j)$. If $\gamma \notin A_0$ then $\varepsilon = 1$ and continue similarly. □

Since p satisfies (*), the Amalgamation lemma is proved. □

Using the Amalgamation lemma it is easy to complete the proof of the theorem.

By a standard Δ -system argument, any uncountable set of conditions contains two elements, p_0 and p_1 , which are twins. So, by Lemma 1.3, they have a common extension p . So P satisfies c.c.c.

If \mathcal{G} is a generic filter, for $\alpha < \omega_1$ and $i < \omega$ put

$$U(\alpha, i) = \bigcup \{U_p(\alpha, i) : p \in \mathcal{G}, \alpha \in A_p, i < n_p\},$$

and let $\mathcal{U}_\alpha = \{U(\alpha, i) : i < \omega\}$ be the base of the point α in $X = \langle \omega_1, \tau \rangle$.

By (P3), a countable family of open subsets is not a base of X . So $w(X) = \omega_1$.

Finally we show that X does not contain an uncountable subspace which has an irreducible base. Assume on the contrary that

$$r \Vdash \text{the subspace } \dot{Y} = \{\dot{y}_\xi : \xi < \omega_1\} \text{ has an irreducible base } \mathcal{B}, \\ \text{and } \{\dot{\mathcal{B}}_{\dot{y}_\xi} : \xi < \omega_1\} \text{ is an irreducible decomposition of } \dot{\mathcal{B}}.$$

We can assume that $r \Vdash \dot{y}_\xi \geq \check{\xi}$. For each $\xi < \omega_1$ pick a condition r_ξ and $k_\xi \in \omega$ such that

$$r_\xi \Vdash \text{“if } V \in \mathcal{B} \text{ with } \dot{y}_\xi \in V \subset U(\dot{y}_\xi, \check{k}_\xi) \text{ then } V \in \mathcal{B}_{\dot{y}_\xi}\text{”}. \quad (8)$$

For each $\xi < \omega_1$ pick a condition $p_\xi \leq r_\xi$, an ordinal $\alpha_\xi \geq \xi$, a name \dot{V}_ξ and a natural number $m_\xi < \omega$ such that $\alpha_\xi \in A_{p_\xi}$ and

$$p_\xi \Vdash \dot{y}_\xi = \check{\alpha}_\xi, \dot{V}_\xi \in \dot{\mathcal{B}}_{\alpha_\xi} \text{ and } U(\check{\alpha}_\xi, \check{m}_\xi) \subset \dot{V}_\xi \subset U(\check{\alpha}_\xi, \check{k}_\xi). \quad (9)$$

By a standard argument find $I \in [\omega_1]^{\omega_1}$ such that

- (i) $m_\xi = m$ and $k_\xi = k$ for each $\xi \in I$,
- (ii) the sequence $\{\alpha_\xi : \xi \in I\}$ is strictly increasing,
- (iii) the conditions $\{p_\xi : \xi \in I\}$ are pairwise twins,
- (iv) $\sigma_{\xi, \eta}(\alpha_\xi) = \alpha_\eta$ for $\{\xi, \eta\} \in [I]^2$, where $\sigma_{\xi, \eta}$ is the twin function.

Pick $\xi < \eta$ from I . By the Amalgamation Lemma there is a common extension p of p_ξ and p_η such that

$$\alpha_\xi \in U_p(\alpha_\eta, m) \wedge U_p(\alpha_\eta, k) \subset U_p(\alpha_\xi, k).$$

Then, by (d2),

$$p \Vdash \check{\alpha}_\xi \in U(\check{\alpha}_\eta, \check{m}) \wedge U(\check{\alpha}_\eta, \check{k}) \subset U(\check{\alpha}_\xi, \check{k}).$$

Then, by (9),

$$p \Vdash \dot{V}_\eta \in \mathcal{B}_{\alpha_\eta} \text{ and } \check{\alpha}_\xi \in U(\check{\alpha}_\eta, \check{m}) \subset \dot{V}_\eta \subset U(\check{\alpha}_\eta, \check{k}) \subset U(\check{\alpha}_\xi, \check{k}),$$

which contradicts (8).

This completes the proof of Theorem 1.2. □

References

- [1] I. JUHÁSZ, L. SOUKUP and Z. SZENTMIKLÓSSY, What makes a space have large weight?, *Topology Appl.*, **57** (1994), 271–285.