

The Journal of Symbolic Logic

<http://journals.cambridge.org/JSL>

Additional services for *The Journal of Symbolic Logic*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Adding linear orders

Saharon Shelah and Pierre Simon

The Journal of Symbolic Logic / Volume 77 / Issue 02 / June 2012, pp 717 - 725

DOI: 10.2178/jsl/1333566647, Published online: 12 March 2014

Link to this article: http://journals.cambridge.org/abstract_S0022481200000839

How to cite this article:

Saharon Shelah and Pierre Simon (2012). Adding linear orders . The Journal of Symbolic Logic, 77, pp 717-725 doi:10.2178/jsl/1333566647

Request Permissions : [Click here](#)

ADDING LINEAR ORDERS

SAHARON SHELAH[†] AND PIERRE SIMON

Abstract. We address the following question: Can we expand an *NIP* theory by adding a linear order such that the expansion is still *NIP*? Easily, if $\text{acl}(A)=A$ for all A , then this is true. Otherwise, we give counterexamples. More precisely, there is a totally categorical theory for which every expansion by a linear order has *IP*. There is also an ω -stable *NDOP* theory for which every expansion by a linear order interprets pseudofinite arithmetic.

A well known open question is whether every unstable *NIP* theory interprets an infinite linear order. We are concerned here with a question somewhat in the same spirit but going in a different direction: Can we expand an *NIP* theory by adding a linear order on the whole universe so that the resulting theory is still *NIP*? We give a negative answer in two strong forms:

- (1) There is an ω -stable *NDOP* theory of depth 2 for which every expansion by a linear order interprets pseudofinite arithmetic (see Section 2.2).
- (2) There is a totally categorical theory for which every expansion by a linear order has *IP*.

In the first section, we mention a few positive statements that are true (and easy): if M is *NIP* and $\text{acl}(A) = A$ for all $A \subset M$, then M can be linearly ordered so as to stay *NIP*. Also if M is ω -categorical, then M can be linearly ordered so as to stay ω -categorical (a well known fact) so we cannot expect to get the strong conclusion of (1) with an ω -categorical theory.

Let us also note that, as proved in [2] (using results from [1]), adding a predicate for a small dense indiscernible sequence preserves *NIP*. In particular any unstable *NIP* theory T has an *NIP* expansion which defines an infinite linear order.

As far as we know, the question we address was first asked by Artem Chernikov. It came up again in discussions with Udi Hrushovski, which led to this work. We would like to thank him for that and for helping the second author understand the results of the first author.

§1. The easy case. We fix a one-sorted relational language \mathcal{L} not containing the symbol $<$, let $\mathcal{L}_{<}$ be the language with a unique binary relation $<$ and let $\mathcal{L}' = \mathcal{L} \cup \mathcal{L}_{<}$. Let T be a complete \mathcal{L} -theory that eliminates quantifiers. Let T'

Received May 6, 2011.

[†]The first author would like to thank the Israel Science Foundation for partial support of this research (Grant no. 710/07). Publication 979 on Shelah's list.

be the \mathcal{L}' -theory generated by T and axioms stating that $<$ defines a dense linear order with no end-points.

We show in this section that if T eliminates \exists^∞ , then T' has a model-companion. Apart maybe from Proposition 1.4, everything here is well known. We follow the exposition of [6] which contains exactly what we need. The ω -categorical case was already observed by Schmerl in [5].

Let M be any structure, and $A \subset M$ a finite subset. We say that a formula $\phi(x_1, \dots, x_n)$ with parameters in A is *large* if it has a solution a_1, \dots, a_n such that: for all i , $a_i \notin \text{acl}(A)$ and for all $i \neq j$, $a_i \neq a_j$. If $\phi(x_1, \dots, x_n; y)$ is a formula, then using elimination of \exists^∞ , it can be checked that the set of d 's such that $\phi(x_1, \dots, x_n; d)$ is large is definable (Fact 2 of [6]).

PROPOSITION 1.1. *Let T be any \mathcal{L} -theory that eliminates quantifiers and \exists^∞ . Then T' admits a model-companion \tilde{T} axiomatized by T along with the statements saying that for every large \mathcal{L} -formula $\phi(x_1, \dots, x_n)$ and quantifier-free large $\mathcal{L}_<$ -formula $\theta(x_1, \dots, x_n)$, the conjunction $\phi(x_1, \dots, x_n) \wedge \theta(x_1, \dots, x_n)$ has a solution.*

Furthermore, in \tilde{T} , the type of an algebraically closed set is given by the union of its \mathcal{L} -type and its $\mathcal{L}_<$ -quantifier-free-type.

PROOF. The first statement is a special case of Lemma 8 of [6]. The proof is straightforward: any model of T' embeds into a model of \tilde{T} by iteratively adding solutions to formulas. Existential closedness is clear.

The second statement is by easy back-and-forth. \dashv

COROLLARY 1.2. *Let $M \models T$ be an \mathcal{L} -structure, and assume that algebraic closure is trivial in M : $\text{acl}(A) = A$ for all $A \subset M$. Assume that M admits elimination of quantifiers in \mathcal{L} . Then there is an expansion T' of T to $\mathcal{L}' = \mathcal{L} \cup \{<\}$ such that $<$ defines a linear order and T' has elimination of quantifiers in \mathcal{L}' .*

PROOF. Take \tilde{T} as above. As $\text{acl}(A) = A$ for all $A \subset M$, the type of any set A is given by its \mathcal{L} -type and its quantifier-free $\mathcal{L}_<$ -type. As T eliminates quantifiers in \mathcal{L} , \tilde{T} eliminates quantifiers in \mathcal{L}' . \dashv

- CONCLUSION 1.3.**
1. *If T is ω -categorical, then T admits an ω -categorical expansion to \mathcal{L}' in which $<$ defines a linear order.*
 2. *If T has trivial algebraic closure and is NIP, then it admits an NIP expansion to \mathcal{L}' in which $<$ defines a linear order.*

PROOF. The second point follows from the Corollary.

For the first point, notice first that any ω -categorical theory eliminates \exists^∞ . More precisely, for finite A , $\text{acl}(A)$ is finite. So for a given integer n , the type of an n -tuple $\bar{a} = (a_1, \dots, a_n)$ in a model of \tilde{T} is given by its \mathcal{L} -type and the ordering of $\text{acl}(\bar{a})$. We see that there are only finitely many possibilities. Therefore \tilde{T} is ω -categorical. \dashv

We end this section with a small observation.

PROPOSITION 1.4. *Assume T eliminates \exists^∞ . Let $M \models T$. If all infinite definable sets of M has the same cardinality λ , then M admits an expansion to a model M' of \tilde{T} as defined above.*

PROOF. Fix a λ -saturated dense linear order $(\Omega, <_\Omega)$ without end points. We will build an injection $f : M \rightarrow \Omega$. Fix an enumeration $(\bar{a}_\alpha : \alpha < \lambda)$ of all pairs $\bar{a}_\alpha = (\phi_\alpha(x_1, \dots, x_n), \theta_\alpha(x_1, \dots, x_n))$ where $\phi_\alpha(\bar{x}) \in \mathcal{L}(M)$ is large and $\theta_\alpha(\bar{x})$ is

an $\mathcal{L}_<$ -quantifier-free formula with parameters in M . We build f by induction on $\alpha < \lambda$ so f will be defined as an increasing union $f = \cup_{\alpha < \lambda} f_\alpha$, where each f_α has domain a subset of M of size $|\alpha|$.

At limit stages, set $f_\alpha = \cup_{\beta < \alpha} f_\beta$.

Assume f_α has been defined. Let A be the set of parameters of θ_α . If needed, start by increasing f_α to an injection f' defined on $Dom(f_\alpha) \cup A$ by defining it on A in an arbitrary way. Let θ' be the formula built from θ_α changing every parameter by its image by f' . If θ' is not large (in particular if it is inconsistent), we let $f_{\alpha+1} = f'$.

Otherwise, we can find a tuple $\bar{a} = (a_0, \dots, a_{n-1})$ in M such that $M \models \phi(\bar{a})$ and no a_k is in the domain of f' . Pick any $\bar{c} = (c_0, \dots, c_{n-1})$ in Ω , such that $\Omega \models \theta'(\bar{c})$ and no c_k is in the range of f' . We define $f_{\alpha+1}$ as $f' \cup \{(a_k, c_k) : k < n\}$.

Once f is defined, we expand M to \mathcal{L}' by letting $a < b$ if and only if $f(a) <_\Omega f(b)$. By construction the resulting structure is a model of \tilde{T} . \dashv

§2. Counterexamples.

2.1. \mathbb{F}_p vector space. We prove the following result.

THEOREM 2.1. *If T defines an infinite dimensional vector space over some prime field \mathbb{F}_p , and T is linearly ordered by $<$, then T has IP.*

First we need to recall the following Ramsey-type result. See for example [3], Section 2.4, Theorem 9.

Here F is a fixed finite field.

THEOREM 2.2. *For all $r, t, k \geq 1$, there is some n such that if the t -dimensional affine subspaces of F^n are r -colored, there exists a k -dimensional affine subspace all of whose t -dimensional affine subspaces have the same color.*

Now we prove Theorem 2.1.

PROOF. We identify $F = \mathbb{F}_p$ with the set $\{0, 1, \dots, p-1\}$ and order F by setting $0 < 1 < \dots < p-1$.

Without loss, T is just an ordered vector space over F . Let $M \models T$ and pick an infinite free family $\langle a_i : i < \omega \rangle$. Let k be any integer and set $t = 1, r = p!$. Finally, let n be given by Theorem 2.2 for those values of r, t, k . We consider the vector space A spanned by (a_1, \dots, a_n) and identify it with F^n using (a_1, \dots, a_n) as canonical base. Let \triangleleft be the lexicographic order on A (identified with F^n).

Let $L \subset A$ be an affine line in A , $L = \{d_0, \dots, d_{p-1}\}$ with $d_0 \triangleleft d_1 \triangleleft \dots \triangleleft d_{p-1}$. We assign to L a 'color' $c(L)$ from the set S of permutations of F in the following way: $c(L)$ is the unique permutation π such that

$$d_{\pi(0)} < d_{\pi(1)} < \dots < d_{\pi(p-1)},$$

where $<$ is the given order on M .

By Theorem 2.2, we can find a k -dimensional affine subspace W of A all of whose 1-dimensional affine subspaces have the same color π . So we can write $W = d_* + W_0$ where W_0 is a linear subspace. Let (b^1, \dots, b^k) be a basis of W_0 such that for each $l \leq k$, b^l is \triangleleft -minimal in $W_0 \setminus Vect(b^1, \dots, b^{l-1})$. For each $l \leq k$, let ω_l be the least i such that the i 'th coordinate of b^l (in the basis (a_1, \dots, a_n)) is non zero. By choice of b^l we have

$$\omega_1 > \omega_2 > \dots > \omega_k.$$

It follows that for each choice of $\mathfrak{s} = (s_1, \dots, s_k) \in F^k$, there is a unique element $d_{\mathfrak{s}}$ of W such that for every i , the ω_i 'th coordinate of $d_{\mathfrak{s}}$ is equal to s_i . For $I \subseteq \{1, \dots, k\}$, let d_I be $d_{\mathfrak{s}}$ for $\mathfrak{s} = (\varepsilon_{1,I}, \dots, \varepsilon_{k,I})$ where $\varepsilon_{i,I} = 1$ if $i \in I$ and 0 otherwise.

We let $\phi_{\pi}(x, y)$ be the formula:

$$\phi_{\pi}(x, y) = \bigwedge_{i=0}^{p-2} x + \pi(i).y < x + \pi(i+1).y.$$

Let $I \subseteq \{1, \dots, k\}$ and $l \leq k$. Consider the line $L = d_I + \text{Vect}(b^l)$. Enumerating L in \triangleleft increasing order gives

$$d_I - \varepsilon_{l,I}b^l \triangleleft d_I - \varepsilon_{l,I}b^l + b^l \triangleleft \dots \triangleleft d_I - \varepsilon_{l,I}b^l + (p-1).b^l.$$

As L has color π we see that $\phi_{\pi}(d_I, b^l)$ holds if and only if $\varepsilon_{l,I} = 0$. This proves that the formula $\phi_{\pi}(x, y)$ has independence rank at least k . As k was arbitrary, and there are only finitely many possibilities for π , there is at least one value of π for which $\phi_{\pi}(x, y)$ has IP. \dashv

Remark 2.3. In the case $p = 2$, the same proof works if instead of assuming that $<$ defines a linear order, we only assume that it defines a tournament (*i.e.*, for all $x \neq y \in M$, exactly one of $x < y$ and $y < x$ holds).

2.2. Interpreting arithmetic. By pseudofinite arithmetic, we mean the (incomplete) theory T_{arith} consisting of formulas true in almost all structures $(\{0, \dots, n\}; +, \times)$. A model interprets pseudofinite arithmetic if it has an elementary extension that interprets a model of T_{arith} . Note that the theory T_{arith} has IP as can be seen by considering the formula $x|y$ (x divides y), so it follows that any structure interpreting pseudofinite arithmetic also has IP.

THEOREM 2.4. *There is an ω -stable theory T , NDOP of depth 2, in a language \mathcal{L} such that: for every model $M \models T$ and every expansion M' of M to $\mathcal{L}' = \mathcal{L} \cup \{<\}$ in which $<$ defines a linear order, M' interprets pseudofinite arithmetic.*

We take as language $\mathcal{L} = \{E, S, R\}$ where E and S are binary predicates and R is quaternary. In the structure we will build, E will define an equivalence relation and each E -class will be made by (S, R) into a regular colored graph as defined later.

We will define T by constructing its prime model M_0 built as a disjoint union of finite regular graphs. Each of those graphs will be exactly one E -class. The n 'th graph will contain no cycle of length $\leq n$. We will choose the finite graphs in such a way that no matter what order is put on them, the n 'th graph interprets $(\{1, \dots, n\}, +, \times)$, by an explicit formula not depending on n . The condition about cycles will ensure that the limit theory of those graphs is ω -stable.

The first observation is that, for every n , there is a structure $(\{1, \dots, N\}, <_1, <_2)$ where both $<_1$ and $<_2$ are linear orders (call this a *bi-order*) that interprets $(\{1, \dots, n\}, +, \times)$. Furthermore, the formulas involved in the interpretation do not depend on n . This is an easy exercise whose solution is given in the appendix. As a consequence, the problem is reduced to that of interpreting two linear orders.

We will work with colored graphs. For us, a colored graph is a structure $(G; S, R)$ where $S(x, y)$ is symmetric anti-reflexive and $R(x_0, y_0, x_1, y_1)$ is a quaternary relation which defines an equivalence relation on pairs $(x, y) \in S$. It should be thought of as saying that $\{x_0, y_0\}$ and $\{x_1, y_1\}$ are two edges in the graph of the same color. We will consider only regular colored graph, namely such that each vertex belongs to exactly one edge of each color. For simplicity, we will introduce the imaginary sort C of colors defined to be the quotient of $\{(x, y) \in G^2 : (x, y) \in S\}$ by the equivalence relation $(x_0, y_0) \sim (x_1, y_1) \iff R(x_0, y_0, x_1, y_1)$. If $k \in C$ is a color and $x \in G$ is a vertex, by the k -neighbor of x , we mean the unique $y \in G$ such that $\{x, y\}$ is an edge of color k .

Denote by \mathcal{L}_{cg} the language $\{S, R\}$ of colored graph. We will use \mathcal{L}'_{cg} to denote the expanded language $\{S, R, <\}$ where $<$ is a binary predicate. Also let $\Phi(x, y, u, v)$ be the \mathcal{L}'_{cg} formula saying that $\{u, v\}$ is an edge and if k is its color, then the k -neighbor of x is $<$ -less than the k -neighbor of y .

In the rest of this subsection, we prove the following proposition.

PROPOSITION 2.5. *Let $N, c \geq 3$ be integers, let also $<_1, <_2$ be two linear orders on $N = \{0, \dots, N-1\}$. Then for every even integer n big enough, there is a finite regular colored graph G_n on n vertices such that for every expansion of G_n to \mathcal{L}'_{cg} making $<$ into a linear order, there are $a, b, u, v \in G_n$ such that the structure $([a, b]; <, \Phi(x, y, u, v))$ is isomorphic to $(N; <_1, <_2)$. Furthermore, G_n has no cycle of length $\leq c$.*

Here $[a, b]$ denotes the interval with end points a and b in the sense of $<$.

We will build the graph G_n by a random procedure and show that with positive probability, we obtain what we want. Actually, we will start by building a colored multigraph $G(\sigma)$ and then modify it to make into an actual regular graph $G'(\sigma)$. By a colored multigraph, we mean a colored graph in which there can be two or more edges (of different colors) between two given vertices.

Let n be even and big enough (we will see during the construction what big enough means). Fix some $1 - \frac{1}{3c} < \alpha < 1$ and let $d = \lfloor n^{1-\alpha} \rfloor$.

Let \mathfrak{S}_n denote the symmetric group on n elements. Our space of events is $\Omega = \mathfrak{S}_n^d$ equipped with the uniform probability law. Let $\sigma = (\sigma_i : i < d)$ be an element of Ω . We define a colored multigraph $G(\sigma)$ as follows: The vertex set of $G(\sigma)$ is $n = \{0, \dots, n-1\}$. The set of colors is $C = \{0, \dots, d-1\}$. For $k < d$, we draw an edge of color k between vertices a and b if and only if for some $l < n/2$, we have $\{\sigma_k(2l), \sigma_k(2l+1)\} = \{a, b\}$.

A cycle of length $r \geq 2$ is a sequence (a_0, \dots, a_{r-1}) of distinct vertices and a sequence (e_0, \dots, e_{r-1}) of distinct edges such that for each $i < r$, e_i is an edge between a_i and a_{i+1} (addition is modulo r). In particular, a cycle of length 2 in $G(\sigma)$ is given by two vertices and two different edges linking them. A cycle is said to be *small* if it is of length $\leq c$.

If $k \in C$ is a color, we define \prec_k by $x \prec_k y$ if and only if the k -neighbor of x is $<$ -less than the k -neighbor of y . We will show that each of the following events occurs with probability converging to 1 as n tends to $+\infty$:

1. The number of small cycles in $G(\sigma)$ is less than d^{2c} , and we can obtain a regular graph $G'(\sigma)$ with no small cycles by changing at most $2d^{2c}$ edges of $G(\sigma)$,

2. For every ordering $<$ of the vertices of $G(\sigma)$, we can find at least d^{3c} values of $(a, b, k) \in G^2 \times C$ such that $([a, b]; <, <_k)$ is isomorphic to $(N; <_1, <_2)$.

2.2.1. Removing small cycles. We perform some surgery to remove small cycles from the multigraph $G(\sigma)$ and obtain a regular graph $G'(\sigma)$ with no small cycles.

We compute the expectancy of the number of small cycles in $G(\sigma)$. Let $2 \leq s \leq c$. Let $H = (V_H, E_H)$ be the graph consisting of a unique cycle of length s . Fix some $f : V_H \rightarrow \{0, \dots, n-1\}$ injective and some $g : E_H \rightarrow \{0, \dots, d-1\}$. Assume that for any two edges $e, e' \in E_H$ having a vertex in common, $g(e) \neq g(e')$. Call the pair (f, g) a subgraph of $G(\sigma)$ if for every edge $e = \{a, b\} \in E_H$, $\{f(a), f(b)\}$ is an edge of $G(\sigma)$ and has color $g(e)$. For a given edge e (and remembering that the pair (f, g) is fixed), the probability p_0 that this occurs is

$$p_0 = \frac{1}{n-1}.$$

If g is injective, then all those events are independent, so the probability p that (f, g) is a subgraph satisfies

$$p \sim \left(\frac{1}{n}\right)^s.$$

If g is not injective, the events are not independent, but nevertheless, looking color by color, the same estimate can easily be seen to be true.

The number of such pairs (f, g) is less than $n^s d^s$. So the expected value of the number of cycles of length s is asymptotically at most

$$n^s d^s \left(\frac{1}{n}\right)^s = d^s.$$

Let X denote the number of small cycles in $G(\sigma)$. Summing over all $s \leq c$, we see that $E(X) = O(1) \cdot d^c$. By the first moment method

$$\text{Prob}(X \geq d^{2c}) \leq O(1) \cdot d^{-c}.$$

In particular, with probability converging to 1 at n tends to $+\infty$, the number of small cycles in $G(\sigma)$ is less than d^{2c} . Consider a σ that has that property.

We now modify the graph $G(\sigma)$ so as to remove all small cycles. We show that this can always be done by changing at most $2d^{2c}$ edges. Working by induction on the number of small cycles, we have to show that if G is a $(d$ -regular colored) graph with less than d^{2c} small cycles, then one can move 2 edges of it so as to decrease the number of small cycles.

Let G be such a graph and take C a cycle of size $s \leq c$. Pick any edge $\{a, b\}$ in C . Assume it is drawn with color k . We want to take an edge $\{a', b'\}$ drawn with color k , erase both those edges, and draw instead, with color k , the edges $\{a, a'\}$ and $\{b, b'\}$. We have to choose $\{a', b'\}$ so that no new small cycle is introduced. It is easy to check that this will happen if the following two properties are satisfied:

- the graph distance between a and a' is at least $c + 2$,
- the edge $\{a', b'\}$ does not belong to a small cycle.

Note that the first condition implies that the distance between b and b' is at least c . We see that the number of edges $\{a', b'\}$ of color k which fail to satisfy those properties is at most $O(1) \cdot (d^{c+2} + d^{2c}) < n/2$, so we can find a', b' as required.

Iterating this procedure starting with our original graph $G(\sigma)$, we obtain a graph $G'(\sigma)$ with no small cycle and such that all but at most $2d^{2c}$ of the edges of $G(\sigma)$ are still edges in $G'(\sigma)$ of the same color.

2.2.2. Interpreting bi-orders. Let $(N; <_1, <_2)$ be a given bi-order. We now prove that we can interpret it with high probability in $G(\sigma)$ (uniformly in N) in many different ways. This will automatically imply that we can also interpret it in $G'(\sigma)$. Let $<$ be any order on $\{0, \dots, n-1\}$ (the set of vertices of $G(\sigma)$), without loss, the usual order. Given $a < n$ and $k \in C$ a color, define the event $\mathcal{E}_{a,k}$ as

$$([a, a+N-1]; <, <_k) \text{ is isomorphic to } (N; <_1, <_2).$$

Fix a value of k . Let $D \subset \{0, \dots, n-1\}$ be a subset of size $< n/4N$ closed under taking the k -neighbor. Assume we know the values of $\sigma_k(x)$ for every $x \in D$ and the values of $\sigma_l(x)$ for $l < k$ and every x . Take some $a < n$ such that $D \cap \{a, a+1, \dots, a+N-1\} = \emptyset$. We look at the values of the k -neighbors of the points $\{a, a+1, \dots, a+N-1\}$. With probability $> 1/2$ those neighbors are disjoint from $\{a, a+1, \dots, a+N-1\}$ so we are considering $2N$ distinct points. Consider the bi-order $([a, a+N-1]; <, <_k)$. Clearly, all $N!$ bi-order have the same probability of occurring. So, knowing σ_k restricted to D and σ_l for $l < k$, the event $\mathcal{E}_{a,k}$ holds with probability at least $\frac{1}{2N!}$.

Now, assume we know the values of σ_l for $l < k$. Take $D_0 = \emptyset$ and $a_0 = 0$. With probability at least $\frac{1}{2N!}$, the event $\mathcal{E}_{a_0,k}$ holds. Next we let D_1 contain $\{a_0, a_0+1, \dots, a_0+N-1\}$ and all k -neighbors of those points. We take some a_1 as in the previous paragraph. Again with probability at least $\frac{1}{2N!}$, $\mathcal{E}_{a_1,k}$ holds. We can iterate this at least $\frac{n}{8N^2}$ times (as D increases by at most $2N$ each time). We do this for each color, one color after the other. Let Y be the number of events $\mathcal{E}_{a,k}$ that hold. Then for every x ,

$$\text{Prob}(Y \leq x) \leq \text{Prob}(B \leq x)$$

where B is a random variable whose law is a binomial distribution with parameters $\frac{dn}{8N^2}$, $\frac{1}{2N!}$. (Recall that the binomial distribution with parameters n, p is defined as the distribution of the sum of n independent random variables being equal to 1 with probability p and to 0 with probability $1-p$.)

We now use the following fact about binomial distributions. It is a special case of Hoeffding's inequality (or of Chernoff's bound). See for example [4].

LEMMA 2.6. *Let B be a random variable whose law is a binomial distribution with parameters n, p . Then for any x we have:*

$$\text{Prob}(B \leq x) \leq \exp\left(-2\frac{(np-x)^2}{n}\right).$$

We conclude that the probability that at least d^{3c} of the events $\mathcal{E}_{a,k}$ succeed is at least

$$1 - \exp\left(-16N^2\frac{\left(\frac{dn}{16N^2N!} - d^{3c}\right)^2}{nd}\right) = 1 - \exp(-n^{2-\alpha} \cdot O(1)).$$

As there are $n! \leq \exp(n \ln n)$ different orders on $\{0, \dots, n-1\}$, we see that with probability converging to 1 as n tends to $+\infty$, for every ordering $<$ of the vertices,

there are at least d^{3c} values of $(a, k) \in n \times d$ for which

$$([a, a + N - 1]; <, \triangleleft_k) \text{ is isomorphic to } (N; <_1, <_2).$$

In particular, if furthermore $G(\sigma)$ has less than d^{2c} small cycles, there is a choice of (a, k) such that no edge of color k having an end point in $\{a, \dots, a + N - 1\}$ is changed during the construction of $G'(\sigma)$. Therefore with this choice of (a, k) , we obtain an interpretation of $(N; <_1, <_2)$ in $G'(\sigma)$ as

$$([a, a + N - 1], <, \Phi(x, y, u, v))$$

where $\{u, v\}$ is any edge of color k .

This ends the proof of Proposition 2.5.

2.3. The full structure. We now conclude with the proof of Theorem 2.4.

We recall that we set $\mathcal{L} = \{E, S, R\}$ where E is a binary relation. Fix an $3 \leq n < \omega$. Let $(N; <_1, <_2)$ be the structure P_n given by Proposition 2.7 below. Let also $(G_n; S, R)$ be the colored graph given by Proposition 2.5 with $c = n$. We define the \mathcal{L} -structure M_0 as follows. The reduct to $\{S, R\}$ is just the disjoint union of the graphs $(G_n; S, R)$. The predicate E is interpreted as an equivalence relation, such that two points $x, y \in M_0$ are E -equivalent if and only if they come from the same G_n .

What does $Th(M_0)$ look like? Let M be elementary equivalent to M_0 . Then in M , E defines an equivalence relation which has ω finite classes. The substructure of M formed by the union of those finite classes is isomorphic to M_0 . An infinite class of M , equipped with (S, R) is a regular colored graph with no cycles and infinite degree. So as a graph, it is a union of trees. Finally, there are no S -relations between points in different E -classes.

We see that to every E -class are associated two regular types: one for a new connected component and one for a new color. The type of a new E -class is itself a regular type. A model is entirely determined up to isomorphism by the number of infinite E -classes and for each such class, the number of connected components and the number of colors. So the resulting theory is ω -stable, *NDOP* of depth 2.

Finally, if $<$ is an expansion of M_0 by a linear order, then in the k 'th equivalence class, we can interpret $(\{0, \dots, k\}; +, \times)$, uniformly in k . In particular the expansion interprets pseudofinite arithmetic.

Appendix: interpreting arithmetic from two linear orders. The following is certainly well known, but we include it for completeness.

PROPOSITION 2.7. *There are formulas $\phi(x, y, z; \bar{t})$, $\psi(x, y, z; \bar{t})$ in the language $\mathcal{L} = \{<_1, <_2\}$ such that for each $k < \omega$, there is a finite \mathcal{L} -structure P_k satisfying:*

- $<_1$ and $<_2$ define linear orders on P_k ,
- there are a_0, a_1, \bar{a} in P_k such that the structure

$$([a_0, a_1]_2; \phi(x, y, z; \bar{a}), \psi(x, y, z; \bar{a}))$$

is isomorphic to $(\{0, \dots, k - 1\}, +, \times)$.

(where $[a, b]_2$ denotes the interval $a \leq_2 x \leq_2 b$).

PROOF. Let $k < \omega$ and set $n = 10k^2$. Let P_k have universe $\{0, \dots, n - 1\}$ and let $<_1$ be the usual order on that set. For $i \in \{0, \dots, k - 1\}$, let $b_i = 7k(i + 1)$. Let $a_1 = b_{k-1}$. The points of P_k greater than b_{k-1} will be called *delimiters*.

Now we construct $<_2$ in the following way: an initial segment is $b_0 <_2 b_1 <_2 \dots <_2 b_{k-1}$. Those points will correspond to $\{0, \dots, k-1\}$ in the interpretation. For $l < n$, $r < k$ we say that l codes for r if $b_{r-1} < l < b_r$ (setting $b_{-1} = -1$). Now we encode the graph of addition. Let $(A_i)_{i < N}$ be an enumeration of all triples $(r, s, t) \in \{0, \dots, k-1\}^3$ such that $r + s = t$. Build a map $f : N \rightarrow P_k^4$ such that for each $i < N$, if $f(i) = (c_r, c_s, c_t, d)$ then c_r, c_s and c_t code for r, s, t respectively and $d = a_1 + i + 1$ is a delimiter. Also impose that no point appears in two tuples $f(i), f(i')$ for $i \neq i'$. Now set $<_2$ such that $c_r <_2 c_s <_2 c_t <_2 d$ each time $f(i) = (c_r, c_s, c_t, d)$, and those four points form an interval of $<_2$. Place those intervals one after another in any order. Let a_2 be the last delimiter placed. Next do the same for the graph of multiplication, and let a_3 be the last delimiter placed there. Finally place the elements that are left in any order after a_3 .

Now it is easy to check that one can interpret $(\{0, \dots, k-1\}; +, \times)$ in this structure, with parameters a_1, a_2, a_3 , and that the formulas involved do not depend either on the choices made nor on k . \dashv

REFERENCES

- [1] JOHN BALDWIN and MICHAEL BENEDIKT, *Stability theory, permutations of indiscernibles, and embedded finite models*, *Transactions of the American Mathematical Society*, vol. 352 (2000), no. 11, pp. 4937–4969.
- [2] ARTEM CHERNIKOV and PIERRE SIMON, *Externally definable sets and dependent pairs*, submitted, Modnet preprint 265.
- [3] R. GRAHAM, B. ROTHSCHILD, and J.H. SPENCER, *Ramsey theory*, 2nd ed., New York: John Wiley and Sons, 1990.
- [4] WASSILY Hoeffding, *Probability inequalities for sums of bounded random variables*, *Journal of the American Statistical Association*, vol. 58 (1963), no. 301, pp. 13–30.
- [5] JAMES H. SCHMERL, *Decidability and \aleph_0 -categoricity of theories of partially ordered sets*, this JOURNAL, vol. 45 (1980), no. 3, pp. 585–611.
- [6] AKITO Tsuboi, *Random amalgamation of simple theories*, *Mathematical Logic Quarterly*, vol. 47 (2001), no. 1, pp. 45–50.

EINSTEIN INSTITUTE OF MATHEMATICS
EDMOND J. SAFRA CAMPUS, GIVAT RAM
THE HEBREW UNIVERSITY OF JERUSALEM
JERUSALEM, 91904
ISRAEL

E-mail: shelah@math.huji.ac.il
URL: <http://shelah.logic.at>

ENS
DÉPARTEMENT DE MATHÉMATIQUES ET APPLICATIONS
45, RUE D'ULM
75005 PARIS, FRANCE

E-mail: pierre.simon.05@normalesup.org
URL: <http://www.normalesup.org/~simon/>