

Set theory without choice: not everything on cofinality is possible

Saharon Shelah^{1,2,*}

¹ Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel**

² Department of Mathematics, Rutgers University, New Brunswick, NJ, USA

Received May 6, 1993 / Revised December 11, 1995

Abstract. We prove in ZF+DC, e.g. that: if $\mu = |\mathcal{H}(\mu)|$ and $\mu > \text{cf}(\mu) > \aleph_0$ then μ^+ is regular but non measurable. This is in contrast with the results on measurability for $\mu = \aleph_\omega$ due to Apter and Magidor [ApMg].

Annotated content

0 Introduction

[In addition to presenting the results and history, we gave some basic definitions and notation.]

1 Exact upper bound

[We define some variants of least upper bound (*lub*, *eub*) in $(^{A^*}\text{Ord}, <_D)$ for D a filter on A^* . We consider $<_D$ -increasing sequence indexed by ordinals δ or indexed by sufficiently directed partial orders I , of members of $(^{A^*}\text{Ord}$ or of members of $(^{A^*}\text{Ord}/D$ (and cases in the middle). We give sufficient conditions for existence involving large cofinality (of δ or I) and some amount of choice. Mostly we look at what the *ZFC* proof gives without choice. Note in particular 1.8, which assumes only *DC* (*ZF* is not mentioned of course), the filter is \aleph_1 -complete and cofinality of δ large and we find an *eub* only after extending the filter.]

2 *hpp*

[We look at various ways to measure the size of the set $\prod_{a \in A^*} f(a)/D$, like supremum length of $<_D$ -increasing sequence in $\prod_{a \in A^*} f(a)$ (called *ehpp* _{D}), or

* Research supported by “The Israel Science Foundation” administered by The Israel Academy of Sciences and Humanities. Publication no 497, done 11/92–1/93.

** Address for correspondence

Mathematics Subject Classification: 03E05, 03E25

in $\prod_{a \in A^*} f(a)/D$ (called hpp_D), or we can demand them to be cofinal (getting epp or pp); when we let D vary on Γ we write $ehpp_\Gamma$ etc. So existence of $<_D$ – eub give downward results on pp .]

3 Nice family of filters

[In this paper we can say little on products of countably many; we can say something when we deal with \aleph_1 -complete filters. So as in [Sh-g], V, we deal with family E of filters on A^* which is nice. Hence suitable ranks from function f from A^* to ordinal and filter $D \in E$ are well defined (i.e. the values are ordinals not infinity). The basic properties of those ranks are done here.

We then define some measures for the size of $\prod_{a \in A^*} f(a)/D$ (i.e. Tw , Ts , T), looking at subsets of $\prod_{a \in A^*} f(a)$ or of $\prod_{a \in A^*} f(a)/D$ which are pairwise \neq_D . In conclusion 3.12 we, under reasonable assumption, prove that some such measures and $\sup\{rk_D^2(f) : D \in E\}$ are equal. In 3.14 we have a parallel of 1.8: sufficient condition for the existence of eub when we allow to increase the filter. We end defining normal filters and a generalization.

The basic point is that for every $f \in {}^{(A^*)}\text{Ord}$ (if E is nice) for some $D \in E$ we have $rk_D^2(f) = rk_D^3(f)$ and in this case the rank determine f/D , and order on rank equal order among such f ; so we can represent $\prod_{a \in A^*} f^*(a)/D$ as the union of $\leq |E|$ well ordered sets.]

4 Investigating strong limit singular

[We deal with $\otimes_{\alpha,R}$, which means that we can regard $R \cap \alpha$ as a substitute of the family of regulars (not just individually) i.e. we can find $\langle e(i) : i < \alpha \rangle$, $e(i)$ is an unbounded subset of i of order type which belongs to R . We give the basic properties in 4.2, then move up from μ to $rk_D^2(\mu)$ with appropriate choice of R . With this we have a parallel of Galvin–Hajnal theorem (4.5). Here comes the main theorem (4.6) assuming DC , μ singular of uncountable cofinality (and E a set of filters on $cf(\mu)$ which is nice), if $\mu = |\mathcal{H}(\mu)|$ (kind of strong limit) then set theory “behave nicely” up to 2^μ : 2^μ is an aleph, there is no measurable $\leq 2^\mu$ and μ^+ is regular.

We end defining some natural ideals in this context of $\otimes_{\mu,R}$.]

5 The successor of a singular of uncountable cofinality

[Here we prove our second main theorem: if $\langle \lambda_i : i < \kappa \rangle$ is increasing continuous (E a nice family of filters on κ) then for at most one λ , for stationarily many $i < \kappa$, the cardinal λ_i^+ has cofinality λ .]

6 Nice E exists

[The theorems so far have as a major assumption the existence of a nice E . Using inner models we show that as in the situation with choice, if there is no nice E then the universe is similar enough to some inner model to answer our questions on the exponentiation and cofinality.]

0 Introduction

Originally I disliked choiceless set theory but ([Sh 176]) discovering (the first modern asymmetry of measure/category: a difference in consistency strength and)

[ZF + DC]

if there is a set of \aleph_1 reals then there is a Lebesgue non-measurable set

I have softened. Recently Gitik suggested to me to generalize the pcf theory to the set theory without choice, or more exactly with limited choice. E.g.: is there a restriction on $\langle \text{cf}(\aleph_n) : n < \omega \rangle$? By Gitik [Gi] ZF+ “every aleph has cofinality \aleph_0 ” is consistent.

It is known that if ZF + DC + AD then there is a very specific pattern of cofinality, but we have no flexibility. So we do not know e.g. if

$$“ZF + DC + (\forall \delta)[\text{cf}(\delta) \leq \aleph_\alpha] + (\forall \beta < \alpha)(\aleph_{\beta+1} \text{ is regular})”$$

is consistent for $\alpha = 1$, or $\alpha = 2$ etc. The general question is what are the restrictions on the cofinality function; really it is appropriate to dedicate this work to Azriel Levy, my teacher and my friend, who has contributed so much to this direction.

Apter repeated the question above and told me of Apter and Magidor [ApMg] in which the consistency of ZF + DC $_{\aleph_\omega}$ + $|\mathcal{H}(\aleph_\omega)| = \aleph_\omega$ + “ $\aleph_{\omega+1}$ is measurable” was proved (for an earlier weaker result see Kafkoulis [Kf]) and was quite confident that a parallel theorem can be similarly proved for \aleph_{ω_1} .

My feeling was that while the inner model theory and the descriptive set theory are not destroyed by dropping AC, modern combinatorial set theory says essentially nothing in this case, and it will be nice to have such a theory.

Our results may form a modest step in this direction. The main result is stated in the abstract.

Theorem 0.1 (ZF + DC) *If μ is singular of uncountable cofinality and is strong limit in the sense that $\mathcal{H}(\mu)$ has cardinality μ then μ^+ is regular and non measurable.*

Note that this work stress the difference between “bounds for cardinal arithmetic for singulars with uncountable cofinality” and the same for countable cofinality.

Another theorem (see section 5) says

Theorem 0.2 *If $\langle \mu_i : i \leq \kappa \rangle$ is an increasing continuous sequence of alephs $> \kappa$, then for at most one λ , $\{i : \text{cf}(\mu_i^+) = \lambda\}$ is a stationary subset of κ (see 3.17 (3)).*

We were motivated by a parallel question in ZFC, asked by Magidor and to which we do not know the answer: can \aleph_{ω_1} be strong limit and for $\ell \in \{1, 2\}$ there is a stationary set $S_\ell \subseteq \omega_1$ such that $\prod_{\delta \in S_\ell} \aleph_{\delta+2}/J_S^{\text{bd}}$ has true cofinality $\aleph_{\omega_1+\ell}$?

It was known that “there are two successive singulars” has large consistency strength.

We do not succeed to solve Woodin's problem (show consistency of “ZF + DC_{ω} + every aleph has cardinality \aleph_0 or \aleph_1 ”), and Specker's problem (show consistency of “ZF + every $\mathcal{P}(\kappa)$ is the union of countably many sets of cardinality $\leq \kappa$ ”). For me the real problems are:

- (a) (ZF) Is there an aleph κ (i.e. suitable definition) such that DC_{κ} implies that the class of regular alephs is unbounded?
- (b) For which $\kappa < \lambda$, does $DC_{<\kappa}$ implies λ is regular?
- (c) (ZF) Is there κ such that DC_{κ} implies that for every μ for a class of alephs λ , the set $\mathcal{P}(\lambda)$ is not the union of $\leq \mu$ sets each of cardinality λ and $\text{cf}(\lambda) > \kappa$?

We try to assume only standard knowledge in set theory but we start by discussing what part of pcf theory survives (see [Sh-g] and history there) so knowledge of it will be helpful; in particular Sect. 6 imitates [Sh-g, V, Sect. 1] so it also uses a theorem of Dodd and Jensen [DJ]. On set theory without full choice see [J1], on cardinal arithmetic and its history [Sh-g]. Lately Apter and Magidor got consistency results complementary to ours and we intend to return to the problems here in [Sh 589, Sect. 5], for more and on 1.5, 2.5 and (c) above, see [Sh: F239].

Definition 0.3 A cardinality is called an aleph if there is an ordinal with that cardinality, or just the cardinality of a set which can be well ordered and then it is identified with the least ordinal of that cardinality.

Notation 0.4 $\alpha, \beta, \gamma, \epsilon, \zeta, \xi, i, j$ are ordinals; δ a limit ordinal; $\lambda, \mu, \kappa, \chi, \theta, \sigma$ are cardinals (not necessarily alephs),

Reg is the class of regular alephs (see Definition 1.1(7)),

D a filter on the set $\text{Dom}(D)$,

$A = \emptyset \text{ mod } D$ means $\text{Dom}(D) \setminus A \in D$,

$D^+ = \{A : A \subseteq \text{Dom}(D), \text{ and } A \neq \emptyset \text{ mod } D\}$,

$D + A = \{X \subseteq \text{Dom}(D) : X \cup (\text{Dom}(D) \setminus A) \in D\}$.

For a filter D , $\theta_D = \min\{\kappa : \text{there is no function } h : \text{Dom}(D) \rightarrow \kappa \text{ such that for every } i < \kappa \text{ we have } h^{-1}(i) \neq \emptyset \text{ mod } D\}$.

\mathcal{I}, \mathcal{J} denote ideals on the set $\text{Dom}(\mathcal{I}), \text{Dom}(\mathcal{J})$ respectively; definitions given for filters D apply also to the dual ideal $\{\text{Dom}(D) \setminus A : A \in D\}$ but $\mathcal{I} + A = \{B \subseteq \text{Dom}(\mathcal{I}) : B \setminus A \in \mathcal{I}\}$.

I, J are directed partial orders or just index sets.

A cone of a partial order I is a subset of the form $\{a \in I : I \models a_0 \leq a\}$ for some $a_0 \in I$.

For a set A let $\theta(A)$ be

$\sup\{\alpha : \alpha \text{ is an ordinal and there is a function from } A \text{ onto } \alpha \text{ or } \alpha \text{ is finite}\}$.

Note that if $|A|$ is an aleph then $\theta(A) = |A|^+$. Also $\theta(A)$ is an aleph. If $g : A \rightarrow \delta$ has unbounded range then $\text{cf}(\delta) \leq \theta(A)$ (and $|\text{Rang}(g)|$ is an aleph $\leq \theta(A)$) see Definition 1.1(7).

Let $\theta^-(A)$ be

$\min\{\alpha : \alpha \text{ an ordinal and there is no one to one function from } \alpha \text{ into } A\};$

clearly $\theta^-(A^*)$ is an aleph and $\theta^-(A) \leq \theta(A)$.

For a directed partial order I , $\mathcal{I}_I^{\text{bd}}$ is the ideal of bounded subsets of I and $\mathcal{G}_I^{\text{bd}}$ is the dual filter (usually I is a regular aleph).

If $f, g : A^* \rightarrow \text{Ord}$ and D is a filter on A^* then

$$f \leq_D g \quad \text{means} \quad \{a \in A^* : f(a) \leq g(a)\} \in D,$$

similarly for other relations ($<_D, =_D, \neq_D$). Note: \leq_D is a quasi order (as maybe $f_0 \leq_D f_1 \leq_D f_0, f_0 \neq f_1$) but $\langle^{(A^*)}\text{Ord}/D, <_D\rangle$ is a partial order and \neq_D is not the negation of $=_D$.

Definition 0.5 Let $|A| \leq^* |B|$ mean that there is a function from B onto A (so $|A| \leq |B| \Rightarrow |A| \leq^* |B|$ but not necessarily the converse). (Note: for well ordered sets $\leq, <^*$ are equal, in fact “ B is well ordered” is enough.)

Definition 0.6

1. $AC_{\lambda, \bar{\mu}}$ is the axiom of choice for every family $\{A_t : t \in I\}$ of sets, $|A_t| \leq \mu_t$, where $\bar{\mu} = \langle \mu_t : t \in I \rangle$, I a set of cardinality $\leq \lambda$. If $\mu_t = \mu$ for all $t \in I$ then we write $AC_{\lambda, \mu}$. We may write $AC_{I, \mu}, AC_{I, \bar{A}}$ instead, similarly below.
2. If in part (1), I is a well ordering (e.g. an ordinal) then $DC_{I, \bar{\mu}}$ is the dependent choice version.
 $DC_{I, \mu}$ is $DC_{I, \bar{\mu}}$ where $(\forall t \in I)\mu_t = \mu$.
3. Let AC_λ mean $\forall \mu AC_{\lambda, \mu}$; and $AC_{<\lambda} \equiv (\forall \mu < \lambda) AC_\mu$; and $AC_{\lambda, <\mu}, AC_{<\lambda, \mu}, AC_{<\lambda, <\mu}, AC_{\lambda, \bar{\mu}}, AC_{<\lambda, \bar{\mu}}, AC_{<\lambda, <\bar{\mu}}$, have the natural meanings.
4. DC_α means $\forall \mu DC_{\alpha, \mu}$, and $DC_{<\alpha}, DC_{\alpha, <\mu}, DC_{<\alpha, \mu}$ and $DC_{<\alpha, <\mu}$ have the natural meanings. DC is DC_{\aleph_0} .

1 Exact upper bound

Definition 1.1

1. A partial order I is $(\leq \lambda)$ -directed (or $(< \lambda)$ -directed) if for every $A \subseteq I$ of cardinality $\leq \lambda$ (of cardinality $< \lambda$) there is an upper bound. If $\lambda = 2$ we omit it. Note that 2-directed is equivalent to $(< \aleph_0)$ -directed.
2. We say that J is cofinal in I if $J \subseteq I$ and $(\forall s \in I)(\exists t \in J)[I \vDash s \leq t]$.
3. I is endless if $(\forall s \in I)(\exists t \in I)[I \vDash s < t]$.
4. We say $\lambda \geq \text{cf}(I)$ if there is a cofinal $J \subseteq I$ of cardinality $\leq \lambda$ and $\lambda = \text{cf}(I)$ if λ is the smallest such λ i.e. $(\forall \lambda_1)(\lambda_1 \geq \text{cf}(I) \rightarrow \lambda_1 \geq \lambda)$ (it does not necessarily exist, but is unique by the Cantor-Bernstein theorem).
5. Let F be a set of functions from A^* to ordinals, D a filter on A^* . We say $g : A^* \rightarrow \text{Ord}$ is a $<_D$ -eub (exact upper bound) of F if:
 - (i) $f \in F \Rightarrow f \leq_D g$

- (ii) If $h : A^* \rightarrow \text{Ord}$, $h < \max\{g, 1\}$
 then for some $f \in F$ we have $h <_D \max\{f, 1\}$ (where $f^* = \max\{f, 1\}$
 means f^* is a function, $\text{Dom}(f^*) = \text{Dom}(f)$ and for every $x \in \text{Dom}(f^*)$
 we have $f^*(x) = \max\{f(x), 1\}$).
6. Let I be $(\leq \lambda)^*$ -directed if in (1) we ask $|A| \leq^* \lambda$. We define similarly " $\text{cf}^*(I) \leq \lambda$ ", and also " I is $(< \lambda)^*$ -directed", " $\text{cf}^*(I) < \lambda$ ", " $\text{cf}^*(I) = \lambda$ " (e.g. $\text{cf}^*(I) = \lambda$ means $(\forall \lambda_1)[\lambda_1 \geq \text{cf}^*(I) \rightarrow \lambda \leq^* \lambda_1]$ & $\text{cf}^*(I) \geq \lambda$, so actually we should say $\text{cf}^*(I) = \lambda / \equiv^*$ where $\lambda_1 \equiv^* \lambda_2$ iff $\lambda_1 \leq^* \lambda_2 \leq^* \lambda_1$. Instead " $(\leq \lambda)^*$ -" we may write " $(\leq^* \lambda)$ -" . Let λ -directed mean $(< \lambda)$ -directed, etc.
7. The ordinal δ is regular if $\delta = \text{cf}(\delta)$ where

$$\text{cf}(\alpha) = \min\{\text{otp}(A) : A \subseteq \alpha \text{ is unbounded}\}.$$

Clearly if I is well ordered then $\text{cf}(I)$ is a regular ordinal, and a regular ordinal is a cardinal.

8. We say A is unbounded in I if $A \subseteq I$ and for no $t \in I \setminus A$ do we have $(\forall s \in A)(s \leq_I t)$.
9. $\text{cf}_-(I) \leq \lambda$ if there is an unbounded $A \subseteq I$, $|A| \leq \lambda$; similarly $\text{cf}_+^*(I) < \lambda$, $\text{cf}_+^*(I) = \lambda$.
10. $\text{hcf}_-(I) \leq \lambda$ (the hereditary $\text{cf}_-(I)$ is $\leq \lambda$) if for every $J \subseteq I$, $\text{cf}_-(J) \leq \lambda$; similarly $\text{hcf}_+^*(I) < \lambda$, $\text{hcf}_+^*(I) = \lambda$.

Definition 1.2 Let I be a $(< \aleph_0)$ -directed (equivalently a directed) partial order (often I will be a limit ordinal δ with its standard order, then we write just δ).

1. $\bar{F} = \langle F_t : t \in I \rangle$ is $<_D$ -increasing (or \leq_D -increasing) if:
- (a) $\bigcup_{t \in I} F_t \subseteq \text{Dom}(D)\text{Ord}$ and $F_t \neq \emptyset$,
- (b) if $f_e \in F_{t_e}$, (for $e < 2$) and $t_0 <_I t_1$ then $f_0/D < f_1/D$ (or $f_0/D \leq f_1/D$).
2. \bar{F}/D is smooth (or \bar{F} is D -smooth) if:
- (a) $\bigcup_{t \in I} F_t \subseteq \text{Dom}(D)\text{Ord}$ and $F_t \neq \emptyset$,
- (c) $f_1, f_2 \in F_t \Rightarrow f_1/D = f_2/D$.
3. Let $\bar{F} = \langle F_t : t \in I \rangle$, we say \bar{F}/D is semi-smooth (or \bar{F} is semi- D -smooth) if:
- (a) $\bigcup_{t \in I} F_t \subseteq \text{Dom}(D)\text{Ord}$ and $F_t \neq \emptyset$,
- (c)' if $I \models "t_0 < t_1 < t_2 < t_3"$ and $f_l \in F_{t_l}$ for $l < 4$ then

$$\{a \in \text{Dom}(D) : f_1(a) < f_2(a)\} \subseteq \{a \in \text{Dom}(D) : f_0(a) < f_3(a)\} \text{ mod } D,$$

4. \bar{F} is almost D -smooth (or \bar{F}/D is almost smooth) is defined similarly but
- (c)'' if $I \models t_0 < t_1$ and $f_0, f'_0 \in F_{t_0}$, and $f_1, f'_1 \in F_{t_1}$ then

$$\{a \in \text{Dom}(D) : f_0(a) < f_1(a)\} = \{a \in \text{Dom}(D) : f'_0(a) < f'_1(a)\} \text{ mod } D,$$

5. $f \in \text{Dom}(D)\text{Ord}$ is a lub (least upper bound) of \bar{F}/D (or $<_D$ -lub of \bar{F}) if:
- (a) $f \in \bigcup_{t \in I} F_t \Rightarrow f \leq_D g$,
- (b) if $g' \in \text{Dom}(D)\text{Ord}$ satisfies (a) too then $g \leq_D g'$.

6. $g \in {}^{\text{Dom}(D)}\text{Ord}$ is the eub of \bar{F}/D (or $<_D$ -eub of \bar{F}) if
- (a) $f \in \bigcup_{t \in I} F_t \Rightarrow f <_D g$,
- (b) if $f^* <_D \max\{g, \mathbf{1}_{A^*}\}$, then $f^* <_D \max\{f, \mathbf{1}_{A^*}\}$; note that if for some $g \in F$ we have $g \neq_D \mathbf{0}_{A^*}$ then this is equivalent to: if $f^* <_D g$ then

$$(\exists f \in \bigcup_{t \in I} F_t)(f^* <_D f)$$

(note: if $\neg(g \neq_D \mathbf{0}_{A^*})$ this holds vacuously).

Observation 1.3 1. We have $\text{cf}(0) = 0$, $\text{cf}(\alpha + 1) = 1$. For a limit ordinal δ , $\text{cf}(\delta) = \text{cf}^*(\delta)$ is regular and infinite; each regular ordinal is an aleph. For a linear order I , $\text{cf}^*(I) \leq \lambda$ iff I is not $(\leq \lambda)^*$ -directed. For a limit ordinal δ we have: $\text{cf}(\delta) \leq^* |A|$ iff $\text{cf}(\delta) < \theta(A)$.

2. If \bar{F} is $<_D$ -increasing then it is semi- D -smooth.
 3. If \bar{F} is $<_D$ -increasing then it is almost D -smooth.
 4. If I is a partial order, $|I|$ an aleph, then there is $J \subseteq I$, linearly and well ordered by $<_I$ with no upper bound in $I \setminus J$.
 5. We are assuming that each F_t is a set; if we consider classes, intersect them with $\mathcal{H}(\chi)$ for χ large enough.
 6. If $\bar{Y} = \langle Y_\alpha : \alpha < \delta \rangle$ is a sequence of subsets of A and $\alpha < \beta < \delta$ implies $Y_\alpha \subseteq Y_\beta$ and $\text{cf}(\delta) \geq \theta(A)$ then \bar{Y} is eventually constant i.e.

$$(\exists \alpha < \delta)(\forall \beta)(\alpha \leq \beta < \delta \rightarrow Y_\beta = Y_\alpha)$$

7. If $\bar{F} = \langle F_\alpha : \alpha < \delta \rangle$ is \leq_D -increasing, $A^* = \text{Dom}(D)$ and $\theta(\mathcal{P}(A^*)/D) \leq \text{cf}(\delta)$, then we can define, in a uniform way, from \bar{F} a club C of δ and $Y/D \in \mathcal{P}(A^*)/D$ such that:
- (a) $\bar{F} \upharpoonright C$ is $<_{D+Y}$ -increasing,
 (b) $f_1, f_2 \in \bigcup_{t \in C} F_t \Rightarrow f_1 =_{D+(A^* \setminus Y)} f_2$.
8. If $\langle F_\alpha : \alpha < \delta \rangle$ is $<_D$ -increasing, f is a \leq_D -lub of $\bigcup_{\alpha < \delta} F_\alpha$
 then $\{a \in A^* : f(a) \text{ a limit ordinal}\} \in D$.

Remark. Not $\theta(\mathcal{P}(A^*)) \geq \theta(\mathcal{P}(A^*)/D)$ as $|\mathcal{P}(A^*)|^* \geq |\mathcal{P}(A^*)/D|$.

Proof. (1)–(6), (8) Check.

(7) For $\alpha < \beta < \delta$ let

$$E_{\alpha, \beta}^1 = \{Y/D : Y \subseteq A^* \text{ and: } Y = \emptyset \text{ mod } D \text{ or } Y \in D^+ \\ \text{and for some } f_1 \in F_\alpha \text{ and } f_2 \in F_\beta \text{ we have } f_1 <_{D+Y} f_2\}$$

$$E_{\alpha, \beta}^0 = \{Y/D : Y \subseteq A^*, Y = \emptyset \text{ mod } D \text{ or } Y \in D^+ \\ \text{and for every } f_1 \in F_\alpha, f_2 \in F_\beta \text{ we have } f_1 <_{D+Y} f_2\}$$

Clearly

$$(*)_1 \ E_{\alpha, \beta}^1 \subseteq \mathcal{P}(A^*)/D, \emptyset/D \in E_{\alpha, \beta}^0 \subseteq E_{\alpha, \beta}^1, \text{ and } \alpha_1 \leq \alpha_2 < \beta_2 \leq \beta_1 \Rightarrow \\ E_{\alpha_2, \beta_2}^\ell \subseteq E_{\alpha_1, \beta_1}^\ell$$

and

(*)₂ if $\alpha_1 < \alpha_2 < \beta_2 < \beta_1$ then $E_{\alpha_2, \beta_2}^1 \subseteq E_{\alpha_1, \beta_1}^0$.

As $\text{cf}(\delta) \geq \theta(\mathcal{P}(A^*)/D)$, for $\ell \in \{0, 1\}$ and $\alpha < \delta$ the sequence $\langle E_{\alpha, \beta}^\ell : \beta \in [\alpha, \delta) \rangle$ is eventually constant (by 1.3(6)), so let $\gamma_\alpha^\ell \in (\alpha, \delta)$ be minimal such that $\gamma_\alpha^\ell \leq \gamma < \delta \Rightarrow E_{\alpha, \gamma}^\ell = E_{\alpha, \gamma_\alpha^\ell}^\ell$.

Let $E_\alpha^\ell = E_{\alpha, \gamma_\alpha^\ell}^\ell$. Clearly

(*)₃ $E_\alpha^0 \subseteq E_\alpha^1$,

(*)₄ $\ell < 2$ & $\alpha < \beta \Rightarrow E_\alpha^\ell \subseteq E_\beta^\ell \subseteq \mathcal{P}(A^*)/D$,

(*)₅ $\alpha < \beta \Rightarrow E_\alpha^1 \subseteq E_\beta^0 \subseteq \mathcal{P}(A^*)/D$.

Applying again 1.3(6), by (*)₄, we find that for some $\alpha(*) < \delta$ we have $\alpha(*) \leq \alpha < \delta \Rightarrow E_\alpha^0 = E_{\alpha(*)}^0$ so by (*)₃ & (*)₅ we get $\alpha(*) < \alpha < \delta \Rightarrow E_\alpha^0 = E_\alpha^1 = E_{\alpha(*)}^1$.

Choose by induction on ε an ordinal $\beta_\varepsilon < \delta$: for $\varepsilon = 0$ let $\beta_\varepsilon = \alpha(*) + 1$, for $\varepsilon = \zeta + 1$ let $\beta_\varepsilon = \max\{\gamma_{\beta_\zeta}^0, \gamma_{\beta_\zeta}^1\} < \delta$ and for ε limit $\beta_\varepsilon = \bigcup_{\zeta < \varepsilon} \beta_\zeta$. So for some

limit ordinal $\varepsilon(*)$ we have: β_ε is defined iff $\varepsilon < \varepsilon(*)$, and $\langle \beta_\varepsilon : \varepsilon < \varepsilon(*) \rangle$ is strictly increasing with limit δ .

Choose $f^* \in F_{\beta_0}, f^{**} \in F_{\beta_1}$ and let $Y^* = \{x \in A^* : f^*(x) < f^{**}(x)\}$. Clearly $Y^* \in E_{\beta_0, \beta_1}^1 = E_{\beta_0}^1 = E_{\alpha(*)}^1 = E_{\alpha(*)}^0$ hence

$$f' \in F_{\beta_0} \text{ \& } f'' \in F_{\beta_1} \Rightarrow f' <_{D+Y^*} f''.$$

So clearly Y^*/D does not depend on the choice of f^*, f^{**} , i.e. $f' \in F_{\beta_0}$ & $f'' \in F_{\beta_1} \Rightarrow \{x \in A^* : f'(x) < f''(x)\} = Y^* \pmod{D}$.

In fact we can replace (β_0, β_1) by any $(\beta_{\varepsilon_0}, \beta_{\varepsilon_1})$ with $\varepsilon_0 < \varepsilon_1 < \varepsilon(*)$. So clearly the conclusion holds with $Y = A^* \setminus Y^*$. $\square_{1.3}$

Claim 1.4

1. If $\langle f_\beta : \beta < \delta \rangle$ is such that: $f_\beta : A^* \rightarrow \text{Ord}$ and¹

$$\delta \geq \theta^-(\bigcup_{\alpha < \theta(A^*)} \mathcal{P}(A^* \times \alpha))$$

then for some $\beta < \gamma < \delta$ we have $f_\beta \leq f_\gamma$.

2. $[AC_{\lambda, |I|}]$ If I is $(\leq \lambda)^*$ -directed, H an increasing function from I to J , $|J| \leq \lambda$ (I, J partial orders), then H is constant on a cone of I .
3. If I is $(< \theta(J))$ -directed, $|I|$ an aleph, H an increasing function from I to J and $\theta(J) \leq \lambda$ then H is constant on a cone of I .
4. $[AC_{\lambda, |I|}, AC_{A, B}]$ If I is $(\leq^* \lambda)$ -directed, H an increasing function from I to $J = {}^A B$, B partially ordered set, $\text{hcf}_*(B) \leq \lambda$ (J ordered by: $f \leq g \iff (\forall a \in A)(f(a) \leq g(a))$), and $|A| \leq \lambda$ then H is constant on a cone.
5. In part (2), instead of $|J| \leq \lambda$, actually $\text{hcf}_*(J) \leq \lambda$ suffices (see Definition 1.1(10)).

¹ See 4.5 below

6. [AC \aleph_0] If D is an \aleph_1 -complete filter on A^* and $\bar{F} = \langle F_\alpha : \alpha < \delta \rangle$ is almost D -smooth, and $\delta \geq \theta^-(\bigcup\{\alpha(\mathcal{P}(A^*)/D) : \alpha < \theta(\mathcal{P}/A^*)/D \text{ or } \alpha < \omega\})$ then for some $\beta < \gamma < \delta$, $f_\beta \in F_\beta$, $f_\gamma \in F_\gamma$ we have $f_\beta \leq_D f_\gamma$.

Proof. 1) Assume not. For each $\beta < \delta$ choose by induction on i , $\alpha(\beta, i) = \alpha_{\beta, i}$ as the first ordinal α such that:

(*) $\alpha < \beta$ and for every $j < i$, $\alpha_{\beta, j} < \alpha$ and for $a \in A^*$

$$[f_{\alpha(\beta, j)}(a) > f_\beta(a) \iff f_{\alpha(\beta, j)}(a) > f_\alpha(a)].$$

So $\alpha_{\beta, i}$ is strictly increasing with i and is $< \beta$, hence $\alpha_{\beta, i}$ is defined iff $i < i_\beta$ for some $i_\beta \leq \beta$. The sequence $\langle \langle \alpha_{\beta, i} : i < i_\beta \rangle : \beta < \delta \rangle$ exists. Now for $\beta < \delta$, $i < j < i_\beta$ let $u_{\beta, i, j} = \{a \in A^* : f_{\alpha(\beta, i)}(a) > f_{\alpha(\beta, j)}(a)\}$. For each $a \in A^*$ and $\beta < \delta$ we have:

$v_\beta(a) = \{i < i_\beta : f_{\alpha(\beta, i)}(a) > f_\beta(a)\}$ is a subset of i_β and

$$\langle f_{\alpha(\beta, i)}(a) : i \in v_\beta(a) \rangle$$

is a strictly decreasing sequence of ordinals; hence $v_\beta(a)$ is finite.

Now $i_\beta = \bigcup_{a \in A^*} v_\beta(a)$ (as if $j \in i_\beta \setminus \bigcup_{a \in A^*} v_\beta(a)$ then $\alpha_{\beta, j}, \beta$ are as required). So we have a function v_β from A^* onto $\{v_\beta(a) : a \in A^*\}$, a set of finite subsets of i_β whose union is i_β . Now this set is well ordered of cardinality $|i_\beta|$ (or both are finite), so $i_\beta < \theta(A^*)$.

Let $A_{\beta, i} = \{a \in A^* : i \in v_\beta(a)\} = \{a \in A^* : f_{\alpha(\beta, i)}(a) > f_\beta(a)\}$; also for no $\beta_1 \neq \beta_2 < \delta$ do we have $\langle \alpha_{\beta_1, i}, A_{\beta_1, i} : i < i_{\beta_1} \rangle = \langle \alpha_{\beta_2, i}, A_{\beta_2, i} : i < i_{\beta_2} \rangle$: as by symmetry we may assume $\beta_1 < \beta_2$; now β_1 is a good candidate for being $\alpha_{\beta_2, i_{\beta_2}}$ so $\alpha_{\beta_2, i_{\beta_2}}$ is well defined and $\leq \beta_1$, contradicting the definition of i_{β_2} . Similarly

$$\beta_1 < \beta_2 < \delta, \quad \forall i < i_{\beta_2} [i < \alpha_{\beta_1} \ \& \ \alpha_{\beta_2, i} = \alpha_{\beta_1, i} \ \& \ A_{\beta_2, i} = A_{\beta_1, i}]$$

is impossible. Clearly if $j < i_\gamma$, $\beta = \alpha_{\gamma, j}$ then $\langle \alpha_{\beta, i} : i < i_\beta \rangle = \langle \alpha_{\gamma, i} : i < j \rangle$.

Now for every $\beta < \delta$ let us define $c_\beta = \{\alpha_{\beta, i} : i < i_\beta\}$. So $|c_\beta| < \theta(A^*)$. Let $X_\beta = \{(i, a) : i < i_\beta \text{ and } a \in A_{\beta, i}\}$ so $X_\beta \subseteq i_\beta \times A^*$ and $i_\beta < \theta(A^*)$. It is also clear that for $\beta_1 \neq \beta_2 (< \delta)$, $X_{\beta_1} \neq X_{\beta_2}$. (Just let f be a one-to-one order preserving function from c_{β_1} onto c_{β_2} , let $\alpha \in c_{\beta_1}$ be minimal such that $f(\alpha) \neq \alpha$ and we get a contradiction to the previous paragraph.)

So there is a one-to-one function from δ into $\bigcup\{\mathcal{P}(A^* \times \alpha) : \alpha < \theta(A^*)\}$. But $\delta \geq \theta^-(\bigcup_{\alpha < \theta(A^*)} \mathcal{P}(A^* \times \alpha))$. So we are done.

2) Let for $t \in J$, $I_t = \{s \in I : H(s) = t\}$. So $\{I_t : t \in J, I_t \neq \emptyset\}$ is an indexed family of $\leq |J|$ nonempty subsets of I , $|I_t| \leq |I|$, so by assumption there is a choice function F for this family. Let $A = \{F(I_t) : t \in J, I_t \neq \emptyset\}$, so $|A| \leq |J|$, so as I is $(\leq \lambda)^*$ -directed and $|J| \leq \lambda$ there is $s(*) \in I$ such that $(\forall t \in J)(F(I_t) \leq_I s(*)$.

Thus $H(s(*)$) is the largest element in the range of H . As H is increasing it must be constant on the cone $\{s : s(*) \leq_I s\}$.

3) Like part (2) (with $F(I_t)$ being the first member of I_t for some fixed well ordering of I).

4) By the proof of part (5) below it suffices to prove $\text{cf}_-^*({}^A B \upharpoonright Y, \leq) \leq \lambda$; where $Y = \text{Rang}(H) \subseteq {}^A B$, clearly Y is $(\leq^* \lambda)$ -directed (proved as in the proof of (2)). Let for $(a, b) \in A \times B$,

$$Y_{(a,b)} = \{h : h \in Y \text{ and } b = h(a)\}.$$

Let for each $a \in A$, $B_a = \{b \in B : \text{for some } h \in Y \text{ we have } h(a) = b\} \subseteq B$.

Case 1 for some $a \in A$, B_a is a non empty subset of A with no last element. B_a has an unbounded subset B^* , $|B^*| \leq^* \lambda$. As Y is $(\leq^* \lambda)$ -directed, to get contradiction it suffices to find Z , $|Z| \leq^* \lambda$, Z not disjoint from any set in $\{Y_{(a,b)} : b \in B^*\}$; by $AC_{\lambda,|Y|}$, as $|Y| \leq^* |I|$, one exists.

Case 2 not case 1.

Let $h^* \in {}^A B$ be $h(a) =$ the maximal element of B_a . If h^* is in Y then it is the maximal member of Y ; so $\text{cf}_-^*({}^A B \upharpoonright Y, \leq) = 1 \leq \lambda$. If not to get contradiction it suffices to have a choice function for $\{Y_{a,h^*(a)} : a \in A\}$ which again exists.

5) Like the proof of part (2). Let $J_1 = \text{Rang}(H)$ and $J_2 \subseteq J_1$ be unbounded in J_1 , $|J_2| \leq \lambda$ and choose F as a choice function for $\{I_t : t \in J_2, I_t \neq \emptyset\}$.

6) Let for $\beta < \gamma < \delta$, $Y_{\beta,\gamma} \in \mathcal{P}(A^*)/D$ be such that : for every $f_\beta \in F_\beta$ and $f_\gamma \in F_\gamma$ we have $Y_{\beta,\gamma} =: \{a \in A^* : f_\beta(a) > f_\gamma(a)\}/D$. We can assume toward contradiction that the conclusion fails, so $\beta < \gamma < \delta \Rightarrow Y_{\beta,\gamma} \neq \emptyset/D$. For each $\beta < \delta$ we define by induction on i an ordinal $\alpha_{\beta,i} = \alpha(\beta, i)$ as the first α such that:

$$(\forall j < i)(\alpha_{\beta,j} < \alpha < \beta) \text{ and } (\forall j < i)(Y_{\alpha_{\beta,i},\alpha} = Y_{\alpha_{\beta,i},\beta}).$$

So clearly $\alpha_{\beta,i}$ increase with i and is $< \beta$, hence for some $i_\beta \leq \beta$ we have $\alpha_{\beta,i}$ is defined iff $i < i_\beta$, and $\langle \langle \alpha_{\beta,i} : i < i_\beta \rangle : \beta < \delta \rangle$ exists. Clearly for $i < j < i_\beta$, $Y_{\alpha_{\beta,i},\alpha_{\beta,j}} = Y_{\alpha_{\beta,i},\beta}$. If for some β and Y the set $\{i < i_\beta : Y_{\alpha_{\beta,i},\beta} = Y\}$ is infinite, let j_n the n -th member. By AC_{\aleph_0} we can find $\langle f_{j_n} : n < \omega \rangle$ with $f_{j_n} \in F_{j_n}$, so $Y_{\alpha_{\beta,j_n},\alpha_{\beta,j_{n+1}}} = Y$, so

$$A_n = \{a \in A^* : f_{\alpha_{\beta,j_n}}(a) > f_{\alpha_{\beta,j_{n+1}}}(a)\} \in Y_{\alpha_{\beta,j_n},\alpha_{\beta,j_{n+1}}} = Y.$$

Hence $\bigcap_{n < \omega} A_n \neq \emptyset$ (as $D + A_0$ is a proper \aleph_1 -complete filter remembering $Y_{\beta,\gamma} \neq \emptyset/D$) and we get a strictly decreasing sequence of length ω of ordinals $\langle f_{\alpha_{\beta,j_n}}(a) : n < \omega \rangle$ for $a \in \bigcap_{n < \omega} A_n$, a contradiction. So $i_\beta < \theta(\mathcal{P}(A^*)/D)$. (In fact let

$b_\beta = \{i < i_\beta : Y_{\alpha_{\beta,i},\beta} \notin \{Y_{\alpha_{\beta,j},\beta} : j < i\}\}$, then $|b_\beta| = |i_\beta|$ or both are finite, and clearly $\text{otp}(b_\beta) < \theta^-(\mathcal{P}(A^*)/D)$, but $\theta^-(\mathcal{P}(A^*)/D)$ is an aleph or finite number, hence $i_n < \omega \vee i_n < \theta^-(\mathcal{P}(A^*)/D)$). So $\beta \mapsto \langle Y_{\alpha_{\beta,i}} : i < i_\beta \rangle$ is a one-to-one function from δ to $\bigcup_{\alpha < \theta(\mathcal{P}(A^*)/D)} \alpha(\mathcal{P}(A^*)/D)$ [why? because if

$\beta_1 \neq \beta_2$ have the same image, without loss of generality $\beta_1 < \beta_2$ and β_1 could have served as $\alpha_{\beta_1,i_{\beta_2}}$, contradiction so the mapping is really one-to-one]. $\square_{1.4}$

Claim 1.5 [DC_κ , κ an aleph $> |\mathcal{P}(A^*)|$ and: $AC_{\mathcal{P}(A^*)}$ or $|I|$ is an aleph]

Assume:

- (i) D is a filter on the set A^*
- (ii) I is a $(\leq |\mathcal{P}(A^*)|)^*$ -directed partial order
(e.g. an ordinal δ , $\text{cf}(\delta) \geq \theta(\mathcal{P}(A^*))$)
- (iii) $F = \{f_t : t \in I\}$ is \leq_D -increasing, $f_t : A^* \rightarrow \text{Ord}$

Then F has a \leq_D -eub

Proof. Let $\beta^* = (\sup \bigcup_{t \in I} \text{Rang}(f_t) + 1)$, it is an ordinal, and let $X = A^*(\beta^*)$. By DC_κ as $|\mathcal{P}(A^*)| < \kappa$, so without loss of generality $\kappa = |\mathcal{P}(A^*)|^+$, $|\mathcal{P}(A^*)|$ an aleph.

We can try to choose by induction on $\alpha < \kappa$ functions $g_\alpha \in X$ such that:

- (a) $(\forall \beta < \alpha)(\neg g_\beta \leq_D g_\alpha)$
- (b) $(\forall t \in I)(f_t \leq_D g_\alpha)$
- (c) $g_0(a) = \sup\{f_t(a) : t \in I\}$

By 1.4(1) the construction cannot continue for κ steps; so as DC_κ holds, for some $\alpha = \alpha(*) < \kappa$ we have $\langle g_\alpha : \alpha < \alpha(*) \rangle$ and we cannot choose $g_{\alpha(*)}$. By clause (c) clearly $\alpha(*) > 0$. Let $B_a =: \{g_\beta(a) : \beta < \alpha(*)\}$ so B_a is a set of ordinals, $|B_a| \leq |\alpha(*)|$ (as $|\alpha(*)|$ is an aleph) and $\langle B_a : a \in A^* \rangle$ exists.

Define $H : I \rightarrow \prod_{a \in A^*} B_a$ as follows:

$$H(t)(a) = \min(B_a \setminus f_t(a)) \quad (\text{well defined as } g_0(a) \in B_a).$$

Clearly $|\text{Rang}(H)| \leq |\mathcal{P}(A^*)|^{A^*} = |\mathcal{P}(A^*)|$ (as $|A^*|$ is an infinite aleph hence $|A^* \times A^*| = |A^*|$) and $I \vDash s \leq t \Rightarrow H(g) \leq H(s) \pmod D$. As I is $(\leq |\mathcal{P}(A^*)|)^*$ -directed, by 1.4(2) if $AC_{\mathcal{P}(A^*)}$ and 1.4(3) if $|I|$ is an aleph we know that H is constant on some cone $\{t \in I : s^* \leq t\}$. Now consider $H(s^*)$ as a candidate for $g_{\alpha(*)}$: it satisfies clause (b), clause (c) is irrelevant so clause (a) necessarily fails, i.e. for some $\beta < \alpha^*$, $g_\beta \leq_D H(s^*)$ so necessarily $\alpha(*) = \beta + 1$, and so g_β is as required. $\square_{1.5}$

Discussion 1.5A. In 1.5 the demand DC_κ is quite strong, implying $\mathcal{P}(A^*)$ is well ordered. Clearly we need slightly less than DC_κ - only $DC_{\alpha(*)}$ but $\alpha(*)$ is not given *a priori* so what we need is more than $DC_{|\mathcal{P}(A^*)|}$. Restricting ourselves to \aleph_1 -complete filters we shall do better (see 1.7).

Claim 1.6 (1) $[AC_{|\mathcal{P}(A^*)|, |I|}]$ If (i), (ii), (iii) of 1.5 hold, and g is a \leq_D -lub of F then

(a) g is a \leq_D -eub of F

(1A) $[AC_{\mathcal{P}(A^* \times A^*), I} + AC_{A^*}]$ Assume in addition

(ii)⁺ I is $(\leq |\mathcal{P}(A^* \times A^*)|)^*$ -directed partial order

Then

(b) for some $A \subseteq A^*$ and $t \in I$ we have:

(α) $A \in D^+$ & $t \leq_I s \Rightarrow f_s \upharpoonright A = g \upharpoonright A \pmod D$,

(β) $a \in A \iff [\text{cf}(g(a)) \geq \theta(\mathcal{P}(A^*)) \vee g(a) = f_t(a)]$,

(γ) if $A \notin D^+$, without loss of generality $A = \emptyset$, (changing g)

(2) If (i), (ii), (iii) of 1.5 hold, $|I|$ an aleph, g a \leq_D -lub of F then (a) above holds and if AC_{A^*} also (b) above holds.

(3) Assume:

- (i) D is a filter on A^*
- (ii) I is a partial order, $\theta(\mathcal{P}(A^*)/D)$ -directed
- (iii) $\bar{F} = \langle F_t : t \in I \rangle$ is \leq_D -increasing
- (iv) $|I|$ is an aleph
- (v) g a \leq_D -lub of F

Then (a) above holds. If AC_{A^*} then also (b) holds.

Even waving AC_{A^*} (but assuming (i)-(v)) we have

(b)' for some $A \subseteq A^*$ and $t(*)$

(α) if $A \in D^+$ and $t(*) \leq t \in I$ and $f \in F_t$ then

$$f \upharpoonright A = g \upharpoonright A \text{ mod } D$$

(β) assume (ii)⁺; for no $C \subseteq \text{Ord}$ do we here $|C| < \theta(\mathcal{P}(A))$ and $\{a \in A^* \setminus A : g(a) = \sup(C \cap g(a))\} \neq \emptyset \text{ mod } D$.

Remark. In 1.6(1A) instead $A^* \times A^*$ we can use $A^* \times \zeta$ for every $\zeta < \theta(\mathcal{P}(A^*)/D)$. In 1.6(3) we could have used $\theta_-(\mathcal{P}(A^*)/D)$ instead $\theta(\mathcal{P}(A^*)/D)$.

Proof. 1) (a) Let $f \in {}^A \text{Ord}$, $f < \max\{g, \mathbf{1}_{A^*}\}$. We define a function $H : I \rightarrow \mathcal{P}(A)$ by $H(t) = \{a \in A^* : f(a) \geq \max\{f_t(a), \mathbf{1}\}\}$. By 1.4(2) we know that H is constant on a cone of I , more exactly H' , $H'(t) = H(t)/D$ is increasing and is constant by the proof of 1.4(2). Let $t \in I$ be in the cone. If $H(t) = \emptyset \text{ mod } D$ we are done, otherwise define f^* to be equal to g on $A^* \setminus H(t)$ and equal to f on $H(t)$. Now f^* contradicts the fact that g is a $<_D$ -lub. Now check.

1A) (b) for $A \subseteq A^*$ let $I_A = \{t \in I : A = \{a \in A^* : f_t(a) = g(a)\}\}$ if not empty, and I otherwise. Let us apply $AC_{\mathcal{P}(A^*), I}$ to $\{I_A : A \in \mathcal{P}(A^*)\}$ getting $\{t_A : A \in \mathcal{P}(A)\}$. Let $t(*) \in I$ be an upper bound of $\{t_A : A \in \mathcal{P}(A)\}$ (exists by assumption (ii)). Let

$$A_0 =: \{a \in A^* : f_{t(*)}(a) \neq g(a)\}$$

and

$$A_1 = \{a \in A_0 : g(a) \text{ is a limit ordinal}\}$$

and lastly

$$A' =: \{a \in A_1 : \text{cf}(g(a)) < \theta(\mathcal{P}(A^*))\}.$$

Now clearly $t(*) \leq_t s \in I \Rightarrow f_s < g \text{ mod } (D + A_0)$ and $A_0 = A_1 \text{ mod } D$. If $A' = \emptyset \text{ mod } D$ we are done, so assume $A' \in D^+$. By AC_{A^*} we can find $\langle C_a : a \in A^* \rangle$, such that for $a \in A'$ we have: C_a an unbounded subset of $g(a)$ of order type $\text{cf}[g(a)]$. Let for $a \in A^* \setminus A'$, $C_a = \{0\}$, and for $h \in \prod_{a \in A^*} C_a$ let

$I_h^* =: \{t \in I : h <_D \max\{f_t, \mathbf{1}_{A^*}\}\}$. Now the assumption of 1.6(1A) implies that of 1.6(1), hence clause (a) holds, and hence I_h^* is not empty. As we have AC_{A^*} clearly $|A^*|$ is an aleph, and $\theta(\mathcal{P}(A^*))$ is an aleph with cofinality $> |A^*|$,

hence $\{\text{otp}(C_a) : a \in A^*\}$ is a bounded subset of $\theta(\mathcal{P}(A^*))$, so we can find a function h^* from $\mathcal{P}(A^*)$ onto $\zeta^* = \sup\{\text{otp}(C_a) : a \in A^*\}$. Let h_a be the unique one-to-one order preserving function from $\text{otp}(C_a)$ onto C_a , so $\langle h_a : a \in A^* \rangle$ exists. For $h \in \prod_{a \in A^*} C_a$ let

$$X_h =: \{(a, b) : a \in A^* \text{ and } h_a^{-1} \circ h(a) = h^*(b)\} \subseteq A^* \times A^*.$$

Clearly the mapping $h \mapsto X_h$ is one-to-one from $\prod_{a \in A^*} C_a$ into $\mathcal{P}(A^* \times A^*)$.

Hence by $AC_{\mathcal{P}(A^*), I}$

$$\left| \prod_{a \in A^*} C_a \right| \leq |\mathcal{P}(A^* \times A^*)| = |\mathcal{P}(A^*)|,$$

so we can find a function G , $\text{Dom}(G) = \prod_{a \in A^*} C_a$, $G(h) \in I_h^*$. By (ii) we get that

$\text{Rang}(G)$ has an \leq_I -upper bound $t(**)$, and we can get a contradiction.

2) Follows by (3) (with $F_t =: \{f_t\}$)

3) (a) Let $<^*$ be a well ordering of I . Let $f \in (A^*)\text{Ord}$, $f < \max\{g, \mathbf{1}_{A^*}\}$. For every $A \in D^+$ let $t_{A/D}$ be the $<^*$ -first $t \in I$ such that for some $f_t \in F_t$ we have $\{a \in A^* : f_t(a) \geq f(a)\} = A \text{ mod } D$. Now $A \mapsto t_{A/D}$ is a mapping (maybe partial) from $\mathcal{P}(A^*)/D$ into I and $|I|$ is an aleph, so

$$|\{t_{A/D} : A \in D^+\}| < \theta(\mathcal{P}(A^*)/D) (\leq \theta(\mathcal{P}(A^*))).$$

But I is $\theta(\mathcal{P}(A^*)/D)$ -directed so there is $t(*) \in I$ such that

$$(\forall A \in D^+)(t_{A/D} \leq_I t(*)$$

Any $f_{t(*)} \in F_{t(*)}$ satisfies $f < \max\{f_{t(*)}, \mathbf{1}_{A^*}\}$.

(b) + (b)' For each $A \in D^+$, let $t_{A/D}$ be the $<^*$ -first $t \in I$ such that

$$A \subseteq \{a \in A^* : f_t(a) = g(a)\} \text{ mod } D$$

if there is one. So clearly

$$\{t_{A/D} : A \in D^+ \text{ and } t_{A/D} \text{ is well defined}\}$$

is a set of $<^*$ -order type $< \theta(\mathcal{P}(A^*)/D)$, hence there is $t(*) \in I$ such that

$$A \in D^+ \ \& \ t_{A/D} \text{ well defined} \Rightarrow t_{A/D} \leq_I t(*) .$$

Let

$$A_0 =: \{a \in A^* : f_{t(*)}(a) \neq g(a)\}$$

$$A_1 =: \{a \in A_0 : f(a) \text{ a limit ordinal}\}$$

and

$$A' =: \{a \in A_1 : \text{cf}(g(a)) < \theta(\mathcal{P}(A^*))\},$$

as in the proof of 1.6(1A) we have for $A = A_0$, clause (α) of (b) and of (b)' holds, and $A_1 = A_0 \text{ mod } D$, and if $A' = \emptyset \text{ mod } D$ we are done. As for clause (b)(β) by AC_{A^*} , if it fails then (β) of (b)' fails, so it suffices to prove (b)'. Now clause (α) holds, and we prove (β) as in 1.6 (1A). $\square_{1.6}$

Claim 1.7 [$DC_{\aleph_0} + AC_{\mathcal{P}(A^*)}$] Assume:

- (i) D is an \aleph_1 -complete filter on a set A^* ,
- (ii) I is a $(\leq |\mathcal{P}(A^*)|)^*$ -directed partial order,
- (iii) $F = \{f_t : t \in I\}$ is \leq_D -increasing, $f_t : A^* \rightarrow \text{Ord}$.

Then F has a \leq_D -eub

Remark 1.7A.

1. Given $\bar{F} = \langle F_t : t \in I \rangle$ is \leq_D -increasing (i.e. $[I \models s < t \ \& \ f \in I_s \ \& \ g \in I_t \Rightarrow f \leq_D g]$) we can use $J = (\bigcup_{t \in I} F_t, \leq_D)$ as the partial order instead I with $f_g = g$ for $g \in \bigcup_{t \in I} F_t$, so claim 1.7 applies to $\bar{F} = \langle F_t : t \in I \rangle$ too; similarly for 1.5. Note that if I is $(\leq \lambda)^*$ -directed so is J . Also 1.6 (1) (in the proof replace H by $H' : H'(t) = H(t)/D$), concerning 1.6 (1A) check, and lastly for 1.6 (2) see 1.6 (3))
2. If we want to demand only $AC_{\mathcal{P}(A^*), \lambda}$, then $\lambda = \prod_{\alpha \in A^*} \beta_\alpha$ is large enough.
3. Note $AC_{\mathcal{P}(A^*)}$ can be replaced by $AC_{\mathcal{P}(A^*)/D}$.

Proof. By 1.6(1) it suffices to find a \leq_D -lub. Let for $a \in A^*$,

$$\beta_a =: \sup\{f_t(a) : t \in I\} + 1 \in \text{Ord}.$$

For every $A \in D^+$, there is no decreasing ω -sequence in $(\prod_{a \in A^*} \beta_a, <_{D+A})$; hence by DC_{\aleph_0} there is a function $g \in \prod_{a \in A^*} \beta_a$ satisfying:

- (i)_A $(\forall t \in I)[f_t \leq_{D+A} g]$
- (ii)_A if g' satisfies (i) then $\neg(g' <_{D+A} g)$.

By $AC_{\mathcal{P}(A^*)}$ we can find $\langle g_A : A \in D^+ \rangle$ with $g_A \in \prod_{a \in A} \beta_a$ that satisfies (i)_A + (ii)_A.

Let $B_a = \{g_A(a) : A \in D^+\} \cup \{\beta_a\}$ so $|B_a| \leq |D^+| \leq |\mathcal{P}(A^*)|$

Let $H : I \rightarrow B^* = \prod_{a \in A^*} \{B_a : a \in A^*\}$ be $H(t)(a) = \min(B_a \setminus f_t(a))$. Clearly H is an order preserving mapping from (I, \leq_I) to $(B^*/D, \leq_D)$. Also

$$|B^*/D| \leq^* |B^*| \leq^* \prod_{a \in A^*} |D^+| = |D^+|^{|A^*|} \leq |\mathcal{P}(A^*)|^{|A^*|}.$$

But as $AC_{\mathcal{P}(A^*)}$ holds clearly A^* is well ordered, hence $|A^*| \times |A^*| = |A^*|$, hence $|\mathcal{P}(A^*)|^{|A^*|} = |\mathcal{P}(A^*)|$.

By 1.4(2) we know that H is constant on a cone, say $\{t : s(*) \leq_I t\}$.

Assume $g^* =: H(s(*))$ is not a \leq_D -lub, so some g' exemplifies it. Let $A = \{a \in A^* : g'(a) < g^*(a)\}$, so $A \in D^+$ and g_A is well defined.

By the choice of g_A and g' clearly $\neg(g' <_{D+A} g_A)$ hence

$$A_1 = \{a \in A^* : g'(a) \geq g_A(a)\} \neq \emptyset \text{ mod } D + A$$

So $A_2 = A_1 \cap A \in D^+$; and $g_A \leq_{D+A_2} g'$. But by the choice of $g^* = H(s(*))$ (as $(\forall a)(g_A(a) \in B_a)$) clearly $g^* \leq_{D+A} g_A$. Together we get $g^* \leq_{D+A_2} g'$, but this contradicts the definition of A as $A_2 \subseteq A$. □_{1.7}

Claim 1.8 $[DC_{\aleph_0}]$ Consider the conditions:

- (i) D is an \aleph_1 -complete filter on A^* ,
- (ii) I is a $(\leq |\mathcal{P}(A^*)/D|)^*$ -directed partial order,
- (iii) $\bar{F} = \langle F_t : t \in I \rangle$ is \leq_D -increasing, so $f : A^* \rightarrow \text{Ord}$ for $f \in \bigcup_{t \in I} F_t$,
- (iv) $g : A^* \rightarrow \text{Ord}$ and $(\forall t \in I)(\forall f \in F_t)(f \leq_D g)$,
- (v) no $g' <_D g$ satisfies (iv).

Then the following implications hold:

- (1) If (i), (iii) then some g satisfies (iv).
- (2) If (i), (iii), (iv) then some $g^* \leq g$ satisfies clauses (iv), (v); if for g clauses (i), (iii), (iv), $\neg(v)$ hold, we can ask also $g^* <_D g$.
- (3) $[AC_{\mathcal{P}(A^*)/D, I}]$ If (i)-(v), and \bar{F} is D -smooth then there is D^* , such that
 - (a) D^* an \aleph_1 -complete filter extending D ,
 - (b) g is $<_{D^*}$ -lub of \bar{F} .
- (4) In (3) we can omit smoothness.
- (5) If (i)-(v) and $|I|$ is an aleph then for some D^* clauses (a), (b) of (3) hold (i.e. we can drop AC in part (3)).

Proof. (1), (2) are contained in the proof of 1.7

(3) Let $D^* = \{A \subseteq A^* : \text{for } D + (A^* \setminus A), \text{ clause (v) fails}\}$. Clearly D^* is \aleph_1 -complete filter and $\emptyset \notin D^*$. Clearly \bar{F} is \leq_{D^*} -increasing and g is a \leq_{D^*} -upper bound of \bar{F} . If g is not a \leq_{D^*} -lub, there is $g' \in {}^{(A^*)}\text{Ord}$, a \leq_{D^*} -upper bound of \bar{F} such that $\neg[g \leq_{D^*} g']$, so

$$A' =: \{a \in A^* : g'(a) < g(a)\} \neq \emptyset \text{ mod } D^*.$$

For each $t \in I$, for every $f_t \in F_t$ let

$$Y_t =: \{a \in A^* : f_t(a) > g'(a)\} / D \in D^* / D$$

(well defined as \bar{F} is D -smooth), and $t \mapsto Y_t$ is an increasing function from I to $(D^* / D, \leq_D)$, by 1.4(2) we get a contradiction.

(4) Use 1.9(3) below to regain smoothness.

(5) Left to the reader (just use 1.4(3) instead 1.4(2)). □_{1.8}

We have used

Observation 1.9 1. $[AC_{\mathcal{P}(A^*)/D, I}$ or $|I|$ is an aleph] If $\bar{F} = \langle F_t : t \in I \rangle$ is $<_D$ -increasing (where $f_t : A^* =: \text{Dom}(D) \rightarrow \text{Ord}$), I is $(\leq |\mathcal{P}(A)/D|)^*$ -directed then for every $g : A^* \rightarrow \text{Ord}$ for some $t^* \in I$ and $A \subseteq A^*$, for all $f \in \bigcup_{t^* \leq t \in I} F_t$

$$\{a \in A^* : f(a) > g(a)\} = A \text{ mod } D.$$

Similarly for $f(a) = g(a)$, $f(a) < g(a)$.

2. In (1) in the case $|I|$ is an aleph, “ I is $(< \theta(\mathcal{P}(A)/D))^*$ -directed” suffices.

3. $[AC_{\aleph_0}]$ Assume $\bar{F} = \langle F_t : t \in I \rangle$ is $<_D$ -increasing. Let I be a partial order $(\leq \aleph_0)^*$ -directed,

$$J_0 = \{ \langle t_n : n < \omega \rangle : t_n \in I \text{ and } t_n <_I t_{n+1} \text{ for } n < \omega \},$$

$$\bar{t}^0 \leq_{J_0} \bar{t}^1 \iff [(\forall n < \omega)(\exists m < \omega)(t_n^0 \leq_I t_m^1)],$$

$$\bar{t}^0 <_J \bar{t}^1 \text{ if } \bar{t}^0 \leq_J \bar{t}^1 \&\neg(\bar{t}^1 \leq_J \bar{t}^0).$$

Let $F_t^* = \{ \sup_{n < \omega} f_n : \langle f_n : n < \omega \rangle \in \prod_{n < \omega} F_{t_n} \}$ and $\bar{F}^* = \langle F_t^* : t \in J \rangle$ where

of course $(\sup_{n < \omega} f_n)(a) = \sup_{n < \omega} f_n(a)$. Then

- (a) \bar{F}^* is \leq_D -increasing and smooth
 (b) g is a \leq_D -lub of \bar{F} iff g is a \leq_D -lub of \bar{F}^*
 (c) g is a \leq_D -eub of \bar{F} iff g is a \leq_D -eub of \bar{F}^*
 (d) I, J are equi-directed, what means: there is an embedding H of I into J (a partial order) such that its range is a cofinal subset of J [use $H(t) = \langle t : n < \omega \rangle$].
4. $[AC_{\aleph_0}]$ Assume $I = \delta$ where δ is a limit ordinal of cofinality $> \aleph_0$, and let $\bar{F} = \langle F_\alpha : \alpha < \delta \rangle$ be $<_D$ -increasing (where each $F_\alpha \subseteq {}^{(A^*)}\text{Ord}$ is not empty). Let $J = \{ \alpha < \delta : \text{cf}(\alpha) = \aleph_0 \}$, and for $\alpha \in J$ let

$$F_\alpha^* = \{ \sup_n f_n : \text{for some strictly increasing sequence } \langle \alpha_n : n < \omega \rangle \text{ of ordinals } < \alpha \text{ we have } \alpha = \bigcup_{n < \omega} \alpha_n, \text{ and } f_n \in F_{\alpha_n} \}.$$

Then $\bar{F}^* = \langle F_\alpha^* : \alpha \in J \rangle$ satisfies clauses (a)-(d) from part (3), and of course J is well ordering.

Proof. Check.

2 hpp

Definition 2.1

1. Let Γ be a set of filters on $A^* = A_\Gamma^* = \text{Dom}(\Gamma)$. For an ordinal α , we let $\text{hpp}_\Gamma(\alpha)$ be the supremum of the ordinals $\beta + 1$ for which there is a witness (\bar{F}, D) , which means:
- $D \in \Gamma$,
 - $F = \langle F_\gamma : \gamma < \beta \rangle$, with $F_\gamma \neq \emptyset$,
 - $\bigcup_{\gamma < \beta} F_\gamma \subseteq {}^{(A^*)}\alpha$,
 - \bar{F} is $<_D$ -increasing.
2. $\text{ehpp}_\Gamma(\alpha)$ is defined similarly, but each F_γ a singleton; the definition of $\text{shpp}_\Gamma(\alpha)$ is similar too, but \bar{F} is smooth. We can replace α by $f \in {}^{(A^*)}\text{Ord}$ in all these cases here (i.e. $F \subseteq \prod_{a \in A^*} f(a)$). If $F_\alpha = \{f_\alpha\}$ we may write $F = \langle f_\alpha : \alpha < \beta \rangle$ instead of \bar{F} .

3. If $\Gamma = \{D\}$ then we write D instead of Γ . We say that Γ is \aleph_1 -complete if each $D \in \Gamma$ is \aleph_1 -complete, and similarly for other properties.
4. We define $pp_\Gamma(\alpha)$ as in (1) but add to **(i)**-**(iv)** also
 (v) there exists $\langle \alpha_a : a \in \text{Dom}(D) \rangle$ such that:
 (α) $\theta(A^*) \leq \alpha_a < \alpha$,
 (β) $\text{cf}_D(\langle \alpha_a : a \in A^* \rangle) = \alpha$ (see below), each α_a a limit ordinal,
 (γ) for every $g \in \prod_{a \in A^*} \alpha_a$ there is $f \in \bigcup_{\gamma < \beta} F_\gamma$ such that $g \leq_D f$.
5. $epp_\Gamma(\alpha)$, $spp_\Gamma(\alpha)$ are defined similarly.
6. For a filter D with the domain A^* ,

$$\begin{aligned} & \text{cf}_D(\langle \alpha_a : a \in A^* \rangle) \\ &= \inf\{\text{otp}(C) : C \subseteq \bigcup_{a \in A^*} \alpha_a \text{ and } \{a \in A^* : \alpha_a = \sup(C \cap \alpha_a)\} \in D\}, \\ & \limsup_D(\langle \alpha_a : a \in A^* \rangle) = \min\{\alpha : \{a \in A^* : \alpha_a \leq \alpha\} \in D\}. \end{aligned}$$

Remark. 1) Note that in 2.1(4) clause (β) (i.e. $\text{cf}_D(\langle \alpha_a : a \in A^* \rangle) = \alpha$) is a replacement to “ $\langle \alpha_a : a \in A^* \rangle$ is a sequence of limit ordinals with $\text{tlim}(\text{cf}(\alpha_a) : a \in A^*) = \alpha$ ”.

2) Note pp stands for pseudo power, h for hereditary, s for smooth, e for element (rather than set).

- Observation 2.2** 1. $hpp_\Gamma(\alpha)$, $ehpp_\Gamma(\alpha)$ and $shpp_\Gamma(\alpha)$ increase with α and Γ ; and $pp_\Gamma(\alpha)$, $epp_\Gamma(\alpha)$ and $spp_\Gamma(\alpha)$ increase with Γ .
2. $ehpp_\Gamma(\alpha) \leq shpp_\Gamma(\alpha) \leq hpp_\Gamma(\alpha) \leq \theta^{(A^*)}(\alpha)$ and $epp_\Gamma(\alpha) \leq pp_\Gamma(\alpha) \leq \theta^{(A^*)}(\alpha)$, and $epp_\Gamma(\alpha) \leq ehpp_\Gamma(\alpha)$ and $pp_\Gamma(\alpha) \leq hpp_\Gamma(\alpha)$.
3. $x_\Gamma(\alpha) = \sup\{x_D(\alpha) : D \in \Gamma\}$ for $x \in \{pp, epp, hpp, ehpp, spp, shpp\}$.
4. $x_D(\alpha) = \sup\{x_D(\beta) : \theta(\text{Dom}(D)) \leq \beta < \alpha\}$ for $x \in \{hpp, ehpp, shpp\}$ if $\theta(\text{Dom}(D)) \leq \text{cf}(\alpha)$.
5. If $a \neq b \Rightarrow \alpha_a \neq \alpha_b$ then $\text{cf}_D(\langle \alpha_a : a \in A^* \rangle) \geq \min_{A \in D} |A|$.

Remark 2.2A.1 If $hpp_\Gamma(\alpha) > \beta$, and $\text{cf}(\beta) \geq \theta(\mathcal{P}(\text{Dom}(D)))$ and relevant criterion for existence of $<_D$ -*eub* holds for $D \in \Gamma$ then for some $\alpha' \leq \alpha$ we have $pp_\Gamma(\alpha') > \beta$. See below. 2) In fact $shpp_\Gamma(\alpha) = hpp_\Gamma(\alpha)$ and $spp_\Gamma(\alpha) = pp_\Gamma(\alpha)$ by 1.9(4); if α is a limit ordinal (hence $x_\Gamma(\alpha)$ is a limit ordinal for x as in 2.2(3)).

Proof. Easy, e.g.

5) let $C \subseteq \bigcup_{a \in A^*} \alpha_a$ exemplify $\text{cf}_D(\langle \alpha_a : a \in A^* \rangle) = \text{otp}(C)$. Let

$$A' = \{a \in A^* : C_a \neq a\} \quad \text{where}$$

$$C_a = \{\beta \in C : \beta \leq \alpha_a \text{ but for no } b \in A^* \text{ do we have } \beta \leq \alpha_b < \alpha_a\}.$$

So there is a one-to-one function from A' into C (simply $a \rightarrow \min(C_a)$) and

$$\text{otp}\{\alpha_a : a \in A'\} < \text{otp}\{\alpha_a : a \in A'\} + \omega,$$

so we can finish.

Definition 2.3

1. I has the true cofinality δ , δ an ordinal if there is a cofinal $J \subseteq I$ and a function h from J onto δ such that $h(f_1) < h(f_2) \Rightarrow f_1 <_I f_2$.
2. I has strict cofinality δ , δ an ordinal if the function h above is one-to-one.

Claim 2.4 1. If I has the strict cofinality δ then I has the true cofinality δ and $\text{cf}(I) \leq \delta$.

2. If I has the true (strict) cofinality δ then I has the true (strict) cofinality $\text{cf}(\delta)$ (which is regular).

3. If $\bar{F} = \langle F_\alpha : \alpha < \delta \rangle$, $\bigcup_{\alpha < \delta} F_\alpha \subseteq {}^{(A^*)}\text{Ord}$, \bar{F} is $<_D$ -increasing and \bar{F} has $<_D$ -eub $\langle \alpha_a : a \in A^* \rangle$ then $\prod_{a \in A^*} \alpha_a / D$ has the true cofinality $\text{cf}(\delta)$.

4. If I has the true cofinality δ_1 and δ_2 then $\text{cf}(\delta_1) = \text{cf}(\delta_2)$.

Remark 2.4A. We did not say “ I has the true cofinality $\lambda \Rightarrow \text{cf}(I) \leq \lambda$ ”. But it is true that: I has true cofinality λ implies $\text{cf}^*(I) \leq^* \lambda$.

Claim 2.5 Let Γ, A^* be as in 2.1(1).

1. $[|A^*| \text{ an aleph, } DC_{|\mathcal{P}(A^*)|}^+]$ Assume $\delta < \text{hpp}_\Gamma(\alpha)$, δ a cardinal and $\text{cf}(\delta) \geq \theta(\mathcal{P}(A^*))$. Then for some $D \in \Gamma$ and $\bar{\alpha} = \langle \alpha_a : a \in A^* \rangle$, with each α_a being a limit ordinal $\leq \alpha$ we have, $\prod_{a \in A^*} \alpha_a / D$ has the true cofinality δ .

2. $[AC_{A^*}]$ If $\prod_{a \in A^*} \alpha_a / D$ has the true cofinality μ then $\prod_{a \in A^*} \text{cf}(\alpha_a) / D$ also has the true cofinality μ .

3. $[|A^*| \text{ an aleph and } DC_{|\mathcal{P}(A^*)|}^+]$ Assume $\omega \leq \alpha < \aleph_\alpha$ and $\aleph_\alpha > \theta(\mathcal{P}(A^*))$. If $\aleph_\gamma \leq \text{hpp}_D(\aleph_\alpha)$ then ${}^{(A^*)}\alpha$ can be mapped onto γ . Similarly for $\aleph_{\alpha_0 + \gamma} \leq \text{pp}_D(\aleph_{\alpha_0 + \alpha})$.

4. Similar claims hold for pp , $shpp$, spp , $ehpp$ and epp .

Proof. 1) By the definition of $\text{hpp}_\Gamma(\alpha)$ we can find $D \in \Gamma$ and a $<_D$ -increasing sequence $\bar{F} = \langle F_\alpha : \alpha < \delta \rangle$, with $F_\alpha \subseteq {}^{(A^*)}\alpha$ non empty.

If $g \in {}^{(A^*)}(\alpha + 1)$ is a $<_D$ -eub of \bar{F} then by 2.4(3), $\alpha_a =: g(a)$ for $a \in A^*$ are as required. Now such g exists by 1.5 which is applicable by 1.7A(1).

2) Let $\bar{F} = \langle F_\alpha : \alpha < \mu \rangle$ exemplify that $\prod_{a \in A^*} \alpha_a$ has true cofinality μ . By

AC_{A^*} we can find $\langle C_a : a \in A^* \rangle \in \mathbf{V}$, C_a a club of α_a of order type $\text{cf}(\alpha_a)$. For $f \in \prod_{a \in A^*} \alpha_a$ let $f^\otimes \in \prod_{a \in A^*} C_a$ be $f^\otimes(a) = \min(C_a \setminus f(a))$, so $f \leq f^\otimes \in \prod_{a \in A^*} C_a$

and $f_1 \leq_D f_2 \Rightarrow f_1^\otimes \leq_D f_2^\otimes$. Now apply 1.3(7) to $\langle \{f^\otimes : f \in F_\alpha\} : \alpha < \mu \rangle$.

3) It is enough to map ${}^{(A^*)}\alpha$ onto $(\gamma - 1) \setminus \alpha$. We define a function H from ${}^{(A^*)}\alpha$ into $\gamma \setminus \alpha$. For $f \in {}^{(A^*)}\alpha$, if $(\prod_{a \in A^*} \aleph_{f(a)}, <_D)$ has true cofinality

$\aleph_{\alpha + \beta + 1} \leq \aleph_\gamma$ then let $H(f) = \beta$. By AC_{A^*} and 2.5 (2) it is enough for every β such that $\aleph_\alpha < \aleph_{\beta + 1} < \aleph_\gamma$ to find $f \in {}^{(A^*)}(\aleph_\alpha)$ such that $(\prod_{a \in A^*} f(a), <_D)$ has

the true cofinality $\aleph_{\beta + 1}$. Now we use part (1) of 2.5.

4) By 2.2 or repeating the proofs. □_{2.5}

Claim 2.6 [$DC_{\aleph_0} + (\forall D \in \Gamma) AC_{\mathcal{P}(\text{Dom}(D))}$] In 2.5(1), if Γ is \aleph_1 -complete then the conclusion holds.

Proof. We can use 1.7 instead of 1.5.

Definition 2.7

$\text{pcf}_\Gamma\{\alpha_a : a \in A^*\} = \{\lambda : \lambda \text{ is the true cofinality of } \prod \alpha_a / D$
for some filter D on A^* which belongs to $\Gamma\}$.

Remark 2.7A. We could phrase 2.5 for pcf_Γ .

Claim 2.8 1. If λ is an aleph, $\mu < \lambda < \text{epp}_D(\mu)$ and $\theta(\text{Dom}(D)) < \lambda$ then λ is not measurable

2. If λ is an aleph, $\mu < \lambda < \text{epp}_D(\lambda)$,
then there is no $(\leq |\text{Dom}(D)| + \mu)$ -complete uniform ultrafilter on λ .

Proof. 1. Let (\bar{F}, D) witness $\lambda < \text{epp}_D(\mu)$, so $f \in {}_\alpha \bigcup F_\alpha \Rightarrow \text{Dom}(f) = A^*$, where $A^* =: \text{Dom}(D)$; so let $F = \{f_\alpha : \alpha < \lambda\}$, where $F_\alpha = \{f_\alpha\}$. Assume D^* is a λ -complete ultrafilter on λ . For each $a \in A^*$, we have a function $g_a : \lambda \rightarrow \mu$, defined by $g_a(\alpha) = f_\alpha(a)$. So g_a is a one-to-one map

from $s' =: \{\alpha < \lambda : (\forall \beta < \alpha)(g_a(\beta) \neq g_a(\alpha))\}$ onto $\text{Rang}(g_a) \subseteq \mu$,

so $|\text{Rang}(g_a)|$ is an aleph $\leq \mu$ (or finite). Hence $|\text{Rang}(g_a)| < \lambda$. By the choice of D^* for some unique γ_a we have $B_a =: \{\alpha < \lambda : f_\alpha(a) = \gamma_a\} \in D^*$, as $\langle f_\alpha : \alpha < \lambda \rangle$ exists also $\langle \gamma_a : a \in A^* \rangle$ exists as well as $\langle B_a : a \in A^* \rangle$. Clearly $|\{\gamma_a : a \in A^*\}| < \theta(A^*)$. As D^* is $(\leq |A^*|)$ -complete, $B^* =: \bigcap_{a \in A^*} B_a \in D^*$, but

$[\alpha, \beta \in B^* \Rightarrow f_\alpha = f_\beta]$, a contradiction.

2. Same proof. $\square_{2.8}$

Remark 2.8A. You can also phrase the theorem in terms of λ -complete filters on λ which are weakly κ -saturated (i.e. for every $h : \lambda \rightarrow \lambda' < \lambda$, for some $C \subseteq \lambda'$ of cardinality $\leq \kappa$, $\{\alpha < \lambda : h(\alpha) \in C\} \in D$). Here instead of $|A^*| < \lambda$ we need: D is uniform and $\neg(\lambda \leq |^{(A^*)}\kappa|)$.

Definition 2.9 Let E be a set of filters on a set A^* .

1. E has the I -lub property (I a directed set) if:
for every $D_0 \in E$ and $F = \{f_t : t \in I\} \subseteq {}^{(A^*)}\text{Ord}$ such that $\{f_t : t \in I\}$ is $<_{D_0}$ -increasing,
there is $D, D_0 \subseteq D \in E$ such that there is a $<_D$ -lub for F .
2. “ E has the I -eub property” is defined similarly. The I -*eub is defined similarly for $<_D$ -increasing $\bar{F} = \langle F_t : t \in I \rangle$. Similarly I -*lub.
3. If $E = \{D\}$ we shall say D has this property.

3 Nice families of filters

Hypothesis 3.0: E is a family of \aleph_1 -complete filters on $A^* = \text{Dom}(E)$, such that (for simplicity) $D \in E$ & $A \in D^+ \Rightarrow D + A \in E$. Our main interest is in nice E (defined below).

Definition 3.1 We call E nice if

$$\oplus_E \quad (\forall f : A^* \rightarrow \text{Ord})(\forall D \in E)(\text{rk}_D^2(f) < \infty)$$

(see below). Let \oplus_{A^*} mean \oplus_E for some E , with $A^* = \text{Dom}(E)$ and $\oplus_\kappa[E] = \oplus_E$ with $A^* = \kappa$.

$$\text{Let } E_{[D_0]} = \{D : D_0 \subseteq D \in E\}.$$

Definition 3.2

1. The truth value of $\text{rk}_D^2(f, E) \leq \alpha$ (for α an ordinal $f : A^* \rightarrow \text{Ord}$, E usually omitted):
 $\text{rk}_D^2(f) \leq \alpha$ if for every $A \in D^+$ and $f_1 <_{D+A} f$ (and $f_1 : A^* \rightarrow \text{Ord}$) there is $D_1, D + A \subseteq D_1 \in E$ and $\beta < \alpha$ such that $\text{rk}_{D_1}^2(f_1) \leq \beta$.
 (So $f =_D \mathbf{0}_{A^*}$ implies $\text{rk}^2(f) \leq 0$)
2. $\text{rk}_D^2(f) = \alpha$ if $\text{rk}_D^2(f) \leq \alpha$ and $\neg[\text{rk}_D^2(f) \leq \beta]$ for $\beta < \alpha$
 $\text{rk}_D^2(f) = \infty$ if $\text{rk}_D^2(f) \leq \alpha$ for no α
 $\text{rk}_D^3(f) = \min\{\text{rk}_{D_1}^2(f) : D \subseteq D_1 \in E\}$
 (really we should have written $\text{rk}_D^2(f, E)$ etc.).

Remark. Why start with rk^2 ? To be consistent with [Sh-g] Ch.V.

Convention 3.3 Let f, g vary on $(A^*)\text{Ord}$ and $D \in E$ and $A, B \subseteq A^*$.

- Claim 3.4**
1. $\text{rk}_D^2(f) \leq \beta, \beta \leq \alpha$ implies $\text{rk}_D^2(f) \leq \alpha$; so $\text{rk}_D^2(f)$ is well defined as an ordinal or ∞ (ZF is enough for the definition).
 2. If $f \leq g$ or just $f \leq_D g$ and $l = 2, 3$ then $\text{rk}_D^l(f) \leq \text{rk}_D^l(g)$ (so $f =_D g$ implies $\text{rk}_D^l(f) = \text{rk}_D^l(g)$).
 3. In 3.2(1) we can demand in addition $f_1 \leq f$.
 4. $\text{rk}_D^3(f) \leq \text{rk}_D^2(f)$.
 5. If $D_1 \subseteq D_2$ then $\text{rk}_{D_1}^3(f) \leq \text{rk}_{D_2}^3(f)$.
 6. For every f, D , for some D_1 such that $D \subseteq D_1 \in E$ we have

$$\text{rk}_D^3(f) = \text{rk}_{D_1}^2(f) = \text{rk}_{D_1}^3(f).$$

7. $\text{rk}_D^2(f) = \sup\{\text{rk}_{D+A}^3(g) + 1 : A \in D^+ \text{ and } g <_{D+A} f\}$.
8. If $A \in D^+$ then $\text{rk}_D^3(f) \leq \text{rk}_{D+A}^3(f) \leq \text{rk}_{D+A}^2(f) \leq \text{rk}_D^2(f)$.
 [Why? By parts (5), (4), (7)+(4) respectively.]
9. If $\text{rk}_D^2(f) = \text{rk}_D^3(f)$ then for every $A \in D^+$

$$\text{rk}_{D+A}^2(f) = \text{rk}_{D+A}^3(f) = \text{rk}_D^2(f) = \text{rk}_D^3(f).$$

10. If $f <_D g$ then $\text{rk}_D^3(f) < \text{rk}_D^3(g)$ or both are ∞ .

[Why? Apply part (6) to g , D and get D_1 . Now

$$\begin{aligned} \text{rk}_D^3(f) &\leq \text{rk}_{D_1}^3(f) < \text{rk}_{D_1}^3(g) + 1 \\ &\leq \sup\{\text{rk}_{D_1+A}^3(h) + 1 : h <_{D_1+A} g \text{ and } A \in D_1^+\} \\ &= \text{rk}_{D_1}^2(g) = \text{rk}_{D_1}^3(g). \end{aligned}$$

(Why the inequalities? by $D \subseteq D_1$, using part (5); trivially; as $f <_D g$ hence $f <_{D_1} g$; by definition of rk^2 ; and by the choice of D_1 respectively.)]

11. $\|f\|_D \leq \text{rk}_D^3(f)$ ($\|\cdot\|_D$ – the Galvin–Hajnal rank which is defined by: $\|f\|_D = \sup\{\|g\|_D + 1 : g <_D f\}$).

[Why? part (10).]

12. If for $l = 1, 2$, $\text{rk}_D^2(f_l) = \text{rk}_D^3(f_l) = \alpha_l$ and $\alpha_1 < \alpha_2$ then $f_1 <_D f_2$.

[Why? If not then for some $A \in D^+$ we have $f_2 \leq_{D+A} f_1$, so by part (2) we have $\text{rk}_{D+A}^2(f_2) \leq \text{rk}_{D+A}^2(f_1)$, but by part (9) $\text{rk}_{D+A}^2(f_2) = \alpha_2 > \alpha_1 = \text{rk}_{D+A}^2(f_1)$, a contradiction.]

13. If for $l = 1, 2$, $\text{rk}_D^2(f_l) = \text{rk}_D^3(f_l) = \alpha_l$ and $\alpha_1 = \alpha_2$ then $f_1 =_D f_2$.

[Why? If not then by symmetry for some $A \in D^+$ we have $f_1 <_{D+A} f_2$, so by part (10) we have $\text{rk}_{D+A}^3(f_1) < \text{rk}_{D+A}^3(f_2)$ and by part (9) we get a contradiction.]

14. If $\text{rk}_{D_0}^2(f) < \infty$ and $\neg(g \geq_{D_0} f)$ then for some $D_1 \in E_{[D_0]}$, $\text{rk}_{D_1}^2(g) < \text{rk}_{D_0}^2(f) < \infty$. [Why? Clearly for some $A \in D_0$, $g <_{D_0+A} f$; now for some D_1 , $D_0 + A \subseteq D_1 \in E$ and $\text{rk}_{D_1}^2(g) = \text{rk}_{D_1}^3(g) = \text{rk}_{D_0+A}^3(g)$, but $\text{rk}_{D_0+A}^3(g) < \text{rk}_{D_0+A}^3(f) \leq \text{rk}_{D_0+A}^2(f) \leq \text{rk}_{D_0}^2(f) < \infty$.]

15. If $\{A_t : t \in I\} \subseteq D^+$ and $(\forall D_1)(D \subseteq D_1 \in E \rightarrow (\exists t \in I)(A_t \in D_1^+))$ (e.g. I finite, $\bigcup_{t \in I} A_t \in D$) then

$$\text{rk}_D^2(f) = \sup\{\text{rk}_{D+A_t}^2(f) : t \in I\} \text{ and } \text{rk}_D^3(f) = \min\{\text{rk}_{D+A_t}^3(f) : t \in I\}$$

[Why? The inequalities \geq, \leq by part (8), the other by the definition (or part (7) and part (9)).]

Observation 3.5 1. (a) If $f = g + 1$, $D \in E$ then $\text{rk}_D^2(f) \leq \text{rk}_D^2(g) + 1$; if in addition $\text{rk}_D^3(g) = \text{rk}_D^2(g)$ then $\text{rk}_D^2(f) = \text{rk}_D^3(g) + 1$.

(b) If every $f(a)$ is a limit ordinal then $\text{rk}_D^3(f)$ is a limit ordinal.

(c) $f =_D \mathbf{0}_{A^*}$ iff $\text{rk}_D^2(f) = 0$.

(d) $\text{rk}_D^3(f) = 0$ iff $\neg(0_{A^*} <_D f)$.

2. If $f = g + 1$ then:

(a) $\text{rk}_D^2(f) \in \{\text{rk}_D^2(g), \text{rk}_D^2(g) + 1\}$,

(b) if $\text{rk}_D^2(f) = \text{rk}_D^2(g) < \infty$ then $\text{rk}_D^2(g)$ is a limit ordinal of cofinality $< \theta(\mathcal{P}(A^*))$ and even $< \theta(\mathcal{P}(A^*)/D)$.

3. If $\text{Rang}(f)$ is a set of limit ordinals, $F \subseteq \prod_{a \in A^*} f(a)$ is such that

$$[g < f \Rightarrow (\exists h \in F)(g \leq h)]$$

then $\text{rk}_D^2(f) = \sup\{\text{rk}_{D+A}^3(h) : h \in F, A \in D^+\}$.

4. If $\delta = \text{rk}_D^2(f)$, $\text{cf}(\delta) \geq \theta(E)$ then² for some D_1 , $D \subseteq D_1 \in E$, $\prod_{a \in A^*} f(a)/D_1$ has true cofinality δ (moreover by a smooth sequence $\langle F_\alpha : \alpha < \text{cf}(\delta) \rangle$)
5. If $\text{rk}_D^2(f) = \alpha < \infty$ then for every $\beta < \alpha$ there are D , g such that $D \subseteq D_1 \in E$, $g <_{D_1} f$ and $\text{rk}_{D_1}^2(g) = \text{rk}_{D_1}^3(g) = \beta$.

Proof. 1)(a) By 3.4(7), 3.4(2), 3.4(8) and common sense respectively

$$\begin{aligned} \text{rk}_D^2(f) &= \sup\{\text{rk}_{D+A}^3(g') + 1 : A \in D^+ \text{ and } g' <_{D+A} f\} \\ &\leq \sup\{\text{rk}_{D+A}^3(g) + 1 : A \in D^+\} \\ &\leq \sup\{\text{rk}_D^2(g) + 1\} = \text{rk}_D^2(g) + 1 \end{aligned}$$

and, for the second phrase, by 3.4(4) and 3.4(10) respectively

$$\text{rk}_D^2(f) \geq \text{rk}_D^3(f) \geq \text{rk}^3(g) + 1.$$

Together we have finished.

(b) By part (c) proved below, $\text{rk}_D^3(f) > 0$, so assume toward contradiction that $\text{rk}_D^3(f) = \alpha + 1$, so (by 3.4(6)) for some D_1 we have $D \subseteq D_1 (\in E)$ and $\text{rk}_D^3(f) = \text{rk}_{D_1}^2(f) = \text{rk}_{D_1}^3(f)$. By 3.4(7) for some $A \in D^+$ and $g <_{D_1+A} f$ we have $\text{rk}_{D_1+A}^3(g) = \alpha$. By 3.4(10), 3.4(10), 3.4(9) and the choice of α respectively

$$\text{rk}_{D_1+A}^3(g) < \text{rk}_{D_1+A}^3(g+1) < \text{rk}_{D_1+A}^3(f) = \text{rk}_D^3(f) = \alpha + 1.$$

So $\text{rk}_{D_1+A}^3(g) < \alpha$ contradicting the choice of A , and g .

(c) If $f \equiv_D 0_{A^*}$ then the supremum in 3.4(7) (or the definition) is taken on an empty set so $\text{rk}_D^2(f) = 0$. If $\neg(f \equiv_D 0_{A^*})$ then $A = \{a \in A^* : f(a) > 0\}$ and $g \in {}^{(A^*)}\text{Ord}$, $g(a) = 0$ appears in the supremum in 3.4(7) (or the definition) so $\text{rk}_D^2(f) \geq 0 + 1 = 1 > 0$.

(d) By (c) and the definition of rk^3 .

2) By 3.4(2), 3.5(a) respectively we have

$$\text{rk}_D^2(g) \leq \text{rk}_D^2(f) \leq \text{rk}_D^2(g) + 1.$$

Thus clause (a) holds. For clause (b) assume $\text{rk}_D^2(f) = \text{rk}_D^2(g)$, and call this ordinal α . If $\alpha = 0$ we get contradiction by 3.5(1)(c) as $\text{rk}_D^2(f) = \alpha$, $f = g + 1$. If $\alpha = \beta + 1$ then for some $A \in D^+$ and $g_1 <_{D+A} g$ we have $\text{rk}_{D+A}^3(g_1) = \beta$ so by 3.4(7), 3.4(10), and the choice of g_1 , A , and the choice of β respectively:

$$\text{rk}_D^2(f) \geq \text{rk}_{D+A}^3(g) + 1 > \text{rk}_{D+A}^3(g_1) + 1 = \beta + 1 = \alpha;$$

contradiction. So α is a limit ordinal and $\{\text{rk}_{D+A}^3(g) : A \in D^+\}$ is an unbounded subset of α hence $\text{cf}(\alpha) < \theta(\mathcal{P}(A^*))$ (in fact $\text{cf}(\alpha) < \theta(\mathcal{P}(A^*)/D)$).

3) By 3.4(7) and 3.4(2).

4) By 3.4(7), and $\text{rk}_D^2(f)$ being a limit ordinal

² We can use this to prove a result parallel to 2.5 (3)

$$\begin{aligned}
\delta = \text{rk}_D^2(f) &= \sup\{\text{rk}_{D+A}^3(g) + 1 : A \in D^+ \text{ and } g <_{D+A} f\} \\
&= \sup\{\text{rk}_{D+A}^3(g) : A \in D^+ \text{ and } g <_{D+A} f \text{ and } g \leq f\} \\
&= \sup\{\text{rk}_{D_1}^3(g) : \text{for some } A \in D^+, g <_{D+A} f \text{ and } g \leq f \\
&\quad \text{and } D + A \subseteq D_1 \in E \text{ and} \\
&\quad \text{rk}_{D_1}^3(g) = \text{rk}_{D_1}^2(g) < \delta\} \\
&= \sup\{\text{rk}_{D_1}^3(g) : D \subseteq D_1 \in E, g <_{D_1} f \text{ and } g \leq f \\
&\quad \text{and } \text{rk}_{D_1}^3(g) = \text{rk}_{D_1}^2(g) < \delta\} \\
&= \sup_{D_1 \in E, D_1 \supseteq D} \sup\{\text{rk}_{D_1}^3(g) : g <_{D_1} f \text{ and } g \leq f \\
&\quad \text{and } \text{rk}_{D_1}^3(g) = \text{rk}_{D_1}^2(g) < \delta\}.
\end{aligned}$$

As $\text{cf}(\delta) \geq \theta(E)$ necessarily for some D_1 we have $D \subseteq D_1 \in E$ and

$$\delta = \sup\{\text{rk}_{D_1}^3(g) : g <_{D_1} f \text{ and } g \leq f \text{ and } \text{rk}_{D_1}^3(g) = \text{rk}_{D_1}^2(g) < \delta\}.$$

For $\alpha < \delta$ let

$$F_\alpha = \{g : g \leq f, g <_{D_1} f, \text{ and } \alpha = \text{rk}_{D_1}^3(g) = \text{rk}_{D_1}^2(g)\}$$

and $S = \{\alpha < \delta : F_\alpha \neq \emptyset\}$, it is necessarily unbounded in δ . Now by 3.4(12), (13), $\bar{F} = \langle F_\alpha : \alpha \in S \rangle$ is $<_{D_1}$ -increasing and smooth.

5) Suppose $\beta < \alpha, f, D$ form a counterexample. Then we prove by induction on $\gamma \geq \beta$ that

(*) $_\gamma$ if $g \in {}^{(A^*)}\text{Ord}$, and $D \subseteq D_1 \in E$ and $g <_{D_1} f$ and $\text{rk}_{D_1}^3(g) \geq \beta$ then $\text{rk}_D^3(g) \geq \gamma$.

For $\gamma = \beta$ clearly (*) $_\gamma$ holds trivially.

For $\gamma > \beta$, we can find D_2 such that $D_1 \subseteq D_2 \in E$ and $\text{rk}_{D_1}^3(g) = \text{rk}_{D_2}^2(g) = \text{rk}_{D_2}^3(g)$, but $\beta \leq \text{rk}_{D_1}^3(g)$ by assumption, so as β, α, f, D form a counterexample $\beta < \text{rk}_{D_1}^3(g)$ so by 3.4(7) applied to $\text{rk}_{D_2}^2(g)$ we can find $A \in D_2^+$ and $g_1 <_{D_2+A} g$ such that $\text{rk}_{D_2+A}^3(g_1) \geq \beta$, so by the induction hypothesis $\text{rk}_{D_1+A}^3(g_1) \geq \bigcup\{\gamma_1 : \beta \leq \gamma_1 < \gamma\}$. Now by 3.4(9) we know $\text{rk}_{D_2+A}^3(g) = \text{rk}_{D_2}^3(g)$ and by 3.4(10), $\text{rk}_{D_2+A}^3(g_1) < \text{rk}_{D_2+A}^3(g)$. Together we get the required conclusion in (*). So $\text{rk}_D^2(f) = \sup\{\text{rk}_{D+A}^3(g) + 1 : A \in D^+ \text{ and } g <_{D+A} f\}$ is α hence $> \beta$ so for some $A \in D^+$ and $g <_{D+A} f$ we have $\text{rk}_{D+A}^3(g) \geq \beta$ hence by (*) it is $> \alpha$, contradiction. $\square_{3.5}$

Definition 3.6 For D a filter on A^* and $f : A^* \rightarrow \text{Ord}$ let

$$\mathbf{T}_D(f) = \{F : F \subseteq \prod_{i \in A^*} f(i) \text{ satisfies } f_1 \in F \ \& \ f_2 \in F \ \& \ f_1 \neq f_2 \Rightarrow f_1 \neq_D f_2\}$$

$$\mathbf{Tw}_D(f) = \{(F, \mathbf{e}) : F \subseteq \prod_{i \in A^*} f(i), \mathbf{e} \text{ is an equivalence relation on } F, \\ \neg f_1 \mathbf{e} f_2 \Rightarrow f_1 \neq_D f_2\},$$

$$\mathbf{T}_D(f) = \{(F, \mathbf{e}) \in \mathbf{Tw}_D(f) : \mathbf{e} \text{ is } =_D \upharpoonright F\},$$

$$\text{Ts}_D(f) = \sup\{|F| : F \in \mathbf{T}_D(f)\},$$

$$Tw_D(f) = \sup\{|F/\mathbf{e}| : F/\mathbf{e} \in \mathbf{Tw}_D(f)\},$$

$$T_D(f) = \sup\{|F/\mathbf{e}| : (F, \mathbf{e}) \in \mathbf{T}_D(f)\}$$

(it is a kind of cardinality; of course the sup do not necessarily exist). We may write $F/\mathbf{e} \in \mathbf{Tw}_D(f)$ instead of $(F, \mathbf{e}) \in \mathbf{Tw}_D(f)$ and also may write $\bar{F} = \langle F_t : t \in I \rangle$ for (F, \mathbf{e}) if $F = \bigcup_t F_t$, and $f \mathbf{e} g \leftrightarrow (\exists t)[\{f, g\} \subseteq F_t]$ and so the F_t 's are pairwise disjoint.

Observation 3.7 1. $F \in \mathbf{Ts}_D(f) \Rightarrow (F, =) \in \mathbf{Tw}_D(f)$.

2. $Ts_D(f) \leq Tw_D(f)$.

3. If $\bar{F} = \langle F_\alpha : \alpha < \alpha^* \rangle$ is $<_D$ -increasing, $\bigcup F_\alpha \subseteq \prod_{a \in A^*} f^*(a)$ then $\text{rk}_D^3(f^*) \geq \alpha^*$; so $\text{hpp}_D(f^*) \leq \text{rk}_D^3(f^*)$; also $\langle F_\alpha : \alpha < \alpha^* \rangle \in \mathbf{Tw}_D(f)$ hence $|\alpha^*| \in Tw_D(f)$.

4. $[AC_{A^*}]$ If $f, g \in {}^{A^*}\text{Ord}$ and D is a filter on A^* and $\{a \in A^* : f(a) = g(a)\} \in D$ and $X \in \{Ts, Tw, T\}$ then $X_D(f) = X_D(g)$.

Claim 3.8 1. $[E \text{ is nice}]$ Assume $F \subseteq ({}^{A^*}\text{Ord})$. We can find $\langle (F_D, h_D) : D \in E \rangle$ such that:

(a) $F = \bigcup_{D \in E} F_D$

(b) $h_D : F_D \rightarrow \text{Ord}$

(c) if $f_1, f_2 \in F_D$ then $h_D(f_1) \leq h_D(f_2) \iff f_1 \leq_D f_2$ and hence

$$h_D(f_1) = h_D(f_2) \iff f_1 =_D f_2, \text{ so}$$

$$h_D(f_1) < h_D(f_2) \iff f_1 <_D f_2$$

(d) if $F \subseteq \prod_{i \in A^*} f(i)$ then $(F_D, =_D) \in \mathbf{Tw}(f)$.

2. Instead of "E is nice" it suffices that $F \subseteq \prod_{a \in A^*} f^*(a)$, $\text{rk}_D^2(f^*) < \infty$.

Proof. 1) Let $F_D = \{f \in F : \text{rk}_D^2(f) = \text{rk}_D^3(f) (< \infty)\}$, $h_D(f) = \text{rk}_D^2(f)$. Now clause (a) holds by 3.4 (6), clause (b) holds as E is nice (see 3.1). For (c): it holds by 3.4(12), 3.4(13). Last clause (d) follows from (c).

2) Similar (see 3.4 (14)).

□_{3.8}

Conclusion 3.9 Let $D_0 \in E$.

1. $[E \text{ is nice}]$ Assume $F \subseteq \prod_{a \in A^*} f^*(a)$ is in $\mathbf{Ts}_{D_0}(f^*)$.

(a) If $|E|$ is an aleph then so is $|F|$.

(b) F can be represented as $\bigcup \{F_D : D_0 \subseteq D \in E\}$ such that F_D 's cardinality is an aleph $< \text{ehpp}_E(f^*)$.

(c) $|F| \leq^* |E| \times \text{ehpp}_E(f^*)$

2. $[E \text{ is nice}]$ Assume (F, \mathbf{e}) is in $\mathbf{Tw}_{D_0}(f^*)$

(any $(F, =_D)$ will do for $F \subseteq ({}^{A^*}\text{Ord})$).

(a) If $|E|$ is an aleph then so is $|F/\mathbf{e}|$.

- (b) F/\mathbf{e} can be represented as $\bigcup\{F_D/\mathbf{e} : D_0 \subseteq D \in E\}$ such that $|F/\mathbf{e}|$ is an aleph $\leq hpp(f^*)$.
- (c) $|F/\mathbf{e}| \leq^* |E| \times hpp_E(f^*)$
3. Instead of “ E is nice”, “ $\text{rk}_{D_0}^2(f^*) < \infty$ ” suffices.

Proof. 1) It follows from 2) as in this case \mathbf{e} is the equality.

2) For $D \in E_{[D_0]}$, let

$$F_D = \{f \in F : \text{rk}_D^2(f) = \text{rk}_D^3(f) \text{ and: } \begin{array}{l} \text{if } f_1 \in f/\mathbf{e} \text{ and } \text{rk}_D^2(f_1) = \text{rk}_D^3(f_1) \\ \text{then } \text{rk}_D^2(f_1) \geq \text{rk}_D^2(f) \}. \end{array}$$

Let $h_D : F_D \rightarrow \text{Ord}$ be $h_D(f) = \text{rk}_D^2(f)$. So (for $f_1, f_2 \in F_D$):

$$h_D(f_1) = h_D(f_2) \text{ iff } f_1/\mathbf{e} = f_2/\mathbf{e}.$$

Also $F/\mathbf{e} = \bigcup\{F_D/\mathbf{e} : D \in E\}$. So clause (b) holds, for clause (c) let $G(D, \alpha)$ be y iff y has the form f/\mathbf{e} where $f \in F_D$, $h_D(f) = \alpha$; so G is a partial function from $|E| \times hpp_E(f^*)$ onto F/\mathbf{e} . Note that clause (a) follows as $|X| \leq^* |Y|$, $|Y|$ is an aleph implies $|X|$ is an aleph. (We can choose a well ordering $<$ of E , we can let $F'_D = \{f : f \in F_D \text{ and letting } f \in F_t, \text{ for no } f' \in f/\mathbf{e} \text{ and } D' < D \text{ do we have } f' \in F_{D'}\}$. Let $h = \bigcup\{h_D \upharpoonright F'_D : D \in E\}$ so letting $h(f/\mathbf{e}) = h_D(f)$ for $f \in F'_D$, clearly h is one-to-one function from F/\mathbf{e} onto a set of ordinals.)

3) Similar proof (remember 3.4 (13)). □_{3.9}

Claim 3.10 1. If F is in $\mathbf{Ts}_D(f)$ and $\text{rk}_D^3(f) = \alpha < \infty$ then $|F| \leq^* |\alpha| \times |E|$.

2. Assume $(F, \mathbf{e}) \in \mathbf{Tw}_D(f)$ and $\text{rk}_D^3(f) = \alpha < \infty$. Then $|F/\mathbf{e}| \leq^* |\alpha| \times |E|$.

Proof. Included in the proof of 3.9.

Claim 3.11 If $\alpha = \text{rk}_{D_0}^2(f^*)$, $D_0 \in E$ then we can find $\langle A_D : D \in E_{[D_0]} \rangle$, $\alpha = \bigcup\{A_D : D \in E_{[D_0]}\}$, and $\langle (F_D, h_D) : D \in E_{[D_0]} \rangle$ with (F_D, h_D) as in 3.8, $F_D \subseteq \prod_{a \in A^*} f^*(a)$ with $\text{Rang}(h_D) = A_D$ and

$$h_D(f) = \alpha \Rightarrow \text{rk}_D^2(f) = \text{rk}_D^3(f) = \alpha.$$

Proof. Because there is “no hole in the possible ranks”. I.e. we apply 3.8 to $F = \prod_{a \in A^*} (f^*(a) + 1)$, and get $\langle (F_D, h_D) : D \in E_{[D_0]} \rangle$. Our main problem is that

some $\beta < \alpha$ is not in $\bigcup_{D \in E_{[D_0]}} \text{Rang}(h_D)$. Then use 3.5(5). □_{3.11}

Conclusion 3.12 Assume E is nice and for simplicity $|E|$ is an aleph, $f \in {}^{(A^*)}\text{Ord}$ and $(\forall a \in \text{Dom}(E)) (f(a) \geq |E|)$.

1. Then the cardinals

$$\sup_{D \in E_{[D_0]}} |\text{rk}_D^2(f)|, \sup_{D \in E_{[D_0]}} (T_D(f)), \sup_{D \in E_{[D_0]}} (Tw_D(f)) \text{ and } \sup_{D \in E_{[D_0]}} \{hpp_D(f)\}$$

are equal.

2. Assume AC_{A^*} . If f_1, f_2 are as in the assumption and $\{a \in A^* : |f_1(a)| = |f_2(a)|\} \in D$ then

$$\sup_{D \in E_{|D_0|}} |\text{rk}_D^2(f_1)|, \quad \sup_{D \in E_{D_0}} |\text{rk}_D^2(f_2)|$$

are the same cardinal.

3. Assume AC_{A^*} . Then $\text{rk}_{D_0}^2(f^+) \leq \sup_{D \in E_{|D_0|}} |\text{rk}_D^2(f)|^+$ where $f^+ \in {}^{(A^*)}\text{Ord}$ is $f^+(a) =: |f(a)|^+$

Proof. 1)

Step A

$$(*) \quad \sup_{D \in E_{|D_0|}} (\text{rk}_D^2(f) + 1) \geq \text{hpp}_{E_{|D_0|}}(f)$$

If $\beta < \text{hpp}_{E_{|D_0|}}(f)$ then we can find $D \in E_{|D_0|}$ and $\bar{F} = \langle F_\alpha : \alpha < \beta \rangle$ such that $F_\alpha \subseteq \prod_{a \in A^*} f(a)$ non empty and \bar{F} is $<_D$ -increasing. We can prove using 3.4(10) by induction on $\alpha < \beta$ that

$$(**) \quad g \in F_\alpha \Rightarrow \text{rk}_D^3(f) \geq \alpha.$$

Now by 3.4(7) we have $\text{rk}_D^2(f) > \alpha$ and $\alpha < \beta$ so $\text{rk}_D^2(f) \geq \beta$.

Step B

$\text{rk}_D^2(f)$ can be represented as the union of E sets each of order type $< \text{hpp}_{E_{|D_0|}}(f)$.
By 3.10.

Step C

$$\sup_{D \in E_{|D_0|}} |\text{rk}_D^2(f)| = \text{hpp}_{E_{|D_0|}}(f).$$

Why? By steps A, B as $|E|$ is an aleph and $f(a) \geq |E|$ for every a .

Step D

$$\text{hpp}_{E_{|D_0|}}(f) \leq T_D(f) \leq Tw_D(f).$$

By definition (true for each D separately).

Step E

$$\sup_{D \in E_{|D_0|}} T_D(f) \leq \sup_{D \in E_{|D_0|}} \text{rk}_D^2(f).$$

Why? By 3.8

- 2) By part (1) and 3.7(4).

- 3) By part (2) and the definition of $\text{rk}_{D_0}^2$. □_{3.12}

Remark 3.12A. If we waive “ $|E|$ is an aleph” but still assume $f(a) \geq \theta(E)$ we get that those three cardinals are not too far.

Claim 3.13 *Assume*

- (A) λ is an aleph, $\langle \lambda_i : i < \delta \rangle$ is a strictly increasing continuous sequence of alephs with limit λ ,

(B) there is $G^* = \bigcup_{i < \delta} G_i$, G_i a one-to-one function from $\prod_{j < i} \lambda_j$ into some $\lambda_i^* < \lambda_{i+1}$ (so if AC_δ , we need just that each G_i exists, i.e. $|\prod_{j < i} \lambda_j| < \lambda_{i+1}$).

Then there is $F \in \mathbf{Ts}_{D_\delta^{bd}}(\langle \lambda_i^* : i < \delta \rangle)$, and $|F| = \prod_{i < \delta} \lambda_i$.

(D_δ^{bd} is the filter generated by the cobounded sets of δ).

Remark. This goes back to Galvin, Hajnal [GH], see [Sh 386, 5.2A(1)].

Proof. Define a function $G : \prod_{i < \delta} \lambda_i \longrightarrow \prod_{i < \delta} \lambda_i^*$ by

$G(f)(i) = G_i(f \upharpoonright i)$ and let $F = \text{Rang}(G)$ □_{3.13}

Claim 3.14 Assume that $D_0 \in E$ and $f_\alpha \in {}^{(A^*)}\text{Ord}$ for $\alpha < \alpha^*$.

1. [E is nice] There are $g \in {}^{(A^*)}\text{Ord}$ and D such that:
 - (a) $D_0 \subseteq D \in E$,
 - (b) $\text{rk}_D^2(g) = \text{rk}_D^3(g)$,
 - (c) g is a $<_D$ -lub of $\{f_\alpha : \alpha < \alpha^*\}$, i.e. :
 - (i) $\alpha < \alpha^* \Rightarrow f_\alpha \leq_D g$
 - (ii) if $g' \in {}^{(A^*)}\text{Ord}$ and $f_\alpha \leq_D g'$ for all $\alpha < \alpha^*$ then $g \leq_D g'$.
 - (d) Moreover for every $A \in D^+$, g is also a $<_{D+A}$ -lub of $\{f_\alpha : \alpha < \alpha^*\}$.
2. If $f_\alpha \leq_{D_0} f^*$ for $\alpha < \alpha^*$, $\text{rk}_{D_0}^2(B^*) < \infty$ then there are $g \in {}^{(A^*)}\text{Ord}$, D satisfying (a)-(d) and we can add
 - (e) $g \leq f^*$.

Proof. 1) Follows by (2), just let $f^*(a) = \sup_{\alpha < \alpha^*} f_\alpha(a)$.

2) Clearly the set

$$K = \{(D, g) : D_0 \subseteq D \in E \ \& \ (\forall \alpha < \alpha^*)(f_\alpha \leq_D g), \text{rk}_D^2(g) < \infty \text{ and } g \leq f^*\}$$

is not empty (because the pair (D_0, f^*) is in it). Choose among those pairs one (D_1, g) with $\text{rk}_{D_1}^3(g)$ minimal. So for some $D, D_1 \subseteq D \in E$ we have $\text{rk}_{D_1}^3(g) = \text{rk}_D^3(g) = \text{rk}_D^2(g)$ (see 3.4(6)), so also $(D, g) \in K$. Clearly (D, g) satisfies (a), (b), (e) and (c)(i). If (c)(ii) fails then let g' exemplify it and so

$$A = \{a \in \kappa : g'(a) < g(a)\} \neq \emptyset \text{ mod } D.$$

But clearly also $(D + A, g')$ is in the family and (see 3.4(10) and 3.4(9) respectively):

$$\text{rk}_{D+A}^3(g') < \text{rk}_{D+A}^3(g) = \text{rk}_D^3(g)$$

- a contradiction.

Clause (d) follows from 3.4(9) replacing D by $D + A$. □_{3.14}

Definition 3.15 1. Let E be as in 3.0, κ an aleph and $\kappa = \text{cf}(\kappa) > \aleph_0$, and D^\oplus a filter on κ . We say that E is [weakly] D^\oplus -normal if there is a function ι witnessing it which means:

ι is a function from $A^* = \text{Dom}(E)$ to κ such that

(a) for every $D \in E$

$$i(D) =: \{A \subseteq \kappa : \iota^{-1}(A) \in D\}$$

is a filter on κ extending D^\oplus ,

(b) every $D \in E$ is [weakly] ι -normal, which means: if $f : A^* \rightarrow \kappa$ is ι -pressing down (i.e. $f(a) < 1 + \iota(a)$ for $a \in A^*$) then on some $A \in D^+$, the function f is constant [bounded].

2. We say that E is [weakly] κ -normal if this holds for some κ as above, filter D^\oplus over κ and function ι .
3. We say that E is [weakly] (κ, S) -normal, where $S \subseteq \kappa$, if this holds for some filter D^\oplus over κ such that $S \in D^\oplus$.
4. We say that D is κ -normal (with ι witnessing it) if $E := \{D + A : A \in D^+\}$ is, with ι witnessing it.

If $\text{Dom}(E) = \kappa$, ι the identity we omit it.

Omitting D from “ D -normal” means omitting clause (a).

Remark 3.16 1. Note: for a filter D on κ the normality is also defined as the closure under diagonal intersection; this is equivalent. But it is not enough that the diagonal intersection of clubs is a club, we need that the diagonal intersection of sets including clubs includes a club.

2. The club filter D_κ^{cl} on a regular $\kappa > \aleph_0$ is not necessarily normal, but there is a minimal normal filter on it, D_κ^\otimes , but possibly $\emptyset \in D_\kappa^\otimes$, see below.

Definition 3.17 1. For an ordinal δ

$$D_\delta^{\text{bd}} = \{A \subseteq \delta : A \text{ bounded in } \delta\}$$

$$D_\delta^{\text{cl}} = \{A \subseteq \delta : A \text{ contains a club of } \delta\}.$$

2. For $\mathcal{P} \subseteq \mathcal{P}(\delta)$ we define $D_{\delta, \zeta}^{\text{nor}}[\mathcal{P}]$ by induction on ζ as follows:

$\zeta = 0$ $D_{\delta, \zeta}^{\text{nor}}[\mathcal{P}]$ is the filter of subsets of δ generated by $\mathcal{P} \cup D_\delta^{\text{cl}}$.

$\zeta > 0$ limit $D_{\delta, \zeta}^{\text{nor}}[\mathcal{P}] = \bigcup_{\xi < \zeta} D_{\delta, \xi}^{\text{nor}}[\mathcal{P}]$.

$\zeta = \xi + 1$ $D_{\delta, \zeta}^{\text{nor}}[\mathcal{P}] = \{\delta \setminus B : \text{there is a function with domain } B, \text{ regressive (i.e. } f(\alpha) < 1 + \alpha), \text{ such that for every } \beta < \delta \text{ the set } \{\alpha \in \delta \setminus B : f(\alpha) = \beta\} = \emptyset \text{ mod } D_{\delta, \xi}^{\text{nor}}[\mathcal{P}]\}$

$$D_\delta^{\text{nor}}[\mathcal{P}] = \bigcup_{\zeta} D_{\delta, \zeta}^{\text{nor}}[\mathcal{P}]$$

Above we replace nor by wnr if for $\zeta = \xi + 1$ we replace $f(\alpha) = \beta$ by $f(\alpha) \leq \beta$. If $\mathcal{P} = \emptyset$ we may omit it.

3. We call $A \subseteq \kappa$ stationary if $A \neq \emptyset \text{ mod } D_\kappa^{\text{nor}}$.

Claim 3.18 1. $D_{\delta, \zeta}^{\text{nor}}[\mathcal{P}]$ increases with ζ and is constant for $\zeta \geq \theta(\mathcal{P}(\delta))$ (so $D_\delta^{\text{nor}}[\mathcal{P}] = D_{\delta, \theta(\mathcal{P}(\delta))}^{\text{nor}}[\mathcal{P}]$), moreover it is constant for $\zeta \geq \zeta_\delta^{\text{nor}}$ for some $\zeta_\delta^{\text{nor}} < \theta(\mathcal{P}(\delta))$. Note that we get $D_\delta^{\text{cl}} \subseteq D_{\delta, 1}^{\text{nor}}[\mathcal{P}]$ even if we redefine $D_{\delta, 0}^{\text{nor}}[\mathcal{P}]$ as the filter generated by $\mathcal{P} \cup D_\delta^{\text{bd}}$. Clearly $D_\delta^{\text{nor}}[\mathcal{P}]$ is the minimal normal filter on δ which includes \mathcal{P} .

2. $D_{\delta, \zeta}^{\text{wnr}}[\mathcal{P}]$ increases with ζ and it is constant for $\zeta \geq \theta(\mathcal{P}(\delta))$ (consequently $D_{\delta}^{\text{wnr}}[\mathcal{P}] = D_{\delta, \theta(\mathcal{P}(\delta))}^{\text{wnr}}[\mathcal{P}]$ and $D_{\delta, \zeta}^{\text{wnr}}[\mathcal{P}] \subseteq D_{\delta, \zeta}^{\text{nor}}[\mathcal{P}]$), moreover for $\zeta \geq \zeta_{\delta}^{\text{wnr}}$ for some $\zeta_{\delta}^{\text{wnr}} < \theta(\mathcal{P}(\delta))$. Also we get $D_{\delta}^{\text{cl}} \subseteq D_{\delta, 1}^{\text{wnr}}$ even if we redefine $D_{\delta, 0}^{\text{wnr}}[\mathcal{P}]$ as the filter generated by $\mathcal{P} \cup D_{\delta}^{\text{bd}}$. Clearly $D_{\delta}^{\text{wnr}}[\mathcal{P}]$ is the minimal weakly-normal filter on δ which includes \mathcal{P}
3. In 1), 2) if $\text{cf}(\delta) = \delta_1$, the filters D_{δ}^{wnr} , $D_{\delta_1}^{\text{wnr}}$ (and also $D_{\delta, \zeta}^{\text{wnr}}$, $D_{\delta_1, \zeta}^{\text{wnr}}$) are essentially the same. I.e. let $h: \delta_1 \rightarrow \delta$ (strictly) increases continuously with unbounded range, then

$$\text{if } A \subseteq \delta, A_1 \subseteq \delta_1, A \cap \text{Rang}(h) = h''(A_1)$$

$$\text{then } A \in D_{\delta, \zeta}^{\text{wnr}}[\mathcal{P}] \iff A_1 \in D_{\delta_1, \zeta}^{\text{wnr}}[\mathcal{P}].$$
4. If δ is not a regular uncountable cardinal then $D_{\delta}^{\text{nor}} = \mathcal{P}(\delta)$.
5. $[AC_{\delta, \mathcal{P}(\delta)} + DC]$ If δ regular uncountable, then in 1), 2) we have $\emptyset \notin D_{\delta}^{\text{nor}}$. Moreover for every regular $\sigma < \delta$,

$$S = \{\alpha < \delta: \text{cf}(\alpha) = \sigma\} \neq \emptyset \pmod{D_{\delta}^{\text{nor}}}.$$

Proof. 1) - 4) Check.

5) We use the variant of the definition starting with D_{δ}^{bd} . Assume toward contradiction that $\delta \setminus S \in D_{\delta}^{\text{nor}}$. We choose by induction on $n < \omega$, an equivalence relation E_n on S such that:

- (i) E_0 is $\{(\alpha, \beta) : \alpha, \beta \in S\}$,
- (ii) E_{n+1} refine E_n ,
- (iii) the function f_n is regressive where

$$f_n(\alpha) = \text{otp}\{\beta : \beta E_n \alpha \text{ but } \beta < \alpha \ \& \ \neg \beta E_{n+1} \alpha \text{ and } \beta = \min(\beta/E_{n+1})\}$$

(so it is definable from E_n, E_{n+1}),

- (iv) for each $\alpha < \delta$ and $n < \omega$

$$\zeta_{\alpha}^{n+1} =: \min\{\zeta : \alpha/E_{n+1} = \emptyset \pmod{D_{\delta, \zeta}^{\text{nor}}}\} < \zeta_{\alpha}^n =: \min\{\zeta : \alpha/E_n = \emptyset \pmod{D_{\delta, \zeta}^{\text{nor}}}\}$$

or both are zero.

We can carry the induction by DC . For $n = 0$ use clause (i) to define E_0 . For the choice of E_{n+1} , for each $\alpha < \delta$, $f_n \upharpoonright (\alpha/E_n)$ as required exists by the inductive definition of $D_{\delta, \zeta}^{\text{nor}}$ (does not matter if we let $D_{\delta, 0}^{\text{nor}}$ be D_{δ}^{cl} or D_{δ}^{bd}), but we have to choose

$$\langle f_n \upharpoonright (\alpha/E_n) : \alpha < \delta, \alpha = \min(\alpha/E_n) \rangle;$$

so we have to make $\leq |\delta|$ choices, each among the family of regressive function on δ . But as we have a pairing function on δ , this is equivalent to a choice of a subset of δ , so $AC_{\delta, \mathcal{P}(\delta)}$ which we assume is enough. Let $C = \{\alpha < \delta: \text{if } \beta < \alpha, m < \omega, \psi_m^{\beta} = 0 \text{ then } \sup(\beta/E_m) < \alpha\}$. Clearly it is closed. Let $\gamma < \delta$.

Now by DC we can choose by induction on $n < \omega$, an ordinal $\alpha_n < \delta$ such that $\alpha_0 \geq \gamma$ and

(*) if $\beta \leq \alpha_n$, $m < \omega$ and $\zeta_m^\beta = 0$ then $\sup(\beta/E_m) < \alpha_{n+1}$ (note β/E_m is bounded in δ).

(Not hard to show that α_{n+1} exists.) Now letting $\bigcup_{n < \omega} \alpha_n < \delta$, belongs to $C \setminus \gamma$; as $\gamma < \delta$ was arbitrary, C is a closed unbounded subset of δ . Let $\alpha(*)$ be the σ -th member of C we get easy contradiction as $\langle \zeta_n^{\alpha(*)} : n < \omega \rangle$ is eventually zero. $\square_{3.18}$

4 Investigating strong limit singular μ

Definition 4.1

1. $\otimes_{\alpha,R}$ means: there is a function e exemplifying it which means:
 $\otimes_{\alpha,R}[e]$ e is a function, $\text{Dom}(e) = \{\delta : \delta < \alpha \text{ a limit ordinal}\}$ and for every limit $\delta < \alpha$, $e(\delta)$ is an unbounded subset of δ such that it is of order type from R . (It follows that $R \supseteq \alpha \cap \text{Reg}$).
2. If $R \cap \alpha$ is the set of infinite regular cardinals $\leq \alpha$ we may omit it (then $\text{otp}(e(\delta)) = \text{cf}(\delta)$).
 If R is the set of regular cardinals $< \alpha$ union with σ (not $\{\sigma\}$!) we write σ instead of R .
3. Let $\otimes_{\alpha,R}^*$ means $\otimes_{\alpha,R \cup (\alpha \cap \text{Reg})}$.

Observation 4.2

1. If $\otimes_{\alpha,R}$, $\sigma \in R$ is not regular and $R' = R \setminus \{\sigma\}$ then $\otimes_{\alpha,R'}$.
2. If there is e satisfying $\otimes'_{\alpha,R}[e]$ below then $\otimes_{\alpha,R}$ holds (for another e)
 $\otimes'_{\alpha,R}[e]$ e is a function, $\text{Dom}(e) = \{\delta : \delta < \alpha \text{ a limit ordinal, } \delta \notin R\}$ and for every limit $\delta \in \text{Dom}(e)$, $e(\delta)$ is an unbounded subset of δ , of order type $< \delta$.
3. Also the converse of (2) is true.
4. If $\otimes_{\alpha_2, \alpha_1 \cup R_1}$ and $\otimes_{\alpha_1, \alpha_0 \cup R_0}$ then $\otimes_{\alpha_2, \alpha_0 \cup R_0 \cup R_1}$.
5. For any ordinal α letting $j_{|\alpha|}$ be 1 if $|\alpha|$ is regular and zero otherwise we have $\otimes_{\alpha, |\alpha|+j}$, hence if $\otimes_{\alpha, \beta}$ then $\otimes_{\alpha, |\beta|+j_\beta}$.
6. If $\otimes_{\alpha, \zeta}^*$ then we can define $\langle f_\beta : \beta \in [\zeta, \alpha] \rangle$, f_β a one-to-one function from β onto $|\beta|$.
7. If $\otimes_{\alpha,R}$ then we can define $\bar{f} = \langle f_\beta : \beta < \alpha \rangle$, f_β a one-to-one function from β onto $\sup(\beta \cap R)$.
8. If $\otimes_{\alpha, \sigma}$ and $\sigma < \lambda^+ \leq \alpha$ (so λ^+ is a successor) then λ^+ is regular.
9. If we have: $R \subseteq \alpha$ closed, $\bar{f} = \langle f_\beta : \beta < \alpha \rangle$, f_β a one to one function from $\sup(R \cap \beta)$ onto β then $\otimes_{\alpha,R}$.
10. If $\otimes_{\alpha,R}$ and $\beta < \gamma < \alpha$ and $[\beta, \gamma] \cap R = \emptyset$ then $|\beta| = |\gamma|$.

Proof. 1) By part 2) it suffices to have $\otimes'_{\alpha,R'}[e]$. Now there is e satisfying $\otimes_{\alpha,R}[e]$. We define a function $e' \supseteq e \upharpoonright (\alpha \setminus R)$ such that $\text{Dom}(e') = \alpha \setminus R'$: just choose for $e'(\sigma)$ a club of σ of order type $\text{cf}(\sigma)$.

2) We are given e such that $\otimes'_{\alpha,R}[e]$. We define $e'(\delta)$ for $\delta \in \alpha$ limit by induction on δ such that $\otimes_{\alpha,R}[e']$ holds.

Case 1: $\delta \in R$

We let $e'(\delta) = \delta$.

Case 2: $\delta \notin R$

We let $\gamma_\delta = \text{otp}(e(\delta))$ so $\gamma_\delta < \delta$. Let g_δ be the unique order preserving function from γ_δ onto $e(\delta)$. Necessarily γ_δ is a limit ordinal (as $e(\delta)$ has no last element), and hence $e'(\gamma_\delta)$ is a well defined unbounded subset of γ_δ of the order type from R . Let

$$e'(\delta) =: \{g_\delta(\beta) : \beta \in e'(\gamma_\delta)\}.$$

3) Straightforward.

4) Let e'_i exemplify $\otimes'_{\alpha_{i+1}, \alpha_i \cup R_i}$ (see (3)), then $e'_0 \cup e'_1$ exemplifies $\otimes'_{\alpha_2, \alpha_0 \cup R_0 \cup R_1}$ and by part 2) we can finish.

5) Let f be a one-to-one function from α onto $|\alpha|$. Define a function e' , $\text{Dom}(e') = \{\delta : |\alpha| + j_{|\alpha|} \leq \delta < \alpha, \delta \text{ a limit ordinal}\}$ which will satisfy $\otimes'_{\alpha, |\alpha| + j_{|\alpha|}}$ (enough by part 2)).

If $\delta = |\alpha|$ choose $e'(\delta)$ as an unbounded subset of $|\alpha|$ of order type $\text{cf}(|\alpha|)$. If $\delta > |\alpha|$, let $\xi_\delta = \min\{\beta : \delta = \sup\{f(\gamma) : \gamma < \beta\}\}$, necessarily it is a limit ordinal. Now, if $\xi_\delta < |\alpha|$ we let $e(\delta) = \{f(\gamma) : \gamma < \xi_\delta\}$. We are left with the case $\xi_\delta = |\alpha|$. In this case define by induction on $i < |\alpha|$, the ordinal $\beta_i = \sup\{g(\gamma) : \gamma < i\}$. As $\xi_\delta = |\alpha|$ clearly $(\forall i < |\alpha|)(\beta_i < \delta)$, also clearly $(\forall j < i)(\beta_i \leq \beta_j)$. Hence $\{\beta_i : i < |\alpha|\}$ has order type $\leq |\alpha|$; as $\delta = \sup\{g(\gamma) : \gamma < |\alpha|\}$, clearly $\delta = \bigcup_{i < |\alpha|} \beta_i$, so $e'(\delta) \stackrel{\text{def}}{=} \{\beta_i : i < |\alpha|\}$ is as required. Hence we have finished

defining e' and the proof is completed.

6) Let $\otimes'_{\alpha, \zeta}[e]$; by 4.2(4), 4.2(5) w.l.o.g $\zeta = |\zeta|$. We define f_β by the induction on β :

If $\beta = \gamma + 1$ let $f_\beta(\gamma) = 0, f_\beta(\epsilon) = 1 + f_\gamma(\epsilon)$ for $\epsilon < \gamma$.

If β is a limit ordinal, first define g_β :

$$g_\beta(\gamma) = (\text{otp}(\gamma \cap e(\beta)), f_{\min(e(\beta) \setminus (\gamma+1))}(\gamma)).$$

So g_β is a one-to-one function from β into $\text{otp}(e(\beta)) \times \sup_{\gamma < \beta} |\gamma|$. As usual we can well order this set by $\langle \cdot \rangle_{lx}$ is the lexicographic order):

$$(i_1, i_2) \langle_{lx}^* (i_2, j_2) \iff (\max\{i_1, i_2\}, i_1, i_2) \langle_{lx} (\max\{j_1, j_2\}, j_1, j_2).$$

Let h_β be the one-to-one order preserving function from $(\text{Rang}(g_\beta), \langle_{lx}^*)$ onto some ordinal γ_β . Let $f_\beta = h_\beta \circ g_\beta$; check that γ_β is a cardinal.

7) Similar proof. (We use the fact that there is a definable function giving for any infinite ordinal α a one-to-one function from $\alpha \times \alpha$ into α : just let $\beta_\alpha \leq \alpha$ be the maximal limit ordinal such that $(\forall \gamma)(\gamma < \beta \rightarrow \gamma \times \gamma < \beta)$, so for some $n, \beta^n > n$, and as above we can define a one-to-one function from α into $\underbrace{\beta \times \dots \times \beta}_n$ and from it into β).

8) Included in the proof of (6).

9) Like the proof of part (5).

10) By (7).

□_{4.2}

Lemma 4.3 Assume $\otimes_{\mu,R}$ and $\mu \geq \theta(E \times E \times \mathcal{P}(A^*))$ and

$$\alpha^* = \sup\{\text{rk}_D^2(\mu_{A^*}, E) : D \in E\} < \infty,$$

and E, A^* are as in hypothesis 3.0 and μ_{A^*} stands for the constant function with domain A^* and value μ . Then \otimes_{α^*, R^*} , where $R \subseteq R^* \subseteq R \cup [\mu, \alpha^*)$ and there is a $\langle Y_\sigma : \sigma \in R^* \setminus R \rangle$ (note that $\sigma \in R^* \setminus R$ is just an ordinal not necessarily an aleph) such that:

(a) Y_σ is a non empty set of pairs $(D, \bar{\delta})$ such that

$$D \in E, \quad \bar{\delta} = \langle \delta_a : a \in A^* \rangle, \quad \delta_a \in R$$

and $\prod_{a \in A^*} \delta_a / D$ has the true cofinality σ (see Def 2.3(1)).

(b) The Y_σ 's are pairwise disjoint. Moreover if $(D_\ell, \bar{\delta}^\ell) \in Y_{\sigma_\ell}$ for $\ell = 1, 2$ and $D_1 = D_2$ then $\bar{\delta}^1 \neq_{D_1} \bar{\delta}^2$.

Remark. Instead of $\text{rk}_D^2(\mu_{A^*}, E) < \infty$ for every $D \in E$ it is enough to assume $\text{rk}_D^2(\mu, E) < \infty$ for some $D \in E$ with

$$\alpha^* = \sup\{r_D^2(\mu_{A^*}, D) : D \in E \text{ and } \text{rk}_D^2(\mu_{A^*}, D) < \infty\}.$$

Proof. For $\alpha < \alpha^*$ and $D \in E$ let

$$F_\alpha^D = \{f \in {}^{A^*}\mu : \text{rk}_D^2(f) = \text{rk}_D^3(f) = \alpha\} \quad \text{and} \quad A_D = \{\alpha < \alpha^* : F_\alpha^D \neq \emptyset\}.$$

So (see 3.8):

- (a) $(\alpha, D) \mapsto F_\alpha^D$ and $D \mapsto A_D$ are well defined (so there are such functions),
- (b) $\alpha^* = \bigcup_{D \in E} A_D$,
- (c) if $f_1, f_2 \in F_\alpha^D$ then $f_1/D = f_2/D$,
- (d) if $f_e \in F_{\alpha_e}^D$ for $e = 1, 2$ and $\alpha_1 < \alpha_2$ then $f_1/D < f_2/D$.

By 4.2(2)+(1) it suffices to prove $\otimes'_{\alpha^*, R^* \cup (\mu+1)}$. Let e be such that $\otimes_{\mu,R}[e]$ holds. For $\delta \in (\mu, \alpha^*)$, we try to define the truth value of $\delta \in R^*$ and $e'(\delta)$ such that $\otimes'_{\alpha^*, R^* \cup (\mu+1)}[e']$ holds. We make three tries; an easier case is when the definition gives an unbounded subset of δ of order type $< \delta$: decide $\delta \notin R^*$ and choose this set as $e'(\delta)$. If not, we assume we fail and continue, and if we fail in all three of them then we decide $\delta \in R^*$, and choose Y_δ .

First try: $e_1(\delta) \stackrel{\text{def}}{=} \{\sup(\delta \cap A_D) : D \in E \text{ and } \sup(\delta \cap A_D) < \delta\}$.

Clearly this set has cardinality $< \theta(E) \leq \mu \leq |\delta|$. So the problem is that it may be bounded in δ . In this case, by (b) above, for some $D \in E$, $\delta = \sup(\delta \cap A_D)$.

Second try: Let $E_\delta = \{D : \delta = \sup(A_D \cap \delta)\}$.

So we can assume $E_\delta \neq \emptyset$. For each $D_0 \in E_\delta$ let

$$E_\delta(D_0) = \{D \in E : D_0 \subseteq D \text{ and there is } f \in ({}^{A^*}\mu) \text{ such that } f/D \text{ is} \\ \leq_D \text{-lub of } \{g/D : g \in F_\beta^{D_0}, \text{ for some } \beta \in A_{D_0} \cap \delta\}\}.$$

For $D \in E_\delta(D_0)$ let

$$F_\delta^{(D_0, D)} = \{f \in {}^{(A^*)}\mu : f/D \text{ is } \leq_D \text{-lub of} \\ \{g/D : g \in F_\beta^{D_0} \text{ for some } \beta \in A_{D_0} \cap \delta\}\}.$$

For $f \in F_\delta^{(D_0, D)}$, let $B(f) = \{a \in A^* : f(a) \text{ a limit ordinal}\}$. Now $B(f) \in D$ by 1.3(8) because of the assumptions f is a \leq_D -lub and $A_{D_0} \cap \delta$ is unbounded in δ and let

$$H(f) = \{g \in A^* \mu : (\forall a \in B(f))(g(a) \in e(f(a))) \ \& \ (\forall a \in A^* \setminus B(f))(g(a) = 0)\}.$$

(remember e is a witness for $\otimes_{\mu, R}$). Note that

- (e) for $D_0 \in E_\delta$, $E_\delta(D_0)$ is not empty [by 3.14(1)],
- (f) if $D_0 \in E_\delta$, $D \in E_\delta(D_0)$ then $F_\delta^{(D_0, D)} \neq \emptyset$,
- (g) if $f_1, f_2 \in F_\delta^{(D_0, D)}$ then $f_1/D = f_2/D$
(as $<_D$ -lub is unique mod D) hence
- (g)' $H_D(f_1) = H_D(f_2)$ where $H_D(f) = \{g/D : g \in H(f)\}$,
- (h) if $D_0 \in E_\delta$ and $f \in F_\delta^{(D_0, D)}$ and $g \in H(f)$ then $g <_D f$,
- (i) if $f \in F_\delta^{(D_0, D)}$ then $|H(f)| \leq \prod_{i \in B_2(f)} e(f(i))$.

We now define, for $f^* \in F_\delta^{(D_0, D)}$ a function $h = h_{\delta, f^*, D_0, D}$ from $H(f^*)$ to δ :

$$h(g) = \min\{\alpha < \delta : \alpha \in A_{D_0} \text{ and there is } f \in F_\alpha^{D_0} \text{ such that } \neg(f <_D g)\}.$$

(by the way, equivalently for every $f \in F_\alpha^{D_0}$). Now

(j) $h(g)$ is well defined.

[Why? As otherwise g exemplify f^* is not a $<_D$ -lub of $\bigcup\{F_\alpha^{D_0} : \alpha \in A_{D_0} \cap \delta\}$ (g is a smaller $<_{D_0}$ -upper bound).]

Also

(k) $\text{Rang}(h_{\delta, f^*, D_0, D})$ is unbounded below δ .

[Why? For every $\alpha < \delta$ there is $\beta \in A_{D_0} \cap (\alpha, \delta)$. Choose $f \in F_\beta^{D_0}$, and define $g \in {}^{(A^*)}\text{Ord}$ by $g(a) = \min(e(f^*(a)) \setminus (f(a) + 1))$. Now $g \in H(f^*)$ and $h(g) > \beta > \alpha$.]

(l) $\text{Rang}(h_{\delta, f^*, D_0, D})$ does not depend on f^* , i.e. is the same for all $f^* \in F_\alpha^{(D_0, D)}$, [by (g)], and we denote it by $t_\delta(D_0, D)$.

(m) h is a nondecreasing function from $(H(f^*), \leq_D)$ to δ .

If $|E|$ is an aleph, choose a fixed well ordering $<_E^*$ of E and if for our δ for some pair (D_0, D) we have $\text{otp}(t_\delta(D_0, D)) < \delta$, choose $e(\delta) = t_\delta(D_0, D)$ for the first such pair; but this assumption on $|E|$ is not really necessary:

If for some $D_0 \in E_\delta$, $D \in E_\delta(D_0)$ we consider (i.e. if this set is O.K., we choose it; easily it is an unbounded subset of δ):

$\{t_\delta(D_0, D) : D_0 \in E_\delta, D \in E_\delta(D_0) \text{ and for any other such } (D'_0, D') \text{ we have } \text{otp}(t_\delta(D_0, D)) \leq \text{otp}(t_\delta(D'_0, D'))\}$.

This is an indexed family of unbounded subsets of δ , indexed by a subset of $E \times E$, all of the same order type, which we call β^* . As we know $\theta(E \times E)$ is a cardinal $\leq \mu < \delta$, by observation 4.3A below it is enough to have $\beta^* < |\delta|$.

Observation 4.3A. 1) If $\mathcal{B} = \{B_c : c \in C\}$, $B_c \subseteq \delta = \sup(B_c)$, δ a limit ordinal, $\sup \text{otp}(B_c) < |\delta|$ and $\theta(C) \leq \delta$ then we can define (uniformly) from δ , \mathcal{B} an $\underset{c \in C}{\text{unbounded subset } B}$ of δ such that $\text{otp}(B) < \delta$.

2) If in part (1) we omit the assumption $\delta = \sup(B_c)$, the conclusion still holds provided that $\neg(*)$ where

(*) for every $\alpha^* < \delta$ we have $\delta = \sup \bigcup \{B_c : \alpha^* < \sup(B_c) \leq \delta\}$.

Proof. For each $i < \delta =: \sup \underset{c \in C}{\text{otp}(B_c)}$ define

$B_i^* = \{\gamma : \text{for some } c \in C \text{ the ordinal } \gamma \text{ is the } i\text{-th member of } B_c\}$.

So there is a partial function from C onto B_i^* , hence $\text{otp}(B_i^*) < \theta(C) \leq \delta$. So if for some i , the set B_i^* is unbounded in δ then let B be B_i^* for the minimal such i . If there is no such i then let $\gamma^* = \sup\{\text{otp}(B_c) : c \in C\}$. By assumption $\gamma^* < |\delta|$, let

$B =: \{\sup(B_i^*) : i < \gamma^*\}$,

so $B \subseteq \delta$, and $|B| \leq^* |\gamma^*|$ hence $|B| \leq |\gamma|$ hence $\text{otp}(B) < |\delta| \leq \delta$ and for every $\beta < \delta$ for some $c \in C$ there is $\gamma \in B_c \setminus \beta$, so for $i = \text{otp}(B_c \cap \gamma)$ we have: $\gamma \in B_i^*$ hence $\gamma \leq \sup(B_i^*) \leq \sup(B)$, so $\sup(B) = \delta$. So B is an unbounded subset of δ of order type $< \delta$.

2) Clearly

$B = \{\sup(B_c) : c \in C \text{ and } \sup(B_c) < \delta\}$

has order type $< \theta(C) \leq \delta$, so if $\delta = \sup(B)$ we are done; if not let $\alpha^* = \sup(B) < \delta$ and $C' = \{c \in C : \delta = \sup(B_c)\}$, $\mathcal{B}' =: \{B_c : c \in C'\}$ are as in the assumption of 4.3A(1). □_{4.3A}

Continuation of the proof 4.3.

Third try: We are left with the case that every $t_\delta(D_0, D)$, when well defined, has order type δ . Continue with our $f^* \in F_\delta^{(D_0, D)}$, $h = h_{\delta, f^*, D_0, D}$. For each $g \in H(f^*)$, we know that there are $A \in D^+$ and $f \in F_{h(g)}^{D_0}$ such that $g \upharpoonright A \leq f \upharpoonright A$. Now turn the table: for $A \in D^+$ let

$H(f^*, A) = \{g \in H(f^*) : \text{for some (=all) } f \in F_{h(g)}^{D_0} \text{ we have } g \leq_A f\}$

(clearly the choice of f is immaterial : some, all are the same). Now:

$(H(f^*, A), \leq_{D+A})$ is mapped by h into $t_\delta(D_0, D) \subseteq \delta$; moreover for $g', g'' \in H(f^*, A)$ we have: $h(g') < h(g'') \Rightarrow g' <_{D+A} g''$ (otherwise the minimality of $h(g'')$ is contradicted). So $(H(f^*, A), \leq_{D+A})$ has the true cofinality $\text{otp}(t_\delta^1(D_0, D, A))$, where $t_\delta^1(D_0, D, A) =: (\text{Rang}(h \upharpoonright H(f^*, A)))$. Now by observation 4.3A if we do not succeed, for some $\beta^* < \delta$ we have

(*) if (D_0, D, A) above and $t_\delta^1(D_\delta, D, A) \not\subseteq \beta^*$ then $\text{otp}(t_\delta^1(D_0, D, A)) = \delta$.

So we assume that for some $\beta^* < \delta$ we have (*). Without loss of generality β^* is minimal. Consider such a triple (D_0, D, A) and the appropriate $f^* = f_\delta^*$. Notice that $H(f^*, A)$ is cofinal in $(H(f^*), <_{D+A})$, (we use clause (d) of 3.14). Now consider whether the quadruple

$$\bar{x} = (D_0, D, A, \langle \text{otp}(e(f_\delta^*(a))) : a \in A^* \rangle / (D + A))$$

was considered by some earlier δ^\otimes , if so, choose minimal δ^\otimes , and we shall finish by the observation 4.3B below. To stress the dependency on δ we may write $H_\delta(f_\delta^*, A)$, and define Col_α for $\alpha < \mu$ such that $\alpha \in \text{Dom}(e)$ onto $e(\alpha)$ as the unique order preserving function from $\text{otp}(e(\alpha))$ onto $e(\alpha)$. Let us define for $g \in H(f_\delta^*, A)$, the function Col_g with domain A^* , $\text{Col}_g(a)$ being $\text{otp}(e(f_\delta^*(a)) \cap g(a))$ if $f_\delta^*(a)$ is a limit ordinal, zero otherwise. Of course Col_g depends on the choice of f_δ^* but Col_g/D does not, and let $H_\delta(f_\delta^*, A) = \{\text{Col}_g : g \in H(f_\delta^*, A)\}$, so $H_\delta(A) =: (H_\delta(f_\delta^*, A) / (D + A), <_{D+A})$ has order type δ and $H_\delta(A)$ is cofinal in $\prod_{a \in A^*} \text{otpf}_\delta^*(a) / (D + A)$, and H and $H_\delta(A)$ were definable from δ , (D_0, D, A) in a fixed way.

Similarly for $H_{\delta^\otimes}(A)$. So observation 4.3B below give us a definition of an unbounded subset $Z_0(D_0, D, A, \delta, \delta^\otimes)$ of $H_\delta(A)$ of order type δ^\otimes , hence also of $\text{otp}(H_\delta(A), <_D) = \delta$ which we call $Z(D_0, D, A, \delta, \delta^\otimes)$. So

$$\{Z(D_0, D, A, \delta, \delta^\otimes) : (D_0, D, A) = \bar{x} \upharpoonright 3 \text{ for some } \bar{x} \in \mathcal{C}\}$$

is a family of unbounded subsets of δ of order type δ^\otimes so by observation 4.3A we are done.

Fourth try: All previous ones failed, in particular there is no δ^\otimes as above. We put δ in R^* , and let Y_δ be the set of quadruple $\langle D_0, D, A, \langle \text{otp}(e(f_\delta^*(a))) : a \in A^* \rangle / D \rangle$ as above (the last one is uniquely determined by the earlier ones).

Observation 4.3 B. If for $l = 1, 2$ we have $Y_l \subseteq {}^{(A^*)}\mu$ is cofinal in $\prod_{a \in A^*} \alpha_a^l / D$ and $[g_1, g_2 \in Y_\ell \Rightarrow (g_1 =_D g_2) \vee (g_1 <_D g_2) \vee (g_2 <_D g_1)]$ and $(Y_\ell / D, <_D)$ is well ordered of order type δ_ℓ and $\delta_1 < \delta_2$ and $\{a \in A^* : \alpha_a^1 = \alpha_a^2\} \in D$ (and $\alpha_a^\ell > 0$ for simplicity) then from the parameters $D, Y_\ell / D, \langle \alpha_a^\ell : a \in A^* \rangle / D$ for $\ell = 1, 2$ we can (uniformly) define one of the following:

- (a) $Y \subseteq Y_1 / D$ cofinal in $\prod_{a \in A^*} \alpha_a^1 / D$ and a function h from Y into $\delta_2, <_D$ -increasing such that $\delta_2 = \sup \text{Rang}(h \upharpoonright Y)$.
- (b) cofinal $X \subseteq \delta_2$ of order type $< \theta(\mathcal{P}(A^*))$.

Proof. For $g_1 \in Y_1$ let

$$K(g_1) =: \{g_2 \in Y_2 : g_1 \leq_D g_2 \text{ and there is no } g'_2 \in Y_2 \text{ such that } g'_2 <_D g_2 \text{ and } g_1 \leq_D g'_2\},$$

$$K_D(g_1) =: \{g_2 / D : g_2 \in K(g_1)\},$$

$$K'_D(g_1/D) = \bigcup \{K_D(g'_1) : g'_1 \in Y_1 \text{ and } g'_1 =_D g_1\}.$$

$\mathbf{k}(g_1/D)$ is (if exists) the \leq_D -minimal $g_2/D \in Y_2$ such that

$$g'_2/D \in K'_D(g_1/D) \Rightarrow g'_2/D \leq_D g_2/D.$$

Now $K(g_1)$ is a subset of Y_2 , $K_D(g_1)$ is a subset of Y_2/D and $g'_1 =_D g''_1 \Rightarrow K_D(g'_1) = K_D(g''_1)$ hence $K_D(g_1) = K'_D(g_1/D)$.

Case 1: for some $g_1/D \in Y_1/D$, the set $K'_D(g_1/D)$ is unbounded in Y_2/D .

We can choose the $<_D$ -minimal such g_1/D , so $K'_D(g_1/D)$ is a well defined cofinal subset of Y_2/D (remember Y_2/D is $<_D$ -well ordered) and easily $\text{otp}(K'_D(g_1/D), <_D) < \theta(\{A/D : A \in D^+\}) \leq \theta(\mathcal{P}(A^*))$.

Case 2: not case 1.

So \mathbf{k} is a well defined function from Y_1/D into Y_2/D , and easily $g'_1/D \leq_D g''_1/D \Rightarrow \mathbf{k}(g'_1/D) \leq_D \mathbf{k}(g''_1/D)$.

Let

$$Y =: \{g_1/D : g_1/D \in Y_1/D \text{ and } [g'_1/D \in Y_1/D \ \& \ g'_1/D <_D g_1/D \Rightarrow \mathbf{k}(g'_1/D) <_D \mathbf{k}(g_1/D)]\},$$

and let $h(g_1/D) = \text{otp}(\{g_2/D \in Y_2 : g_2/D <_D \mathbf{k}(g_1/D)\})$, so $h : Y \rightarrow \delta_2$. Now Y, h are as required. $\square_{4.3B}$

$\square_{4.3}$

Claim 4.4 $[DC + \oplus_\kappa[E]]$ Assume $\mu > A^* = \text{cf}(\mu) > \aleph_0$, $\mu > \theta(E) + \kappa^+$, $\otimes_{\mu,R}$ and $\lambda = |\text{rk}_D^2(\mu, E)|$. Then $\lambda < \aleph_\gamma$, for some $\gamma < \theta(E \times E \times \mathcal{P}(A^*) \times |R|^{A^*})$.

Proof. By 4.3 and 4.2(10). $\square_{4.4}$

Observation 4.5 1. $\theta(\bigcup_{x \in X} A_x) \leq (\theta(X) + \sup_{x \in X} \theta(A_x))^+$.

2. If λ is regular, $\lambda \geq \theta(X)$, $\lambda \geq \theta(A_x)$ (for $x \in X$) then $\theta(\bigcup_{x \in X} A_x) \leq \lambda$. So e.g. $\theta(A \times A) = \theta(A)$ if $\theta(A)$ is regular and $\theta(A \times B) \leq (\theta(A) + \theta(B))^+$, and $\theta(A \times B) = \max\{\theta(A), \theta(B)\}$ (or all three are finite) when the later is regular.

Proof. 1) As $A \leq^* B \Rightarrow \theta(A) \leq \theta(B)$ and $\bigcup_{x \in X} A_x \leq^* \bigcup_{x \in X} (\{x\} \times A_x)$ clearly without loss of generality $\langle A_x : x \in X \rangle$ is pairwise disjoint. Let $A = \bigcup_{x \in X} A_x$ and

$f : A \xrightarrow{\text{onto}} \alpha$, so $\gamma_x \stackrel{\text{def}}{=} \text{otp}(\text{Rang}(f \upharpoonright A_x)) < \theta(A_x) \leq \sup_x \theta(A_x)$ and call the latter λ . So $\alpha = \bigcup \{B_\beta : \beta < \lambda\}$ where

$$B_\beta =: \{\alpha : \text{for some } x \in X \text{ we have } \alpha \in \text{Rang}(f \upharpoonright A_x) \text{ and } \beta = \text{otp}(\alpha \cap \text{Rang}(f \upharpoonright A_x))\}.$$

Let

$$A^\beta =: \{a \in \bigcup_{x \in X} A_x : \text{for the } x \in X \text{ such that } a \in A_x \text{ we have: } \beta \text{ is the order type of } \{f(b) : b \in A_x \text{ and } f(b) < f(a)\}\}.$$

Let E be the relation on A defined by $a E b \iff \bigvee_{x \in X} \{a, b\} \subseteq A_x$ so E is an equivalence relation, and the A_x are the equivalence classes.

Now $f \upharpoonright CA^\beta \cap A_x$ is constant, so $f \upharpoonright A^\beta$ respects E , so f induces a function from A^β/E onto B_β . So

$$|B_\beta| < \theta(A^\beta/E) = \theta(\{A_x : A^\beta \cap A_x \neq \emptyset\}) \leq \theta(X).$$

Hence $\text{otp}(B_\beta) < \theta(X)$. So $|\alpha| \leq \sum_{\beta < \lambda} |B_\beta| \leq \lambda \times \theta(X)$ and

$$\theta\left(\bigcup_{x \in A} A_x\right) \leq (\lambda + \theta(X))^+ = [\sup_{x \in X} \theta(A_x) + \theta(X)]^+.$$

2) We repeat the proof above. Clearly $\gamma_x < \theta(A_x) \leq \lambda$, so $x \mapsto \gamma_x$ is a function from X to λ , so as λ is regular, necessarily $\gamma^* = (\sup_{x \in X} \gamma_x) < \lambda$. So B_β is defined for $\beta < \gamma^*$ only and again $|B_\beta| < \theta(X) \leq \lambda$. Hence $\text{otp}(B_\beta) < \lambda$ and hence by “ $\gamma^* < \lambda$, λ regular” we have $\beta^* = \sup_{\beta < \gamma^*} \text{otp}(B_\beta) < \lambda$ and we finish as above.

The “e.g.” follows (with $X = A$, $A_x = A \times \{x\}$, $\lambda = \theta(A)$). $\square_{4.5}$

Theorem 4.6 [DC] *Assume*

- (a) $\mu > \text{cf}(\mu) = \kappa > \aleph_0$,
- (b) $|\bigcup_{\alpha < \mu} \mathcal{P}(\alpha)| = \mu$,
- (c) $\oplus_\kappa(E)$. (I.e. E is a nice family of filters on κ .)

Then

- (α) 2^μ is an aleph,
- (β) μ^+ is regular, and $\otimes_{2^\mu, \{\delta < 2^\mu : \text{cf}(\delta) > \mu\}}$; if $\mu = \aleph_\gamma$ then $\otimes_{2^\mu, R}$, where $R \cap \mu = \text{Reg} \cap \mu$ and $|R \setminus \mu| \leq \theta(\gamma^\kappa \times |E|^2)$,
- (γ) $\mu < \lambda \leq 2^\mu \Rightarrow \lambda$ not measurable,

Remark 4.6A. Instead of (b), \otimes_μ suffices, if 2^μ is replaced by μ^κ and $|E|$ is well ordered.

Proof. By the proof of 3.13 (and clause (b) of the assumption) there is F exemplifying $2^\mu \in \mathbf{Ts}_{D^\kappa}(\mu)$ as if $\lambda_i < \mu = \bigvee_{j < k} \lambda_j$ then $2^\mu = \bigvee_{i < \kappa} \mathcal{P}(\lambda_i)$.

Let

$$A_D =: \{\alpha : F_\alpha^D \neq \emptyset\} \quad \text{and} \quad F_\alpha^D =: \{f \in F : \text{rk}_D^2(f) = \text{rk}_D^3(f) = \alpha\}.$$

Note that F_α^D is a singleton denoted by f_α^D , so $F = \bigcup_{D \in E} F_D$ where $F_D = \{f_\alpha^D : \alpha \in A_D\}$, so the F_D are not necessarily pairwise disjoint. By clause (b) we can prove that as $\kappa < \mu$ also $|\mathcal{P}(\kappa)| < \mu$ hence $|\mathcal{P}(\mathcal{P}(\kappa))| < \mu$ and hence $|E| < \mu$ so $|E|$ is an aleph. As $|E|$ is an aleph we can well order E (say by $<_E$) and hence can well order F :

$$f <^* g \text{ iff } \alpha(f) < \alpha(g) \text{ or } \alpha(f) = \alpha(g) \text{ and } D(f) <_E D(g)$$

where $\alpha(f)$ is the unique α such that for some D we have $f = f_{\alpha(f)}^D$ and $D(f)$ is the $<_E$ -first D which is suitable. As F is well ordered, $|F|$ is an aleph, but F was chosen by 3.13 such that $|F| = 2^\mu$, so we have proved clause (α) .

If $\lambda \in (\mu, 2^\mu]$, there is $F' \subseteq F$ of cardinality λ so as in 2.8, λ is not measurable so clause (γ) holds.

Now we shall deal with clause (β) .

By assumption (b) clearly \otimes_μ , so 4.3 applies and we get \otimes_{2^μ, R^*} for R^* as there. Now suppose that for $\langle \sigma_i : i < \kappa \rangle \in {}^\kappa(\text{Reg} \cap \mu)$, and $D \in E$, $\prod_{i < \kappa} \sigma_i / D$ has

the true cofinality σ . If for some $A \in D^+$, $\sup_{i \in A} \sigma_i < \mu$, $\prod \sigma_i / (D + A)$ is well ordered by (b), so $\sigma < \mu$. So assume $\text{tlim}_D \sigma_i = \mu$ (i.e. $\mu = \limsup_D \langle \sigma_i : i < \kappa \rangle = \liminf_D \langle \sigma_i : i < \kappa \rangle$), then $\prod \sigma_i / D$ is μ^+ -directed so $\text{cf}(\sigma) > \mu$, hence $\text{cf}(\sigma) \geq \mu^+$.

We conclude $R^* \cap \mu^+ \subseteq R$, hence, by 4.3, $\otimes_{\mu^+, R}$, so, by 4.2(8), μ^+ is regular. This gives the first phrase in clause (β) , the second is straightforward. $\square_{4.6}$

Discussion 4.6B:

1. If $|E| + |\mathcal{P}(\kappa)|$ is an aleph and the situation is as in 4.6, then we can choose $A \subseteq \theta(2^\mu)$ which codes the relevant instances of $\otimes_{\mu^+, R}$, \otimes_{λ, R^*} (or \otimes_{2^μ, R^*}) and then work in $L[A]$ and apply theorems on cardinal arithmetic (see in [Sh-g]) as in 4.4 (so we can ask on weakly inaccessible etc.), but we have to translate them back to \mathbf{V} , with A ensuring enough absoluteness.
2. If $|E| + |\mathcal{P}(\kappa)|$ is not an aleph, we can force this situation not collapsing much, see Sect. 6.

Definition 4.7 For an ordinal δ let

$$ID_\delta^1 = \{A \subseteq \delta : \otimes_{\delta, \delta \setminus A} \text{ holds}\},$$

$$ID_{\delta, \alpha}^1 = \{R \subseteq \delta : R \setminus \alpha \in ID_\delta^1\},$$

$$ID_{\delta, R}^2 = \{A \subseteq \delta : \text{there is a function } e \text{ with domain } A \text{ such that} \\ \text{the requirement in } \otimes_{\delta, R} [e] \text{ holds for } \alpha \in A\},$$

$$ID_{\delta, R, \alpha}^2 = ID_{\delta, R \cup \alpha}^2.$$

Omitting R means $\text{Reg} \cap \delta$.

Claim 4.8 1. $ID_{\delta, 0}^1 = ID_\delta^1$ and $\delta \in ID_{\delta, R}^2 \iff \otimes_{\delta, R}$.

2. $ID_{\delta, \alpha}^1, ID_\delta^2$ are ideals of subsets of δ .

3. Assume $\alpha_1 < \alpha_2 < \alpha_3 \in \text{Ord}$ and $ID_{\delta_{l+1}, R, \alpha_l}^2$ for $l = 1, 2$. Then $ID_{\alpha_3, R, \alpha_1}^2$.

4. $\delta \in ID_\delta^2$ if $AC_{\delta, <|\delta|}$.

5. $[AC_{|\alpha|}] ID_{\delta, R, \alpha}^2$ is $|\alpha|^+$ -complete (see 4.3A, we need to choose the witnesses e_1).

6. $A \in ID_\delta^1 \iff A \in ID_{\delta, \delta \setminus A}^2$

7. $[AC_{|\alpha|}] ID_{\delta, \alpha}^1$ is $|\alpha|^+$ -complete.

We think that those ideals are very interesting.

5 The successor of a singular of uncountable cofinality

Claim 5.1 Assume $\text{cf}(\mu) = \kappa > \aleph_0$, $\mu = \bigcup_{i < \kappa} \mu_i$ where μ_i increasing continuous, μ an aleph (μ_i may be merely an ordinal). If $f^* \in {}^\kappa \text{Ord}$, $f^*(i) = |\mu_i|^+$ then $\|f^*\|_{D_\kappa^{\text{bd}}} \geq \mu^+$ (hence $\text{rk}_D^2(f^*) \geq \mu^+$ when $D \in E$ extend D_κ^{bd}).

Proof. Let $\alpha < \mu^+$ and we shall prove

(*) there is $\langle f_\beta : \beta \leq \alpha \rangle$, $f_\beta \in \prod_{i < \kappa} f^*(i)$, such that $\beta < \gamma < \alpha \Rightarrow f_\beta <_{D_\kappa^{\text{bd}}} f_\gamma$

Let g be a one-to-one function from α into μ . For every $\beta \leq \alpha$ let

$$f_\beta(i) =: \text{otp}\{\gamma < |\mu_i|^+ : g(\gamma) < \beta\}.$$

□_{5.1}

Observation 5.2 [DC + AC $_\kappa$ + $\oplus_\kappa[E]$, E normal, κ an aleph] Assume

(a) $\langle \mu_i : i \leq \kappa \rangle$ is strictly increasing continuous sequence of alephs,

(b) $f \in \prod_{i < \kappa} (\mu_i^+ + 1)$, $D_0 \in E$ and $\text{rk}_{D_0}^2(f, E) = \text{rk}_{D_0}^3(f, E) = \mu^+$,

(c) $\text{Tw}_D(\mu_i) < \mu$ for $D \in E$ and $i < \kappa$.

Then $\{i : f(i) = \mu_i^+\} \in D$.

Proof. Otherwise without loss of generality $(\forall i)(0 < f(i) < \mu_i^+)$ hence $(\forall i)(\text{cf}[f(i)] < \mu_i)$. By AC $_\kappa$ there is $\langle g_i : i < \kappa \rangle$, g_i is a one-to-one function from $f(i)$ into μ_i . By the normality and the possibility to replace D by $D + A$ (for any $A \in D^+$), for every $f_1 \in \prod_{i < \kappa} f(i)$ and $D_1, D_0 \subseteq D_1 \in E$, for some $A \in D_1^+$, and $j < \kappa$ we have $(\forall i \in B)(g_i \circ f_1(i) < \mu_j)$, so we conclude

$$\mu^+ = \bigcup \{W_{D,j,A} : D_0 \subseteq D \in E, j < \kappa, B \in D\},$$

$\langle F_{D,j,A,\alpha} : \alpha \in W_{D,j,A} \rangle$ is $<_D$ -increasing and D -smooth, where

$$F_{D,j,A,\alpha} = \{f_1 : f_1 \in \prod_{i < \kappa} f(i), [i \in A \Rightarrow g_i \circ f_1(i) < \mu_j] \text{ and } \text{rk}_D^2(f_1, E) = \text{rk}_D^3(f_1, E)\},$$

$$W_{D,j,A} =: \{\alpha : F_{D,j,A,\alpha} \neq \emptyset\}.$$

As $\text{Tw}_D(\mu_j) \leq \mu$, clearly $|W_{D,j,A}| \leq \mu$. So

$$\mu^+ = \bigcup \{W_{D,j,A} : D_0 \subseteq D \in E, j < \kappa, A \subseteq \kappa\}$$

with each set of cardinality $< \mu$, as $\theta(E) < \mu$, $\theta(\mathcal{P}(\kappa)) < \mu$ (as in 4.3), we get a contradiction. □_{5.2}

Conclusion 5.3 Under the hypothesis and the assumptions (a)-(c) of 5.2, for the $f^* \in {}^\kappa \text{Ord}$ defined by $f^*(i) =: \mu_i^+$, for every $D \in E$ we have: $\text{rk}_D^2(f, E) = \text{rk}_D^3(f, E) = \mu^+$.

Theorem 5.4 $[DC + AC_\kappa + \oplus_\kappa(E) + \aleph_0 < \kappa = \text{cf}(\kappa), E \text{ normal or just weakly normal as witnessed by } \iota]$ Assume (a)–(c) of 5.2 and

(d) $S = \{i < \kappa : \text{cf}(\mu_i^+) = \lambda\}$ is stationary (see 3.17 (2)), moreover $\iota^{-1}(S) \in D_0 \in E$ for some D_0 .

Then $\text{cf}(\mu_\kappa^+) = \lambda$

Proof. By 5.1 + 3.14 we know $\text{rk}_D^3(\langle \mu_i^+ : i < \kappa \rangle, E) \geq \mu^+$ so by 3.4(6) there are $f \in \prod_{i < \kappa} (\mu_i^+ + 1)$ and $D \in E$ such that $S \in D$, $\text{rk}_D^2(f) = \text{rk}_D^3(f) = \mu^+$. Without loss of generality each μ_i is singular; apply 5.2. By AC_κ there is $\langle e_i : i < \kappa \rangle$, $e_i \subseteq \mu_i^+$ cofinal, $\text{otp}(e_i) = \text{cf}(\mu_i^+)$. Now as in the proof of 5.3 we know $\{\text{rk}_D^3(h) : D \in E, h \in \prod_{i < \kappa} e_i\}$ is cofinal in μ^+ , hence

$$\{\text{rk}_D^3(h) : D \in E, h(i) \text{ the } \gamma\text{-th member of } e_i \text{ for } i \in S, 0 \text{ otherwise}\}$$

is a cofinal subset of μ^+ .

□_{5.4}

Conclusion 5.5 $[as 5.4+(a), (b), (c) \text{ of } 5.2]$ If $\langle \mu_i : i \leq \kappa \rangle$ is a strictly increasing continuous sequence of alephs then for at most one $\lambda > \kappa$ the set $\{i < \kappa : \text{cf}(\mu_i^+) = \lambda\}$ is stationary

Remark. We can prove that for all $D \in E$, $\prod_{i < \kappa} \text{cf}(\mu_i^+)/D$ has the same cofinality.

Question 5.6. Under the hypothesis and assumptions of 5.4 can we in addition have a stationary $S \subseteq \kappa$ such that $\langle \text{cf}(\mu_i^+) : i \in S \rangle$ is with no repetition?

6 Nice E exists

Observation 6.1 For an aleph λ and a set A^* , $\lambda^{|A^*|}$ is equal to

$$\sum_{\beta < \theta(A^*)} \lambda^{|\beta|} \times |\beta|^{|A^*|}$$

provided that $\aleph_0 \leq |A^*|$ (otherwise replace $|\beta|^{|A^*|}$ by $|\{f \in A^* : \text{Rang}(f) = \beta\}|$).

Convention 6.2 κ, σ are alephs, σ an uncountable cardinal such that $DC_{<\sigma}$, f, g, h will be functions from κ to Ord .

Definition 6.3 Assume $\text{cf}(\kappa) \geq \theta$.

1. $P_{\kappa, \alpha}^\sigma = \{f : f \text{ a function from some ordinal } \gamma < \sigma \text{ into } \kappa \cup {}^\kappa\alpha\}$ partially ordered by \subseteq is regarded as a forcing notion and $G_{P_{\kappa, \alpha}^\sigma}$ or simply G denote the generic.
2. $\dot{h} = \bigcup G_{P_{\kappa, \alpha}^\sigma}$ (is forced to be a function from σ onto $\kappa \cup {}^\kappa\alpha$).

3. $\bar{Q}[GP_{\kappa,\alpha}^\sigma] = \langle \alpha_i : i < \sigma \rangle$ lists in increasing order

$$\{\sup(\kappa \cap \text{Rang}(\underline{h} \upharpoonright \gamma)) : \gamma < \sigma\}.$$

4. $\underline{Y} = \{\langle \gamma, \underline{h}(\gamma) \rangle : \gamma < \sigma, \underline{h}(\gamma) \in \kappa\} \cup \{\langle \gamma, i, \beta \rangle : \gamma < \sigma, i < \kappa \text{ and } \underline{h}(\gamma) \in {}^\kappa\alpha, (\underline{h}(\gamma))(i) = \beta\}.$

5. $\underline{D}_{\kappa,\theta}^\sigma$ is the club filter on the ordinal κ in $K[\underline{Y}]$ (K - the core model, see Dodd and Jensen [DJ]).

6. We can replace $\alpha, {}^\kappa\alpha$ by $f \in {}^\kappa\text{Ord}$, $\prod_{i < \kappa} (f(i) + 1)$ respectively.

Remark. On forcing for models of ZF only see e.g. [Bu].

Observation 6.4 1. $D_{\kappa,\alpha}^\sigma = \underline{D}_{\kappa,\alpha}^\sigma \cap \mathcal{P}(\kappa)^\mathbf{V}$ does not depend on the generic subset G of $P_{\kappa,\alpha}^\sigma$.

2. If κ is regular in $\mathbf{V}^{P_{\kappa,\alpha}^\sigma}$ (e.g. κ successor in \mathbf{V}) then $D_{\kappa,\alpha}^\sigma$ is normal, minimal among the normal filters on κ .

Claim 6.5 Assume that

$\boxtimes_{\sigma,\kappa,\alpha}$ for some/every $G \subseteq P_{\kappa,\alpha}^\sigma$ generic over \mathbf{V} , in $K[\underline{Y}]$ there are arbitrarily large Ramsey cardinals³.

Then $\text{rk}_{D_{\kappa,\lambda}^\sigma}^2(\alpha) < \infty$ for $E = E_0$ the family of filters on κ , or $E = E_1$ the family of normal filters on κ in the case of 6.4(2), or even

$$E = E_2 = \{D : \Vdash_{P_{\kappa,\alpha}^\sigma} \text{“there is a normal filter } D' \text{ on } \kappa \text{ such that } D' \cap \mathcal{P}(\kappa)^\mathbf{V} = D \text{”}\}.$$

Proof. As $P_{\kappa,\alpha}^\sigma$ is a homogeneous forcing, “some G ” and “every G ” are equivalent. The proof is as in [Sh-g], Ch.V 1.6, 2.9. $\square_{6.5}$

Remark. In [Sh-g] Ch.V we almost get away with $K(Y)$, $Y \subseteq (2^{\aleph_1})^+$. We shall return to this later.

Claim 6.6 Assume that $\boxtimes_{\sigma,\kappa,\alpha}$ fails.

1. If in \mathbf{V} , $\lambda = \mu^+$, μ is singular $> \theta^{(\sigma > (\kappa\alpha))}$ then λ is regular but not measurable.

2. Bounds to cardinal arithmetic : if $\mu \geq \kappa$ then $\lambda^\mu \leq 2^\mu \times \lambda^+$

Proof. Let $G \subseteq P_{\kappa,\alpha}^\theta$ be generic over \mathbf{V} . First we shall prove that

(*)₁ λ is still a cardinal in $\mathbf{V}[G]$.

³ Or considerably less

Assume not. So $\mathbf{V}[G] \models |\lambda| = |\mu|$, and hence for some P -name \underline{H} and $p \in P$,

$$p \Vdash_P \text{“}\underline{H} \text{ is a one-to-one function from } \mu \text{ onto } \lambda\text{”}.$$

For $\alpha < \mu$ let

$$A_\alpha = \{\beta < \lambda : p \not\Vdash_P \underline{H}(\alpha) \neq \beta\}.$$

So $\langle A_\alpha : \alpha < \mu \rangle \in \mathbf{V}$ and easily $\text{otp}(A_\alpha) < \theta(P_{\kappa, \alpha}^\sigma)$ (map $q \in P$ to β if $p \leq q$, $q \Vdash \underline{H}(\alpha) = \beta$, and map $q \in P$ to $\min(A_\alpha)$ otherwise). But $\theta(P_{\kappa, \alpha}^\sigma) \leq \mu$, so (as the sets are well ordered) this implies

$$\mathbf{V} \models \lambda = \left| \bigcup_{\alpha < \mu} A_\alpha \right| \leq \sum_{\alpha < \mu} \text{otp}(A_\alpha) \leq \sum_{\alpha < \mu} \theta(P_{\kappa, \alpha}^\sigma) \leq \mu \times \mu \leq \mu;$$

a contradiction.

Really we have proved

$(*)_1^+$ every aleph $\geq \theta(\sigma(\kappa\alpha))$ in \mathbf{V} is a cardinal in $\mathbf{V}[G]$.

Next we prove

$(*)_2$ λ is regular in \mathbf{V} .

Assume not. So there are $A_2 \subseteq \lambda = \sup(A_2)$, $\text{otp}(A_2) = \text{cf}(\lambda)$, $A_0 \subseteq \mu = \sup(A_0)$, $\text{otp}(A_0) = \text{cf}(\mu)$. Let $\mu^* = (\mu^+)^{K[Y[G]']}$ and let in \mathbf{V} we have $A_1 \subseteq \mu^* = \sup(A_1)$, $\text{otp}(A_1) = \text{cf}(\mu^*)$. Let now $\bar{A} = \langle A_0, A_1, A_2 \rangle$. In $\mathbf{V}[G]$, μ, λ are still cardinals. Hence in $K[\underline{Y}[G]]$ too μ, λ are cardinals and even in $K[\underline{Y}[G], \bar{A}]$ we have λ is a cardinal. Also $\mu^* \in (\mu, \lambda)$, hence $\text{cf}(\mu^*) < \mu$ hence $K[\underline{Y}[G], \bar{A}] \models \text{cf}(\mu^*) < \mu$. Now $K[\underline{Y}[G]]$, $K[\underline{Y}[G], \bar{A}]$ are models of ZFC , and in the latter $\text{cf}(\mu^*) \leq \text{otp}(A_2) < \mu$.

As $K[\underline{Y}[G]]$ does not have unboundedly many Ramsey cardinals and $\mu > \aleph_1^{\mathbf{V}} = \aleph_1^{K[\underline{Y}[G]']}$, by Dodd and Jensen [DJ] there is $B \in K[\underline{Y}[G]]$ such that $K[\underline{Y}[G]] \models \text{“}|B| < (\mu^+)^{K[\underline{Y}[G], \bar{A}]}$ ” and $A_1 \subseteq B$, and so we get contradiction to $K[\underline{Y}[G]] \models \text{“}\mu^* \text{ a successor cardinal } > \mu\text{”}$.

Next:

$(*)_3$ if μ is singular, λ is not measurable;

[as by the proof of $(*)_2$, $K[\underline{Y}[G]] \models \text{“}\lambda = \mu^+\text{”}$ so there is e as in Sect. 4]. $\square_{6.6}$

Claim 6.7 $[AC_{\mathcal{P}(\mathcal{P}(\kappa))} + DC]$ Assume E is a family of filters on κ , $D_* = \min(E)$ (that is $D_* \in E$, and $[D \in E \Rightarrow D_* \subseteq D]$) and for some $f^* \in {}^\kappa \text{Ord}$, $\text{rk}_{D_*}^2(f^*) = \infty$. Then $\text{rk}_{D_*}^2(\alpha) = \infty$ for some $\alpha, \alpha < \theta(\mathcal{P}(\mathcal{P}(\kappa)))$ (and essentially $\alpha < \theta(\mathcal{P}(\kappa))$) (i.e. α stands for the function from κ to Ord which is constantly α).

Proof. We first prove that there is such $\alpha < \theta(\mathcal{P}(\mathcal{P}(\kappa)))$. If $D \in E$, $\text{rk}_D^2(f) = \infty$ then for every ordinal α there are $A = A_{f, \alpha} \in D^+$ and $g = g_{f, \alpha} <_{D+A_{f, \alpha}} f$ such that $\text{rk}_{D+A}^3(g) > \alpha$. Without loss of generality $g_{f, \alpha} \leq f$; so we have only a set of possible $(A_{f, \alpha}, g_{f, \alpha})$ (for given f).

By the “ F ” of ZF there are $A_f, g_f <_{D+A_f} f, g_f \leq f$ such that $\text{rk}_{D+A_f}^3(g_f) = \infty$, so $\text{rk}_{D_1}^2(g_f) = \infty$ for every $D_1, D + A_f \subseteq D_1 \in E$. Note:

$$\kappa + \kappa = \kappa, \quad |\mathcal{P}(\kappa)|^2 = |\mathcal{P}(\kappa)|, \quad |\mathcal{P}(\mathcal{P}(\kappa))|^2 = |\mathcal{P}(\mathcal{P}(\kappa))|$$

(really $|A| + 1 = |A| \Rightarrow |\mathcal{P}(A)|^2 = |\mathcal{P}(A)|$). By $AC_{\mathcal{P}(\mathcal{P}(\kappa))}$:

(*) if $F \subseteq {}^\kappa\text{Ord}$ has cardinality $< |\mathcal{P}(\mathcal{P}(\kappa))|$ then we can find

$$\{ \langle A_{f,D}, g_{f,D} \rangle : f \in F, D \in E, \text{rk}_D^2(f) = \infty \}$$

such that $\text{rk}_{D+A_f,D}^3(g_{f,D}) = \infty, g_{f,D} <_{D+A_f,D} f, g_{f,D} \leq f$.

By DC we can find $\langle F_n : n < \omega \rangle, F_n \subseteq {}^\kappa\text{Ord}$, such that in (*) for $F = F_n$ we have $F_{n+1} = F_n \cup \{g_{f,D} : f \in F_n, D \in E\}$, and $F_0 = \{f^*\}$. Let $F = \bigcup F_n$, and let

$\bar{A} = \langle A_i : i < \kappa \rangle$ be defined by $A_i = \{f(i) : f \in F\}$. Clearly $|F_n| \stackrel{n < \omega}{\leq^*} |\mathcal{P}(\mathcal{P}(\kappa))|$ and hence $|A_i| \leq^* |\mathcal{P}(\mathcal{P}(\kappa))|$. Hence $h^*(i) =: \text{otp}(A_i) < \theta(\mathcal{P}(\mathcal{P}(\kappa)))$ and $\text{sup Rang}(h^*) < \theta(\mathcal{P}(\mathcal{P}(\kappa)))$. Now for every $f \in F$ we define $h_f \in \prod_{i < \kappa} g(i)$

by $h_f(i) = \text{otp}(A_i \cap f(i)) < g(i)$. It is now easy to check that $\text{rk}_{D^*}^2(h_{f^*}) = \infty$ as required.

Next we prove that for some $\alpha < \theta(\mathcal{P}(\kappa))$ we have $\text{rk}_{D^*}^2(\alpha) = \infty$. As $AC_{\mathcal{P}(\mathcal{P}(\kappa))}$, clearly $\mathcal{P}(\kappa)$ can be well ordered, so let $h_0 : \mathcal{P}(\kappa) \rightarrow |\mathcal{P}(\kappa)|$ be one-to-one onto $|\mathcal{P}(\kappa)|$, an aleph. By the first part there is $\alpha^* < \theta(\mathcal{P}(\mathcal{P}(\kappa)))$ with $\text{rk}_{D^*}^2(\alpha^*) = \infty$. Hence we can find $h_1^* : \mathcal{P}(\mathcal{P}(\kappa)) \xrightarrow{\text{onto}} \alpha^* + 1$.

Observation 6.7A. $[AC_\kappa]$ If $\alpha < \theta(\mathcal{P}(A^*))$, $|A^*| \times \kappa = |A^*|$ then ${}^\kappa(\alpha + 1)$ is a set of cardinality $\leq^* \mathcal{P}(A^*)$.

[Why? Let $h_1 : \mathcal{P}(A^*) \xrightarrow{\text{onto}} \alpha + 1$ and let $h_2 : A^* \times \kappa \rightarrow A^*$ be one to one. For each $f \in {}^\kappa(\alpha + 1)$ we can choose $\bar{B} = \langle B_i : i < \kappa \rangle, h_1(B_i) = f(i)$, so $B_i \subseteq \mathcal{P}(A^*)$, and $\langle B_i : i < \kappa \rangle$ encodes f and it in turn can be coded by $\{(i, x) : i < \kappa \text{ and } x \in B_i\} \subseteq \kappa \times A^*$, but it is not clearly if we have also the function $f \mapsto \bar{B}^f$. However we can define a function H , $\text{Dom}(H) = \mathcal{P}(A^*)$ and $\text{Rang}(H) \subseteq {}^\kappa(\alpha + 1)$ by

$$(H(A))(i) = h_1(\{a \in A^* : h_2(a, i) \in A\}).$$

Clearly H is a function from $\mathcal{P}(A)$ onto ${}^\kappa(\alpha + 1)$, hence $|{}^\kappa(\alpha + 1)| \leq^* |\mathcal{P}(A^*)|$.

□_{6.7A}

Of course, apply 6.7A to $A^* = \mathcal{P}(\kappa)$.

Also $|\text{Fil}(\kappa)| \leq |\mathcal{P}(\mathcal{P}(\kappa))|$ and $|\mathcal{P}(\mathcal{P}(\kappa))|^2 = |\mathcal{P}(\mathcal{P}(\kappa))|$, so by $AC_{\mathcal{P}(\mathcal{P}(\kappa))}$ we can find $Y_0 = \langle \langle A_{f,D}, g_{f,D} \rangle : f \in {}^\kappa(\alpha^* + 1), D \in E \rangle$ such that:

(*) if $\text{rk}_D^2(f) = \infty$ then $\text{rk}_{D+A_f,D}^3(g_{f,D}) = \infty, A_{f,D} \in D^+, g_{f,D} \in {}^\kappa(\alpha^* + 1), g_{f,D} <_{D+A_f,D} f$

Similarly by $AC_{\mathcal{P}(\mathcal{P}(\kappa))}$ we can find $Y_1 = \langle \langle d_{f,D} : f \in {}^\kappa(\alpha^* + 1), D \in E \rangle$ such that:

$$D \subseteq d_{f,D} \in E \text{ and } \text{rk}_D^3(f) = \text{rk}_{d_D}^2(f).$$

We now define the model \mathfrak{C} with the universe $\mathcal{P}(\mathcal{P}(\kappa))$ - just put all the information needed below.

Clearly $|\mathfrak{C}| = |\mathcal{P}(\mathcal{P}(\kappa))|$, so we have a choice function H^* on the family of definable (in \mathfrak{C}) non empty subsets of \mathfrak{C} . So for $A \subseteq \mathfrak{C}$ we have the Skolem hull. Now we define by induction on $\alpha \leq \kappa^+$ (an aleph) submodels M_α of \mathfrak{C} increasing continuously in α and of cardinality $\leq |\mathcal{P}(\kappa)|$.

For $\alpha = 0$: the Skolem hull of $\{\gamma : \gamma \leq |\mathcal{P}(\kappa)|\}$

For α limit: $\bigcup_{\beta < \alpha} M_\beta$

For $\alpha = \beta + 1$: the Skolem hull of $|M_\alpha| \cup \{x : x \in {}^\kappa M_\alpha\}$

Now ${}^\kappa(M_{\kappa^+}) \subseteq M_{\kappa^+}$ (as κ^+ is regular as $|\mathcal{P}(\kappa)|$ is an aleph). Let $H : M_{\kappa^+} \cap \text{Ord} \xrightarrow{\text{onto}} \beta^* < |\mathcal{P}(\kappa)|^+$ be order preserving. So as we are assuming that the conclusion fails, for every $f \in {}^\kappa(\alpha^* + 1) \cap M_{\kappa^+}$,

$$H \circ f^* \in {}^\kappa \beta^* \text{ satisfies } \text{rk}_{D_*}^2(H \circ f^*) < \infty.$$

Whereas $\text{rk}_{D_*}^2(f^*) = \infty$. We can conclude as in [Sh 386, 1.13]. $\square_{6.7}$

We can generalize 6.1-6.7

Claim 6.8 $[DC_{<\sigma}, \sigma = \text{cf}(\sigma) > \aleph_0]$ Assume:

(a) D_* is a σ -complete filter on A^*

(b) $A^* = A_* \times \sigma$, and $\iota : A^* \rightarrow \sigma$ is $\iota(a, \alpha) = \alpha$

(c) D^* is the following filter on A^* ; for $A \subseteq A^*$:

$$A \in D^* \iff \{i < \sigma : \{a \in A_* : (i, a) \in A\} \in D_*\} \in D_\sigma \text{ where}$$

$$D_\sigma \text{ is the minimal normal filter on } \sigma.$$

(d) $E =$ the family of ι -normal filters on A^* .

Then for every $f \in {}^{A^*} \alpha^*$, $\text{rk}_D^2(f) < \infty$, provided that $\otimes_{|A^*|, \alpha^*}^\sigma$.

Concluding Remark. We can deal with $\text{fc}(\omega > \omega)$ as in [Sh 386], [Sh 420]. Also concerning 6.7 change $|A^*|$.

Also as in [Sh 386] we can define ${}^1\text{rk}^2(f, \mathcal{E})$ and so improve 3.12 (as in [Sh 386, Sect. 2]).

References

- [ApMg] Apter, A., Magidor, M.: Instances of dependent choice and the measurability of $\aleph_{\omega+1}$. Ann. App. Logic (submitted)
- [Bu] Bull, E.: Consecutive large cardinals. Ann. Math. Logic **15**, 161-191 (1978)
- [DJ] Dodd, A., Jensen, R. B.: The core model. Ann. Math. Logic **20**, 43-75 (1981)
- [Gi] Gitik, M.: All uncountable cardinals can be singular. Israel J. Math. **35**, 61-88 (1980)
- [GH] Galvin, F., Hajnal, A.: Inequalities for cardinal powers. Ann. Math. **101**, 491-498 (1975)
- [J] Jech, T.: Set Theory. New York: Academic Press 1978
- [J1] Jech, T.: The Axiom of Choice. In: Studies in Logic and the Foundations of Mathematics, 75. Amsterdam: North-Holland 1973

- [Kf] Kafkoulis, G.: Homogeneous sequences of cardinals for OD partition relations. Doctoral Dissertation, Cal. Tech (1989)
- [Sh-g] Shelah, S.: Cardinal Arithmetic. In: Gabbai, D. M., Macintyre, A., Scott, D. (eds.) Oxford Logic Guides, vol. 29. Oxford: Oxford University Press 1994
- [Sh 176] Shelah, S.: Can you take Solovay inaccessible away? *Israel J. Math.* **48**, 1-47 (1984)
- [Sh 386] Shelah, S.: Bounding $pp(\mu)$ when $cf(\mu) > \aleph_0$ using ranks and normal ideals. *Cardinal Arithmetic, Chapter V*. In: Gabbai, D. M., Macintyre, A., Scott, D. (eds.) Oxford Logic Guides, vol. 29. Oxford: Oxford University Press 1994
- [Sh 420] Shelah, S.: Advances in Cardinal Arithmetic. In: Sauer, N. W. et al (eds.) *Finite and infinite combinatorics in sets and logic*, pp. 355-383. Amsterdam: Kluwer 1993
- [Sh 506] Shelah, S.: The pcf-theorem revisited. In: Graham, R., Nešetřil, J (eds.) *Mathematics of Paul Erdős*, vol 2. Berlin Heidelberg New York: Springer
- [Sh:F239] Shelah, S.: Universes without choice (in preparation)