




ELSEVIER

Available online at www.sciencedirect.com



Advances in Mathematics 192 (2005) 403–426

ADVANCES IN
Mathematics

<http://www.elsevier.com/locate/aim>

Comparing the uniformity invariants of null sets for different measures

Saharon Shelah^{a,1,2} and Juris Steprāns^{b,*,3}

^a *Department of Mathematics, Rutgers University, Hill Center, Piscataway, NJ, 08854-8019, USA*

^b *Department of Mathematics, York University, 4700 Keele Street, Toronto, Ont., Canada M3J 1P3*

Received 29 July 2003; accepted 23 April 2004

Communicated by R.D. Mauldin

Abstract

It is shown to be consistent with set theory that the uniformity invariant for Lebesgue measure is strictly greater than the corresponding invariant for Hausdorff r -dimensional measure where $0 < r < 1$.

© 2004 Elsevier Inc. All rights reserved.

MSC: 03E17; 28A78

Keywords: Lebesgue measure; Hausdorff measure; Null set; Proper forcing; Product forcing; Uniformity cardinal invariant

1. Introduction

The uniformity invariant for Lebesgue measure is defined to be the least cardinal of a non-measurable set of reals, or, equivalently, the least cardinal of a set of reals

*Corresponding author. Fax: +416-736-5757.

E-mail addresses: shelah@math.rutgers.edu (S. Shelah), steprans@mathstat.yorku.ca (J. Steprāns).

¹ Current address Institute of Mathematics, Hebrew University, Givat Ram, Jerusalem 91904, Israel.

² Supported by the United States-Israel Binational Science Foundation and by Grant NSF-DMS97-04477 of the National Science Foundation of the United States. This is number 809 in his personal numbering system.

³ Supported by the Natural Sciences and Engineering Research Council of Canada and was completed while enjoying the hospitality of Rutgers University. The second author expresses his thanks to D. Fremlin and A. Miller for enduring a lengthy series of lectures based on an early version of this paper, a version even more awkward than the current.

which is not Lebesgue null. This has been studied intensively for the past 30 years and much of what is known can be found in [1] and other standard sources. Among the well known results about this cardinal invariant of the continuum is that it can equally well be defined using Lebesgue measure on \mathbb{R}^n without changing the value of the cardinal. Indeed, equivalent definitions will result by using any Borel probability measure on any Polish space. However, the question of the values of uniformity invariants for other, non- σ -finite Borel measures is not so easily answered. This paper will deal with the most familiar class of such measures, the Hausdorff measures for fractional dimension. Observe that by the previous remarks, the least cardinal of any non-measurable subset of any σ -finite set will be the same as the uniformity invariant for Lebesgue measure. In other words, this paper will be concerned with the uniformity invariant of the ideal of σ -finite sets with respect to a Hausdorff measure.

It will be shown that given any real number r in the interval $(0, 1)$ it is consistent with set theory that every set of reals of size \aleph_1 is Lebesgue measurable yet there is a set of reals of size \aleph_1 which is not a null set with respect to r -dimensional Hausdorff measure. This answers Question FQ from D. Fremlin’s list of open questions [2]. However, the motivation was an attempt to resolve the following question posed by P. Komjáth.

Question 1.1. Suppose that every set of size \aleph_1 has Lebesgue measure zero. Does it follow that the union of any set of \aleph_1 lines in the plane has Lebesgue measure zero?

It is worth noting that this is really a geometric question since [9] provides a negative answer to the version of the problem in which lines are replaced by their topological and measure theoretic equivalents. To see the relationship between Question 1.1 and the topic of this paper consider that it is easy to find countably many unit squares in the plane such that each line passes through either the top and bottom or the left and right sides of at least one of these squares. It is therefore possible to focus attention on all lines which pass through the top and bottom of the unit square. For any such line L there is a pair (a, b) such that both the points $(a, 0)$ and $(1, b)$ belong to L . If the mapping which sends a line L to this pair (a, b) is denoted by β then it is easy to see that β is continuous and that if $S \subseteq [0, 1]^2$ is a square of side ε then the union of $\beta^{-1}S$ has measure ε while S itself has measure ε^2 . In other words, the Lebesgue measure of the union of the lines belonging to $\beta^{-1}X$ is no larger than the one-dimensional Hausdorff measure of X for any $X \subseteq [0, 1]^2$. In other words, a negative answer to Question 1.1 would imply that there is $X \subseteq [0, 1]^2$ of size \aleph_1 which is not null with respect to linear Hausdorff measure even though every set of reals of size \aleph_1 is null. The consistency of this will be a consequence of Corollary 6.2.

The proof will rely partially on arguments from [7,8] in which a single stage of a forcing iteration that would achieve the desired model was described. The material in Sections 3 and 4 is a reorganized and simplified version of Sections 4–6 of [8] which has been suitably modified for the current context. The approach taken here differs

from the earlier attempt in that the forcing used is finite branching rather than infinite branching as in [8] and this allows the use of product forcing along lines similar to those in [3]. The new ingredient needed is described in Section 5. The arguments presented in Section 6 use ideas explained in greater detail in [6], however familiarity with that paper is not required in order to follow the reasoning presented here.

To be somewhat more precise, the forcing used consists of creatures (as defined in [6]) which consist of approximations to a cover of the ground model reals by an open set of small Lebesgue measure—each branching node consists of finite union of intervals, the some of whose lengths is less than some prescribed value. This is easily seen to guarantee that the ground model reals form a null set, but another mechanism is required to ensure that they do not form a set of finite Hausdorff- s measure for some fixed $s < 1$. This mechanism consists of a norm, denoted by ν in Section 4, which guarantees that all approximations to a name for an open set witnessing that the ground model has small Hausdorff- s measure can be localized sufficiently well. For example, it will certainly be required of a creature that its open sets form a cover because, otherwise, the generic open set obtained from the forcing will not generically cover the ground model. Hence an assignment to each member of the creature of an open set of small Hausdorff- s capacity should allow a thinning out of the creature so that the sets of small Hausdorff- s capacity are well approximated while maintaining that the creature still covers the reals. This is essentially the content of the definition of having norm at least 1. Higher norms are simply combinatorial product constructions allowing this property to be redundantly encoded in the creature so that it is preserved when decreasing the norm. This was done in [8] using infinite branching trees. The defect of this approach is that it does not allow a product construction such as in [3] to be used. The approach presented in this paper uses a compactness argument which does allow finite branching trees to be used. The compactness argument can be applied only to the norm denoted by ν^∞ in Section 4.

The logical requirements guiding the organization of the paper may not have resulted in optimal organization for the purposes of comprehension. Some readers may prefer to start with Section 6 after having read Section 2, Definitions 3.2 and 4.2. The main point to keep in mind is that the norm ν is the one which will be used in establishing the key result Lemma 6.2. However, ν^∞ is needed for the compactness argument and ρ is the form of the norm which allows the combinatorics of Section 3 to be applied.

2. Notation

If $X \subseteq \mathbb{R}$ and $r \in (0, 1)$ then the infimum of all $\sum_{i=0}^\infty (b_i - a_i)^r$ where $\{(a_i, b_i)\}_{i=0}^\infty$ is a cover of X by intervals of length less than ε is often denoted by $\mathcal{H}_\varepsilon^r(X)$. The r -dimensional Hausdorff capacity of X is denoted by $\mathcal{H}_\infty^r(X)$ and is defined to be the infimum of all $\sum_{i=0}^\infty (b_i - a_i)^r$ where $\{(a_i, b_i)\}_{i=0}^\infty$ is a cover of X by arbitrary intervals.

The r -dimensional Hausdorff measure of a set X is denoted by $\mathcal{H}^r(X)$ and, when defined, is equal to its outer measure $\lim_{\varepsilon \rightarrow 0} \mathcal{H}_\varepsilon^r(X)$. Since it can be shown⁴ that $\mathcal{H}_\infty^r(X) = 0$ if and only if $\mathcal{H}^r(X) = 0$, in order to establish the main result it suffices to show that it is consistent with set theory that every set of reals of size \aleph_1 is a Lebesgue null set yet there is a set of reals of size \aleph_1 which is not a null set with respect to r -dimensional Hausdorff capacity. Actually, it will be more convenient to work with measures and capacities on the Cantor set and to replace intervals by dyadic intervals. This amounts to dealing with net measures as described in [5]. Since r -dimensional net measures have the same null sets as r -dimensional Hausdorff measures this will not be of significance to the results of this paper.

Notation 2.1. For the rest of this paper let r be a fixed real number such that $0 < r < 1$.

Notation 2.2. Let \mathbb{C} denote $2^{\mathbb{N}}$ with the usual product topology. Let \mathbb{Q} be the tree

$$\{f \upharpoonright n : f \in \mathbb{C} \text{ and } n \in \mathbb{N}\}$$

and for $a \in \mathbb{Q}$ let $[a] = \{x \in \mathbb{C} : a \subseteq x\}$ and for $A \subseteq \mathbb{Q}$ let $[A] = \bigcup_{a \in A} [a]$. Let

$$\mathbb{C}_\varepsilon^\infty = \left\{ A \subseteq \mathbb{Q} : \sum_{a \in A} 2^{-|a|r} \leq \varepsilon \right\}$$

and let $\mathbb{C}_\varepsilon = \{A \in \mathbb{C}_\varepsilon^\infty : |A| < \aleph_0\}$. For consistency of notation, let $[\mathbb{Q}]^{< \aleph_0}$ be denoted by \mathbb{C}_∞ . Let $\mathbb{C}_\varepsilon^1 = \{A \subseteq \mathbb{Q} : \sum_{t \in A} 2^{-|t|} \leq \varepsilon\}$. For $X \subseteq \mathbb{C}$ define $\lambda_\infty^r(X) = \inf\{\varepsilon : (\exists A \in \mathbb{C}_\varepsilon^\infty) X \subseteq [A]\}$. The usual product measure on \mathbb{C} will be denoted by λ .

Notation 2.3. Some notation concerning trees will be established. By a sequence will always be meant a function $f : n \rightarrow X$ where $n \in \mathbb{N}$ and X is some set. Sequences will occasionally be denoted as n -tuples (x_1, x_2, \dots, x_n) and, in particular, singleton sequences will be denoted by (x) . If t and s are sequences the concatenation of s followed by t will be denoted by (s, t) . This is consistent with considering X^k to be a set of sequences because if $x \in X^k$ and $y \in Y^m$ then $(x, y) \in X^k \times Y^m$. If T is a tree then T is a set of sequences closed under restriction to initial segments. If $t \in T$ then $\mathcal{S}_T(t) = \{x : (t, x) \in T\}$. Furthermore, $T \langle t \rangle = \{s \in T : t \subseteq s \text{ or } s \subseteq t\}$. If $m \in \mathbb{N}$ then $T[m] = \{t \in T : |t| = m\}$ and $T[< m] = \{t \in T : |t| < m\}$ and $T[\leq m] = \{t \in T : |t| \leq m\}$.

Notation 2.4. For any $X \subseteq \mathbb{C}^m$ and $z \in \mathbb{C}^k$ the set $\{x \in \mathbb{C}^{m-k} : (z, x) \in X\}$ will be denoted by $X(z, \cdot)$. Similarly, if $F : \mathbb{C}^m \rightarrow X$ is a function and $z \in \mathbb{C}^k$ then $F(z, \cdot)$ will represent the function $F(z, \cdot) : \mathbb{C}^{m-k} \rightarrow X$ defined by $F(z, \cdot)(x) = F(z, x)$.

⁴See Lemma 4.6 in [5] for example.

3. Continuous mappings between Lebesgue and Hausdorff measures

The goal of this section is to exploit the difference between Lebesgue measure and r -dimensional Hausdorff measure. It will be shown that for any continuous function from the reals to the reals there are arbitrary small sets in the sense of Lebesgue measure whose pre-image is as large as desired in the sense of r -dimensional Hausdorff measure. For reasons which will reveal themselves in Section 4 it will also be necessary to consider similar results for products of the reals.

Definition 3.1. If $\delta > 0$ is a real number and $X \subseteq \mathbb{C}$ define $\square_\delta(X)$ by

$$\square_\delta(X) = \inf \{ \lambda_\infty^r(X \setminus Z) : Z \subseteq \mathbb{C} \text{ and } \lambda(Z) < \delta \}.$$

Lemma 3.1. *If $1 > \gamma > 0$ and $\varepsilon > 0$ then there is $\eta > 0$ such that for any measurable $E \subseteq \mathbb{C}$ and there is $m_0 \in \mathbb{N}$ such that for any measurable $D \subseteq E$ and for all $m \geq m_0$:*

$$\square_\eta(D) \geq \square_\gamma(E)$$

provided that $\lambda(D \cap [s]) \geq \lambda(E \cap [s])\varepsilon$ for each $s \in \mathbb{Q}[m]$.

Proof. Let $\eta > 0$ be sufficiently small that

$$\frac{\gamma}{2 - \gamma} > \frac{4\eta}{\varepsilon\gamma}$$

and then let $m \in \mathbb{N}$ be so large that the inequality

$$\frac{2^{m(1-r)}\varepsilon^r\gamma^r}{4^r} \left(\frac{\gamma}{2 - \gamma} - \frac{4\eta}{\varepsilon\gamma} \right) > \square_\gamma(E)$$

is satisfied.

Suppose that $\lambda(Z) < \eta$ and $A \subseteq \mathbb{Q}$ are such that $D \subseteq Z \cup [A]$ and

$$\sum_{a \in A} 2^{-|a|r} < \square_\gamma(E)$$

and note that, without loss of generality, it may be assumed that $Z \subseteq D$. Let $A^* = \mathbb{Q}[\leq m] \cap A$ and let $E^* = E \setminus [A^*]$. Let $B_0 = \{b \in \mathbb{Q}[m] : [b] \cap [A^*] = \emptyset\}$ and note that $E^* \subseteq [B_0]$ and that if $b \in B_0$ then $E \cap [b] = E^* \cap [b]$. Now let

$$B_1 = \left\{ b \in B_0 : \lambda(E \cap [b]) > \frac{1}{2} \frac{\lambda(E^*)}{2^m} \right\}.$$

Notice that $\lambda(E^*) \geq \gamma$ because $\sum_{a \in A^*} 2^{-|a|r} < \square_\gamma(E)$ and $E \subseteq E^* \cup [A^*]$. It follows from a Fubini argument that

$$|B_1| \geq \frac{2^m \lambda(E^*)}{2 - \lambda(E^*)} \geq \frac{2^m \gamma}{2 - \gamma}. \tag{3.1}$$

Next, let

$$B_2 = \left\{ b \in B_1 : \lambda(Z \cap [b]) > \frac{\varepsilon \lambda(E \cap [b])}{2} \right\}$$

and in order to see that

$$|B_2| \leq \frac{2^{m+2} \eta}{\varepsilon \gamma} \tag{3.2}$$

assume the opposite. Then the following sequence of inequalities

$$\eta > \lambda(Z) \geq \sum_{b \in B_2} \lambda(Z \cap [b]) \geq \frac{\varepsilon}{2} \sum_{b \in B_2} \lambda(E \cap [b]) \geq \varepsilon \sum_{b \in B_2} \frac{\lambda(E^*)}{2^{m+2}} \geq |B_2| \frac{\varepsilon \gamma}{2^{m+2}} \geq \eta$$

yields a contradiction. (The fourth inequality uses that $B_2 \subseteq B_1$.)

Let $B_3 = B_1 \setminus B_2$ and observe that it follows from Inequalities 3.1 and 3.2 that

$$|B_3| \geq \frac{2^m \gamma}{2 - \gamma} - \frac{2^{m+2} \eta}{\varepsilon \gamma} = 2^m \left(\frac{\gamma}{2 - \gamma} - \frac{4\eta}{\varepsilon \gamma} \right). \tag{3.3}$$

It follows that

$$\begin{aligned} \sum_{a \in A} 2^{-|a|r} &\geq \sum_{b \in B_3} \left(\sum_{a \in A \setminus A^*, a \supseteq b} 2^{-|a|r} \right) \geq \sum_{b \in B_3} (\lambda(D \cap [b] \setminus Z))^r \\ &\geq \sum_{b \in B_3} (\lambda(D \cap [b]) - \lambda(Z \cap [b]))^r. \end{aligned}$$

Since $B_3 \cap B_2 = \emptyset$ it follows, using the hypothesis on D , that the last term dominates

$$\begin{aligned} \sum_{b \in B_3} (\lambda(E \cap [b]) \varepsilon - (\varepsilon/2) \lambda(E \cap [b]))^r &\geq \frac{\varepsilon^r}{2^r} \sum_{b \in B_3} (\lambda(E \cap [b]))^r \geq \frac{\varepsilon^r}{2^r} \sum_{b \in B_3} \left(\frac{\lambda(E^*)}{2^{m+1}} \right)^r \\ &\geq \frac{\varepsilon^r}{4^r} |B_3| \left(\frac{\lambda(E^*)}{2^m} \right)^r \geq \frac{2^{m(1-r)} \varepsilon^r \gamma^r}{4^r} \left(\frac{\gamma}{2 - \gamma} - \frac{4\eta}{\varepsilon \gamma} \right) \end{aligned}$$

and hence $\sum_{a \in A} 2^{-|a|r} > \square_\gamma(E)$ which is impossible. \square

If $X \subseteq \mathbb{C}$ then $F : X \rightarrow \mathbb{C}$ will be said to have small fibres if and only if $\lambda(F^{-1}\{x\}) = 0$ for each $x \in \mathbb{C}$. The proof of Theorem 3.1 and the lemmas preceding it will rely on

decomposing an arbitrary continuous function into a piece that has small fibres and a piece which has countable range.

Lemma 3.2. *Let $0 < \mu < 1$ and suppose that $\{X_s\}_{s \in \mathbb{Q}}$ is an indexed family of mutually independent $\{0, 1\}$ -valued random variables, each with mean μ . Suppose that $C \subseteq \mathbb{C}$ is a measurable set and that $F_j: C \rightarrow \mathbb{C}$ is a measurable function with small fibres for $1 \leq j \leq n$. For any $\varepsilon > 0$ for all but finitely many $m \in \mathbb{N}$ the probability that*

$$\lambda \left(\bigcap_{j=1}^n \left(\bigcup_{s \in \mathbb{Q}[m], X_s=1} F_j^{-1}[s] \right) \right) > \frac{\mu^n \lambda(C)}{2}$$

is greater than $1 - \varepsilon$.

Proof. This is Lemma 3.1 in [7] or Lemma 6.2 in [8] except that it is stated here for \mathbb{C} rather than $[0, 1]$. \square

Lemma 3.3. *Suppose that*

- $E \subseteq \mathbb{C}$ is a measurable set;
- \mathcal{F} is a finite family of measurable functions with small fibres from E to \mathbb{C} ;
- $\gamma > 0, 1 > \mu > 0$.

Then there is $\eta > 0$ such that for any $\varepsilon > 0$ and for any mutually independent, $\{0, 1\}$ -valued random variables $\{X_s\}_{s \in \mathbb{Q}}$ with mean μ and for all but finitely many $m \in \mathbb{N}$ the probability that the inequality

$$\square_\eta \left(\bigcap_{F \in \mathcal{F}} F^{-1} \left(\bigcup_{s \in \mathbb{Q}[m], X_s=1} [s] \right) \right) \geq \square_\gamma(E) \tag{3.4}$$

holds is greater than $1 - \varepsilon$.

Proof. Let $|\mathcal{F}| = n$. Use Lemma 3.1 to choose $\eta > 0$ and an integer k such that if $D \subseteq E$ is a measurable set such that for each $s \in \mathbb{Q}[k]$

$$\lambda(D \cap [s]) \geq \frac{\mu^n}{2} \lambda(E \cap [s])$$

then $\square_\eta(D) \geq \square_\gamma(E)$. Let $\varepsilon > 0$. Now use Lemma 3.2 to conclude that for each $t \in \mathbb{Q}[k]$ for all but finitely many $m \in \mathbb{N}$ and any mutually independent, $\{0, 1\}$ -valued random variables $\{X_s\}_{s \in \mathbb{Q}[m]}$ with mean μ , the probability that

$$\lambda \left([t] \cap \bigcap_{F \in \mathcal{F}} F^{-1} \left(\bigcup_{s \in \mathbb{Q}[m], X_s=1} [s] \right) \right) \geq \frac{\mu^n}{2} \lambda(E \cap [t])$$

is greater than $1 - \varepsilon 2^{-k}$. Hence the probability that this holds for all $t \in \mathbb{Q}[k]$ is greater than $1 - \varepsilon$ and so the hypothesis on k guarantees that Inequality 3.4 holds with at least the same probability. \square

Lemma 3.4. *If $\eta > 0$ and $M \subseteq \mathbb{C}^{d+1}$ is measurable then so is the mapping from \mathbb{C}^d to \mathbb{R} defined by $x \mapsto \square_\eta(M(x, \cdot))$.*

Proof. Let $B \subseteq M$ be an F_σ such that $\lambda(M \setminus B) = 0$. Let $Z = \{z \in \mathbb{C}^d: \lambda(M(z, \cdot) \setminus B(z, \cdot)) > 0\}$ noting that, of course, $\lambda(Z) = 0$. For $x \in \mathbb{C}^d \setminus Z$ observe that $\square_\eta(M(x, \cdot)) = \square_\eta(B(x, \cdot))$. Hence it suffices to show that the mapping $x \mapsto \square_\eta(B(x, \cdot))$ is measurable. In other words, it must be shown that $\{x: \square_\eta(B(x, \cdot)) < \xi\}$ is measurable for any $\xi \in \mathbb{R}$. But $\square_\eta(B(x, \cdot)) < \xi$ if and only if there are sequences $\{a_j\}_{j=0}^\infty \subseteq \mathbb{Q}$ and $\{b_j\}_{j=0}^\infty \subseteq \mathbb{Q}$ such that

$$\sum_{j=0}^\infty 2^{-|a_j|} < \eta \quad \text{and} \quad \sum_{j=0}^\infty 2^{-|b_j| r} < \xi,$$

$$B(x, \cdot) \subseteq \bigcup_{j=0}^\infty [a_j \cup b_j]$$

and, since B is an F_σ , this shows that $\{x: \square_\eta(B(x, \cdot)) < \xi\}$ is Σ_1^1 and, hence, measurable. \square

Corollary 3.1. *Suppose that C is a measurable subset of \mathbb{C}^{d+1} and \mathcal{F} is a finite family of measurable functions from C to \mathbb{C} such that $F(x, \cdot)$ has small fibres for each $x \in \mathbb{C}^d$ and $F \in \mathcal{F}$. If $\mu > 0$ and $\gamma > 0$ then there is $\eta > 0$ such that for all $\varepsilon > 0$ there is some $A \in \mathbb{C}_\mu^1$ such that the Lebesgue measure of*

$$\left\{ x \in \mathbb{C}^d: \square_\eta \left(\bigcap_{F \in \mathcal{F}} F(x, \cdot)^{-1}[A] \right) \geq \square_\gamma(C(x, \cdot)) \right\}$$

is at least $1 - \varepsilon$.

Proof. This is a standard application of Fubini’s Theorem using Lemma 3.3 and the Law of Large Numbers once it has been observed that the set in question is measurable. But this follows directly by applying Lemma 3.4 to both sides of the inequality and then taking the difference of measurable functions. \square

Lemma 3.5. *Let $E \subseteq \mathbb{C}$ be a measurable set and \mathcal{F} a finite family of measurable functions from E to \mathbb{C} . Then for any $\gamma > 0$ and any $\mu > 0$ there is $\eta > 0$*

and $A \in \mathbb{C}_\mu^1$ such that

$$\square_\eta \left(\bigcap_{F \in \mathcal{F}} F^{-1}[A] \right) \geq \square_\gamma(E).$$

Proof. Let $|\mathcal{F}| = n$. For each $F \in \mathcal{F}$ let $Y_F = \bigcup \{F^{-1}\{y\} : \lambda(F^{-1}\{y\}) > 0\}$ and let \bar{F} be defined by

$$\bar{F}(z) = \begin{cases} F(z) & \text{if } z \in E \setminus Y_F, \\ z & \text{if } z \in Y_F. \end{cases}$$

Since each \bar{F} is measurable and has small fibres it is possible to use Lemma 3.3 to conclude that there is $\eta > 0$ and an integer m and mutually independent, $\{0, 1\}$ -valued random variables $\{X_s\}_{s \in \mathbb{Q}[m]}$ with mean $\mu/3$ such that the probability that

$$\square_{2\eta} \left(\bigcap_{F \in \mathcal{F}} \bar{F}^{-1} \left(\bigcup_{s \in \mathbb{Q}[m], X_s=1} [s] \right) \right) \geq \square_\gamma(E)$$

is greater than $\frac{1}{2}$.

Since the mean of each X_s is $\mu/3$ it is possible to choose m so large that the probability that

$$\lambda \left(\bigcup_{s \in \mathbb{Q}[m], X_s=1} [s] \right) < \frac{\mu}{2}$$

is also greater than $\frac{1}{2}$. Hence there is $A' \in \mathbb{C}_{\mu/2}^1$ such that

$$\square_{2\eta} \left(\bigcap_{F \in \mathcal{F}} \bar{F}^{-1}[A'] \right) \geq \square_\gamma(E).$$

Now for each $F \in \mathcal{F}$ choose a finite set $A_F \subseteq \mathbb{C}$ such that $\lambda(Y_F \setminus F^{-1}A_F) < \eta/n$. Then let $A \in \mathbb{C}_\mu^1$ be such that $[A'] \cup \bigcup_{F \in \mathcal{F}} A_F \subseteq [A]$. It follows that

$$\square_\eta \left(\bigcap_{F \in \mathcal{F}} F^{-1}[A] \right) \geq \square_\eta \left(\bigcap_{F \in \mathcal{F}} \bar{F}^{-1}[A] \setminus \bigcup_{F \in \mathcal{F}} (Y_F \setminus F^{-1}A_F) \right) \geq \square_\gamma(E)$$

because $F^{-1}[A] \supseteq \bar{F}^{-1}[A] \setminus (Y_F \setminus F^{-1}A_F)$ for each $F \in \mathcal{F}$. \square

For the next definition recall Definition 3.1.

Definition 3.2. If $\Theta : \mathbb{N} \rightarrow \mathbb{R}^+$, $\Gamma : \mathbb{N} \rightarrow \mathbb{R}^+$ and $X \subseteq \mathbb{C}^d$ then the relation $\square_{\Theta, \Gamma}(X)$ will be defined to hold by induction on d . If $d = 1$ then $\square_{\Theta, \Gamma}(X)$ if and only if

$\square_{\Theta(0)}(X) \geq \Gamma(0)$ whereas, if $d > 1$, then $\square_{\Theta, \Gamma}(X)$ holds if and only if $\square_{\Theta(d-1)}(\{x \in \mathbb{C} : \square_{\Theta, \Gamma}(X(x, \cdot))\}) \geq \Gamma(d-1)$. Define $\Theta^+(i) = \Theta(i+1)$ and $\Gamma^+(i) = \Gamma(i+1)$.

The next lemma establishes that the top-down and bottom-up definitions of the relation $\square_{\Theta, \Gamma}$ are the same.

Lemma 3.6. *Let $d \geq 2$. For any $\Theta : \mathbb{N} \rightarrow \mathbb{R}^+$, $\Gamma : \mathbb{N} \rightarrow \mathbb{R}^+$ and $X \subseteq \mathbb{C}^{d+1}$ the relation $\square_{\Theta, \Gamma}(X)$ holds if and only if $\square_{\Theta^+, \Gamma^+}(\{z \in \mathbb{C}^d : \square_{\Theta(0)}(X(z, \cdot)) \geq \Gamma(0)\})$.*

Proof. Proceed by induction on d and observe that the case $d = 2$ is immediate from Definition 3.2. Assuming the lemma established for d let $X \subseteq \mathbb{C}^{d+2}$. Then the following sequence of equivalences establishes the lemma:

$$\begin{aligned} & \square_{\Theta, \Gamma}(X), \\ & \square_{\Theta(d+1)}(\{z \in \mathbb{C}^1 : \square_{\Theta, \Gamma}(X(z, \cdot))\}) \geq \Gamma(d+1), \\ & \square_{\Theta(d+1)}(\{z \in \mathbb{C}^1 : \square_{\Theta^+, \Gamma^+}(\{w \in \mathbb{C}^d : \square_{\Theta(0)}(X(z, \cdot)(w, \cdot)) \geq \Gamma(0)\})\}) \geq \Gamma(d+1), \\ & \square_{\Theta^+(d)}(\{z \in \mathbb{C}^1 : \square_{\Theta^+, \Gamma^+}(\{w \in \mathbb{C}^d : \square_{\Theta(0)}(X((z, w), \cdot)) \geq \Gamma(0)\})\}) \geq \Gamma^+(d), \\ & \square_{\Theta^+, \Gamma^+}(\{(z, w) \in \mathbb{C}^1 \times \mathbb{C}^d : \square_{\Theta(0)}(X((z, w), \cdot)) \geq \Gamma(0)\}), \\ & \square_{\Theta^+, \Gamma^+}(\{z \in \mathbb{C}^{d+1} : \square_{\Theta(0)}(X(z, \cdot)) \geq \Gamma(0)\}). \quad \square \end{aligned}$$

Lemma 3.7. *If $\square_{\Theta, \Gamma}(X)$ holds and $X \subseteq \mathbb{C}^d$ and $\eta < \Theta(i)$ for each $i < d$ and $A \subseteq \mathbb{C}^d$ is such that $\lambda(A) < \eta^d$ then $\square_{\Theta-\eta, \Gamma}(X \setminus A)$ holds where $(\Theta - \eta)(i) = \Theta(i) - \eta$.*

Proof. Proceed by induction on d using Fubini’s Theorem. \square

Theorem 3.1. *Let $\Gamma : \mathbb{N} \rightarrow \mathbb{R}^+$ and $\Theta : \mathbb{N} \rightarrow \mathbb{R}^+$ be functions such that $\Theta(i) + \Gamma(i) < 1$ for each i . Suppose that C is a closed subset of \mathbb{C}^d such that $\square_{\Theta, \Gamma}(C)$ holds and that \mathcal{F} is a finite family of continuous functions from C to \mathbb{C} . If $\mu > 0$ then there is $A \in \mathbb{C}_\mu^1$ and $\eta : \mathbb{N} \rightarrow \mathbb{R}^+$ such that $\square_{\eta, \Gamma}(\bigcap_{F \in \mathcal{F}} F^{-1}[A])$.*

Proof. Proceed by induction on d , noting that if $d = 1$ then this follows from Lemma 3.5 by setting $\gamma = \Theta(0)$ in that lemma. So assume that the lemma has been

established for d and that C is a closed subset of \mathbb{C}^{d+1} , $\square_{\theta, \Gamma}(C)$ holds and \mathcal{F} is a finite family of continuous functions from C to \mathbb{C} and that $\mu > 0$. For each $F \in \mathcal{F}$ let

$$Y_F = \{(x, y) \in \mathbb{C}^d \times \mathbb{C} : \lambda(\{z \in \mathbb{C} : F(x, y) = F(x, z)\}) > 0\}$$

and note that Y_F is closed since F is continuous. Now, for $F \in \mathcal{F}$ define $\bar{F} : C \rightarrow \mathbb{C}$ by

$$\bar{F}(x, y) = \begin{cases} F(x, y) & \text{if } (x, y) \notin Y_F, \\ y & \text{otherwise.} \end{cases}$$

Observe that $\bar{F}(x, \cdot)$ has small fibres for each $x \in \mathbb{C}^d$. By Corollary 3.1 there is $\delta > 0$ such that for all $\varepsilon > 0$ there is some $A \in \mathbb{C}_{\mu/2}^1$ such that the Lebesgue measure of

$$B(A) = \left\{ x \in \mathbb{C}^d : \square_{\delta} \left(\bigcap_{F \in \mathcal{F}} \bar{F}(x, \cdot)^{-1}[A] \right) \geq \square_{\theta(0)}(C(x, \cdot)) \right\}$$

is at least $1 - \varepsilon$.

Since each of the relations Y_F is Borel, it is possible to appeal to Blackwell's Selection Theorem⁵ to find a finite family of Borel functions \mathcal{G} from \mathbb{C}^d to \mathbb{C} such that

$$\sum_{F \in \mathcal{F}} \int_{x \in \mathbb{C}^d} \lambda(Y_F(x, \cdot) \setminus F(x, \cdot)^{-1}\{G(x)\}_{G \in \mathcal{G}}) dx < \frac{\delta^{d+1}}{2^{d+1}}.$$

It is then possible to find a closed set $D \subseteq \mathbb{C}^d$ such that

- (1) $(1 - \lambda(D))^{1/d} < \theta(i + 1)$ for each $i < d$;
- (2) $\lambda(D) > 1 - \delta^d / 2^d$;
- (3) $|\mathcal{F}| \cdot \lambda(Y_F(x, \cdot) \setminus F(x, \cdot)^{-1}\{G(x)\}_{G \in \mathcal{G}}) < \delta / 2$ for each $F \in \mathcal{F}$ and $x \in D$;
- (4) each $G \in \mathcal{G}$ is continuous on D .

Letting $C^* = \{z \in \mathbb{C}^d : \square_{\theta(0)}(C(z, \cdot)) \geq \Gamma(0)\}$ it follows that

$$\square_{\theta^+ - (1 - \lambda(D))^{1/d}, \Gamma^+}(D \cap C^*)$$

holds because Lemma 3.6 implies that $\square_{\theta^+, \Gamma^+}(C^*)$ does. Condition 1 guarantees that $\theta^+(i) > (1 - \lambda(D))^{1/d}$ for each i and C^* is easily seen to be closed. It is therefore possible to apply the induction hypothesis to $\theta^+ - (1 - \lambda(D))^{1/d}$, Γ^+ , \mathcal{G} and $D \cap C^*$ to get $A_0 \in \mathbb{C}_{\mu/2}^1$ and $\eta : \mathbb{N} \rightarrow \mathbb{R}^+$ such that

$$\square_{\eta, \Gamma^+} \left(D \cap C^* \cap \bigcap_{G \in \mathcal{G}} G^{-1}[A_0] \right).$$

⁵See Corollary 18.7 on p. 122 of [4].

Let ι be such that $0 < \iota < \eta(i)$ for $i < d$. Now choose $A_1 \in \mathbb{C}_{\mu/2}^1$ such that $\lambda(B(A_1)) > 1 - \iota^d$. Then

$$\square_{\eta-\iota, \Gamma^+} \left(D \cap C^* \cap B(a_1) \cap \bigcap_{G \in \mathcal{G}} G^{-1}[A_0] \right)$$

holds by Lemma 3.7. Note that by Condition 3

$$\square_{\delta/2} \left(\bigcap_{F \in \mathcal{F}} \bar{F}(x, \cdot)^{-1}[A_1] \setminus \bigcup_{F \in \mathcal{F}} (Y_F(x, \cdot) \setminus F(x, \cdot)^{-1}\{G(x)\}_{G \in \mathcal{G}}) \right) \geq \square_{\Theta(0)}(C(z, \cdot))$$

for every $x \in D \cap B(a_1)$ and, moreover, $\square_{\Theta(0)}(C(z, \cdot)) \geq \Gamma(0)$ for $x \in C^*$. Hence

$$\square_{\delta/2} \left(\bigcap_{F \in \mathcal{F}} \bar{F}(x, \cdot)^{-1}[A_1] \setminus \bigcup_{F \in \mathcal{F}} (Y_F(x, \cdot) \setminus F(x, \cdot)^{-1}\{G(x)\}_{G \in \mathcal{G}}) \right) \geq \Gamma(0)$$

for every $x \in D \cap C^* \cap B(a_1)$. Let $\bar{\eta}$ be defined by

$$\bar{\eta}(i) = \begin{cases} \eta(i-1) - \iota & \text{if } d \geq i > 0, \\ \eta(i-1) & \text{if } d < i, \\ \delta/2 & \text{if } i = 0 \end{cases}$$

and define Z to be the set of all $(x, w) \in \mathbb{C}^{d+1}$ such that the following three conditions are satisfied:

$$x \in D \cap C^* \cap B(a_1) \cap \bigcap_{G \in \mathcal{G}} G^{-1}[A_0] \tag{3.5}$$

$$(\forall F \in \mathcal{F}) \bar{F}(x, w) \in [A_1] \tag{3.6}$$

$$(\forall F \in \mathcal{F}) F(x, w) \notin Y_F(x, \cdot) \text{ or } (\exists G \in \mathcal{G}) F(x, w) = G(x) \tag{3.7}$$

It follows from Lemma 3.6 that $\square_{\bar{\eta}, \Gamma}(Z)$. Hence it suffices to observe that if $(x, w) \in Z$ then $F(x, w) \in [A_0 \cup A_1]$. To see this note that if $F(x, w) = \bar{F}(x, w)$ this is immediate from 3.6. Otherwise $F(x, w) \in Y_F$ and hence it follows from 3.7 that there is some $G \in \mathcal{G}$ such that $F(x, w) = G(x)$. From 3.5 it can be concluded that $G(x) \in [A_0]$. \square

Notation 3.1. For the rest of the paper, fix a pair of functions $\Gamma : \mathbb{N} \rightarrow \mathbb{R}^+$ and $\Theta : \mathbb{N} \rightarrow \mathbb{R}^+$ such that

$$\Theta(n) > \Theta(n+1) \text{ and } \lim_{n \rightarrow \infty} \Theta(n) = 0, \tag{3.8}$$

$$(\forall j) \Theta(j) + \Gamma(j) < 1, \tag{3.9}$$

$$(\forall j) \Theta(j) + \Gamma(j + 1) < 1, \tag{3.10}$$

$$\text{the range of } \Gamma \text{ is a dense subset of } (0, 1). \tag{3.11}$$

Definition 3.3. For any $d \in \mathbb{N}$ and for $X \subseteq \mathbb{C}^d$ define $\square(X) = \square_{\Theta, \Gamma}(X)$ and define $\square_*(X)$ to hold if and only if there is some $\eta : \mathbb{N} \rightarrow \mathbb{R}^+$ such that $\square_{\eta, \Gamma}(X)$ holds.

4. A preliminary norm

This section will introduce three norms⁶ on subsets of \mathbb{C}_∞ which will be denoted by ρ , v^∞ and v . The norms of [6] typically enjoy some form of sub-additivity but this will not be the case for any of these three, at least not explicitly. Nevertheless, Lemma 4.2 can be considered a substitute for this. The only norm used in the definition of the partial order in Section 6 will be v . The role of the norm ρ will be to establish a connection between v^∞ and the results of Section 3. The norm v^∞ is an intermediary between ρ and v and, furthermore, it has the advantage of allowing the compactness argument of Section 5 to work.

Notation 4.1. For any Polish space X let $\mathcal{K}(X)$ denote the space of compact subsets of X with the Hausdorff metric and let $\mathcal{C}(X)$ denote the space of continuous \mathbb{C} -valued functions with the uniform metric.

Definition 4.1. Define ρ to be a function from $\mathcal{P}(\mathbb{C}_\infty)$ to $\mathbb{N} \cup \{\infty\}$ by first defining $\rho^*(\mathcal{X})$ to be

$$\{d \geq 1 : (\forall C \in \mathcal{K}(\mathbb{C}^d))(\forall F \in \mathcal{C}(C))(\exists x \in \mathcal{X}) \square(C) \Rightarrow \square_*(F^{-1}x)\}$$

and let $\rho(\mathcal{X}) = \sup(\rho^*(\mathcal{X}))$. If $\rho^*(\mathcal{X}) = \emptyset$ but $\mathcal{X} \neq \emptyset$ then define $\rho(\mathcal{X}) = 0$.

Observe that each set $\rho^*(\mathcal{X})$ is an initial segment of the integers. In order to see this note that if $d + 1 \in \rho^*(\mathcal{X})$ and $C \in \mathcal{K}(\mathbb{C}^d)$ and $F \in \mathcal{C}(C)$ then, by considering $F' : \mathbb{C} \times C \rightarrow \mathbb{C}$ defined by $F'(x, c) = F(c)$, it is easily shown that $d \in \rho^*(\mathcal{X})$. The following is a direct Corollary of Theorem 3.1.

Corollary 4.1. If $\mu > 0$ then $\rho(\mathbb{C}_\mu^1) = \infty$.

⁶This is the terminology of [6]. The exact meaning of the term “norm” is not required for the arguments used here.

Lemma 4.1. *If $d \geq 1$ and functions θ and γ are given then $\{C \subseteq \mathcal{K}(\mathbb{C}^d) : \square_{\theta, \gamma}(C)\}$ is a closed set.*

Proof. Proceed by induction on d . The case $d = 1$ is easy since an open set witnessing the failure of $\square_{\theta, \gamma}(C)$ witnesses this failure for any of its subsets and, thus, for a neighbourhood of compact sets. Assuming the result for d let $C \subseteq \mathbb{C}^{d+1}$ be compact and suppose that $\square_{\theta, \gamma}(C)$ fails. Using Definition 3.2, find $\mathcal{C}_1 \in \mathcal{C}_{\bar{\gamma}}$ for some $\bar{\gamma} < \gamma(d)$ and $\mathcal{B}_1 \in \mathcal{C}_{\bar{\theta}}^1$ for some $\bar{\theta} < \theta(d)$ such that $\{x \in \mathbb{C} : \square_{\theta, \gamma}(C(x, \cdot))\} \subseteq [\mathcal{B}_1 \cup \mathcal{C}_1]$.

For $x \in \mathbb{C} \setminus [\mathcal{B}_1 \cup \mathcal{C}_1]$ it is possible to use the induction hypothesis to find a finite, and perhaps empty, \mathcal{C}_x such that $C(x, \cdot) \subseteq [\mathcal{C}_x]$ and $\square_{\theta, \gamma}(B)$ fails for any $B \subseteq [\mathcal{C}_x]$. The compactness of C yields an integer k_x such that $C(y, \cdot) \subseteq [\mathcal{C}_x]$ for each $y \in \mathbb{C}$ such that $y \upharpoonright k_x = x \upharpoonright k_x$. Choose a finite $X \subseteq \mathbb{C}$ such that $\{[x \upharpoonright k_x]\}_{x \in X}$ covers $\mathbb{C} \setminus [\mathcal{B}_1 \cup \mathcal{C}_1]$. It is then easy to see that for any

$$C' \subseteq ([\mathcal{B}_1 \cup \mathcal{C}_1] \times \mathbb{C}) \cup \left(\bigcup_{x \in X} [x \upharpoonright k_x] \times [\mathcal{C}_x] \right),$$

$\square_{\theta, \gamma}(C')$ fails. This provides a neighbourhood of C disjoint from $\{C' \subseteq \mathcal{K}(\mathbb{C}^d) : \square_{\theta, \gamma}(C')\}$ as required. \square

Corollary 4.2. *For any $d \geq 1$ the set $\{C \subseteq \mathcal{K}(\mathbb{C}^d) : \square(C)\}$ is closed and $\{C \subseteq \mathcal{K}(\mathbb{C}^d) : \square_*(C)\}$ is an F_σ .*

Proof. For the first assertion simply apply Lemma 4.1 with $\theta = \Theta$ and $\gamma = \Gamma$. For the second use Definition 3.1 to see that $\square_*(C)$ holds if and only if there is some $\eta : \mathbb{N} \rightarrow \mathbb{R}^+$ such that $\square_{\eta, \Gamma}(C)$ holds. But notice that η can be assumed to be a constant function with rational value. It follows immediately from Lemma 4.1 that $\{C \subseteq \mathcal{K}(\mathbb{C}^d) : \square_*(C)\}$ is an F_σ . \square

Notation 4.2. The notation $\langle z, F \rangle$ will be introduced for any $z \in \mathbb{C}$ and any function $F : W \rightarrow \mathcal{P}(\mathbb{Q})$ to denote $\{w \in W : z \notin [F(w)]\}$.

Corollary 4.3. *Suppose that $\mathcal{X} \subseteq \mathbb{C}_\infty$ and $B : \mathcal{X} \rightarrow \mathcal{P}(\mathbb{C}_\infty)$. Then $\{z \in \mathbb{C} : \rho(\langle z, B \rangle) < j\}$ is analytic for any integer j .*

Proof. From Corollary 4.2 it follows that the assertion

$$(\exists x \in \mathcal{X}) \square(C) \Rightarrow \square_*(F^{-1}x)$$

is Borel for $\mathcal{X} \in \mathcal{P}(\mathbb{C}_\infty)$. Hence the corollary follows from Definition 4.1. \square

Lemma 4.2. *Suppose that $\mathcal{X} \subseteq \mathbb{C}_\infty$ and $\rho(\mathcal{X}) = j \geq 2$. Then*

$$\square_{\theta(j-2)}(\{z \in \mathbb{C} : \rho(\langle z, F \rangle) < j - 1\}) < \Gamma(j - 1)$$

for each function F from \mathcal{X} to $\mathbb{C}_{\Gamma(j-1)/2}^\infty$.

Proof. Let $S = \{z \in \mathbb{C} : \rho(\langle z, F \rangle) < j - 1\}$ and assume that the lemma fails. In other words, $\square_{\theta(j-2)}(S) \geq \Gamma(j - 1)$. Let

$$E \subseteq \mathbb{C} \times \mathcal{K}(\mathbb{C}^{j-1}) \times \mathcal{C}(\mathbb{C}^{j-1})$$

be the set of all triples (z, C, f) such that

$$\square(C), \tag{4.1}$$

$$(\forall x \in \langle z, F \rangle) \neg \square_*(C \cap f^{-1}x). \tag{4.2}$$

It follows from Definition 4.1 that S is contained in the domain of E . From Lemma 4.1 and Corollary 4.2 it follows that E is a Borel set. From Corollary 4.3 it follows that S is measurable and so it is possible to use the von Neumann Selection Theorem and Lusin’s Theorem to find a closed set $S^* \subseteq S$ and a continuous function $T \subseteq E$ with domain S^* such that $\lambda(S \setminus S^*) < \theta(j - 2) - \theta(j - 1)$. Let $T(s) = (C_s, f_s)$ and observe that the continuity of T guarantees that $C^* = \bigcup_{s \in S^*} \{s\} \times C_s$ is a closed set. Moreover, $\square_{\theta(j-1)}(S^*) \geq \square_{\theta(j-2)}(S) \geq \Gamma(j - 1)$. A calculation using Definition 3.2 reveals that $\square(C^*)$ holds. Let g be defined on C^* by $g(c_1, c_2, \dots, c_j) = f_{c_1}(c_2, c_3, \dots, c_j)$ recalling that $j \geq 2$.

Since $\rho(\mathcal{X}) \geq j$ it is possible to find $x \in \mathcal{X}$ such that $\square_*(g^{-1}x)$. Let

$$W = \{z \in S^* : \square_*(g^{-1}x(z, \cdot))\} = \{z \in S^* : \square_*(g(z, \cdot)^{-1}x)\}$$

and note that it follows that $\lambda'_\infty(W) \geq \Gamma(j - 1)$. Since $F(x) \in \mathbb{C}_{\Gamma(j-1)/2}$ it is possible to choose $w \in W \setminus [F(x)]$. Then $x \in \langle w, F \rangle$ and so, by Condition 4.2 in the definition of E , it follows that $\square_*(g(w, \cdot)^{-1}x)$ fails since $g(w, \cdot) = f_w$. This contradicts that $w \in W$. \square

Definition 4.2. The norms v and v^∞ will be defined for the subsets of \mathbb{C}_∞ by first using induction to define an associated sequence of sets:

$$\mathcal{N}_0 = \mathcal{N}_0^\infty = \{\mathcal{X} \subseteq \mathbb{C}_\infty : \mathcal{X} \neq \emptyset\},$$

$$\mathcal{N}_1 = \mathcal{N}_1^\infty = \{\mathcal{X} \subseteq \mathbb{C}_\infty : [\cup \mathcal{X}] = \mathbb{C}\},$$

$$\mathcal{N}_{j+1} = \{\mathcal{X} \subseteq \mathbb{C}_\infty : (\forall F : \mathcal{X} \rightarrow \mathbb{C}_{\Gamma(j)/2}) \square_{\theta(j-1)}(\{z \in \mathbb{C} : \langle z, F \rangle \notin \mathcal{N}_j\}) < \Gamma(j)\}, \tag{4.3}$$

$$\mathcal{N}_{j+1}^\infty = \{\mathcal{X} \subseteq \mathbb{C}_\infty : (\forall F : \mathcal{X} \rightarrow \mathbb{C}_{\Gamma(j)/2}^\infty) \square_{\theta(j-1)}(\{z \in \mathbb{C} : \langle z, F \rangle \notin \mathcal{N}_j^\infty\}) < \Gamma(j)\}. \tag{4.4}$$

Then define $v(\mathcal{X})$ to be the supremum of all j such that $\mathcal{X} \in \mathcal{N}_j$ and $v^\infty(\mathcal{X})$ to be the supremum of all j such that $\mathcal{X} \in \mathcal{N}_j^\infty$.

It must be noted that $\mathcal{N}_j \supseteq \mathcal{N}_{j+1}$ for each integer j and, indeed, if \mathcal{X} belongs to \mathcal{N}_{j+1} then it belongs to \mathcal{N}_j in spades. To see this first observe that an easy inductive argument, using nothing more than de Morgan’s Rule, shows that each family \mathcal{N}_j is closed under supersets. Since $\Gamma(j) + \Theta(j - 1) < 1$ by Inequality 3.10, it follows that if $\mathcal{X} \in \mathcal{N}_{j+1}$ then taking any $F : \mathcal{X} \rightarrow \mathbb{C}_{\Gamma(j)/2}$ there is at least one $z \in \mathbb{C}$ such that $\langle z, F \rangle \in \mathcal{N}_j$. Since $\langle z, F \rangle \subseteq \mathcal{X}$ it is immediate that $\mathcal{X} \in \mathcal{N}_j$. Similar reasoning shows that $\mathcal{N}_j^\infty \supseteq \mathcal{N}_{j+1}^\infty$ as well. Hence the supremum in Definition 4.2 is taken over an initial segment of the integers. This will be used implicitly in what follows.

Corollary 4.4. *If $\mathcal{X} \subseteq \mathbb{C}_\infty$ then $\rho(\mathcal{X}) \leq v^\infty(\mathcal{X})$.*

Proof. Proceed by induction on $j = \rho(\mathcal{X})$. The case $j = 0$ is trivial but the case $j = 1$ is less so. To see that $\rho(\mathcal{X}) = 1$ implies that \mathcal{X} covers \mathbb{C} let $z \in \mathbb{C}$. Note that $\square(\mathbb{C})$ holds by Condition 3.10 of Notation 3.1. Letting F be the function on \mathbb{C} with constant value z it follows from the definition of $\rho(\mathcal{X}) = 1$ that there is some $x \in \mathcal{X}$ such that $\square_*(F^{-1}x)$ holds. In particular, $F^{-1}x \neq \emptyset$ and so if $w \in F^{-1}x$ then $z = F(w) \in x$.

Therefore, it can be assumed that $2 \leq j + 1 = \rho(\mathcal{X})$ and that the lemma holds for j . In order to show that $v^\infty(\mathcal{X}) \geq j + 1$ let $F : \mathcal{X} \rightarrow \mathbb{C}_{\Gamma(j)/2}^\infty$. By Lemma 4.2 it must be that

$$\square_{\Theta(j-1)}(\{z \in \mathbb{C} : \rho(\langle z, F \rangle) < j\}) < \Gamma(j).$$

From the induction hypothesis it follows that if $v^\infty(\langle z, F \rangle) < j$ then $\rho(\langle z, F \rangle) < j$. Hence

$$\square_{\Theta(j-1)}(\{z \in \mathbb{C} : v^\infty(\langle z, F \rangle) < j\}) < \Gamma(j)$$

and this establishes that $v^\infty(\mathcal{X}) \geq j + 1$. \square

5. Finding finite sets with large norm

It will be shown that for any j there is a finite \mathcal{X} such that $v^\infty(\mathcal{X}) \geq j$. This will establish that the $\mathfrak{C}(k)$ required in the definition of the partial order \mathbb{P} in Section 6 actually do exist. Each of the next lemmas is a step towards this goal.

Lemma 5.1. *If $\mathcal{X} \subseteq \mathbb{C}_\infty$ then $v^\infty(\mathcal{X}) \leq v(\mathcal{X})$.*

Proof. This follows from the definitions by an argument using induction on j to show that if $v^\infty(\mathcal{X}) \geq j + 1$ then $v(\mathcal{X}) \geq j + 1$. \square

Lemma 5.2. *If $\{\mathcal{A}_n\}_{n=0}^\infty$ is an increasing sequence of finite subsets of \mathbb{C}_∞ then*

$$v^\infty\left(\bigcup_{n=0}^\infty \mathcal{A}_n\right) = \lim_{n \rightarrow \infty} v^\infty(\mathcal{A}_n).$$

Proof. Proceed by induction on j to show that if $v^\infty(\mathcal{A}_n) < j$ for each n then $v^\infty(\bigcup_{n=0}^\infty \mathcal{A}_n) < j$. For $j = 0$ this is trivial and if $j = 1$ this is simply a restatement of the compactness of \mathbb{C} . Therefore assume that $j \geq 1$, that the lemma is true for j , that $v^\infty(\mathcal{A}_n) \leq j$ for each n yet $v^\infty(\bigcup_{n=0}^\infty \mathcal{A}_n) > j$. Let $F_n : \mathcal{A}_n \rightarrow \mathbb{C}_{\Gamma(j)/2}^\infty$ witness that $v^\infty(\mathcal{A}_n) \not\geq j + 1$. In other words, using Equality 4.4 of Definition 4.2, $\square_{\Theta(j-1)}(S_n) \geq \Gamma(j)$ where $S_n = \{z \in \mathbb{C} : v^\infty(\langle z, F_n \rangle) < j\}$.

Claim 1. *If $\{\delta_n\}_{n=0}^\infty$ is a sequence of positive reals and $\{A_n\}_{n=0}^\infty$ is a sequence of elements of \mathbb{C}_1^∞ then there are two increasing sequences of integers $\{k_n\}_{n=0}^\infty$ and $\{m_n\}_{n=0}^\infty$ such that, letting*

$$D_n = A_{m_n} \cap \mathbb{Q}[\langle k_n \rangle]$$

the following hold:

$$\text{if } i \geq n \text{ then } D_n = D_i \cap \mathbb{Q}[\langle k_n \rangle], \tag{5.1}$$

$$\lambda\left(\bigcup_{i=n}^\infty [A_{m_i} \setminus D_n]\right) < \delta_n. \tag{5.2}$$

Moreover, the increasing sequence $\{k_n\}_{n=0}^\infty$ can be chosen from any given infinite set K .

Proof. Let K be given and, using the fact that $0 < r < 1$, let $\{k_n\}_{n=0}^\infty \subseteq K$ be such that for each n

$$\sum_{i=n}^\infty 2^{-k_i(1-r)} \leq \delta_n/2$$

and then choose the sequence $\{m_n\}_{n=0}^\infty$ such that Conclusion 5.1 holds. Then Conclusion 5.2 follows from the following inequalities:

$$\begin{aligned} \lambda\left(\bigcup_{i=n}^\infty [A_{m_i} \setminus D_n]\right) &\leq \lambda\left(\bigcup_{i=n}^\infty [A_{m_i} \setminus D_i] \cup [D_{i+1} \setminus D_i]\right) \\ &\leq \sum_{i=n}^\infty \lambda([A_{m_i} \setminus D_i]) + \lambda([D_{i+1} \setminus D_i]) \end{aligned}$$

$$\begin{aligned}
 &\leq \sum_{i=n}^{\infty} \left(\sum_{t \in A_{m_i} \setminus D_i} 2^{-|t|} + \sum_{t \in D_{i+1} \setminus D_i} 2^{-|t|} \right) \\
 &\leq \sum_{i=n}^{\infty} \left(\sum_{t \in A_{m_i} \setminus D_i} 2^{-|t|r} 2^{-|t|(1-r)} + \sum_{t \in D_{i+1} \setminus D_i} 2^{-|t|r} 2^{-|t|(1-r)} \right) \\
 &\leq \sum_{i=n}^{\infty} \left(2^{-k_i(1-r)} \left(\sum_{t \in A_{m_i} \setminus D_i} 2^{-|t|r} + \sum_{t \in D_{i+1} \setminus D_i} 2^{-|t|r} \right) \right) \\
 &\leq \sum_{i=n}^{\infty} 2^{-k_i(1-r)} (1 + 1) \leq \delta_n. \quad \square
 \end{aligned}$$

Before continuing, let $\{a_i\}_{i=0}^{\infty}$ enumerate $\bigcup_{n=0}^{\infty} \mathcal{A}_n$ and, without loss of generality, assume that $\mathcal{A}_n = \{a_\ell\}_{\ell=0}^n$. Using the claim and its final clause and a diagonalization argument, find two increasing sequences of integers $\{k_n\}_{n=0}^{\infty}$ and $\{m_n\}_{n=0}^{\infty}$ such that for each i and each $n \geq i$, letting $F_{n,i} = F_{m_n}(a_i) \cap \mathbb{Q}[\leq k_n]$, Conclusions 5.1 and 5.2 of the claim hold for a summable sequence of δ . To be more precise, $F_{n+1,i}$ is an end extension of $F_{n,i}$ and

$$\lambda \left(\bigcup_{j=n}^{\infty} [F_{m_j}(a_i) \setminus F_{n,i}] \right) < 2^{-i-n}$$

for any $n \geq i$. Define $F(a_i) = \bigcup_{n=i}^{\infty} F_{n,i}$ and note that $F(a_i) \in \mathbb{C}_{\Gamma(j)/2}^{\infty}$ since $F_{n,i} \in \mathbb{C}_{\Gamma(j)/2}$ for each n and the $F_{n,i}$ are increasing with respect to n .

Now let $S = \{z \in \mathbb{C} : v^{\infty}(\langle z, F \rangle) < j\}$. Because

$$v^{\infty} \left(\bigcup_{n=0}^{\infty} \mathcal{A}_n \right) \geq j + 1,$$

it follows that $\square_{\Theta(j-1)}(S) < \Gamma(j)$ and, hence, it is possible to choose Y such that $\lambda'_{\infty}(S \setminus Y) < \Gamma(j)$ and $\lambda(Y) < \Theta(j - 1)$. Choose M so large that letting

$$W = \bigcup_{i=0}^{\infty} \left(\bigcup_{n=\max(M,i)}^{\infty} [F_{m_n}(a_i) \setminus F_{\max(M,i),i}] \right),$$

it follows that $\lambda(W) < \Theta(j - 1) - \lambda(Y)$.

Now define $F^*(a_i)$ by

$$F^*(a_i) = \begin{cases} F_{M,i} & \text{if } i \leq M, \\ F_{i,i} & \text{if } i > M \end{cases}$$

and note that $F^*(a_i) \subseteq F(a_i)$. Now let $S_m^* = \{z \in \mathbb{C} : v^\infty(\langle z, F^* \upharpoonright \mathcal{A}_m \rangle) < j\}$ and note that $S_m^* \supseteq S_{m+1}^*$ and they are all compact. Indeed,

$$S_m^* = \bigcup_{\mathcal{B} \subseteq \mathcal{A}_m, v^\infty(\mathcal{B}) < j} \left(\bigcap_{a \in \mathcal{A}_m \setminus \mathcal{B}} [F^*(a)] \right)$$

and each $[F^*(a)]$ is compact.

Claim 2. *If $z \notin W$ and $i \geq M$ then $\langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle \subseteq \langle z, F_{m_i} \rangle$.*

Proof. Let $a_\ell \notin \langle z, F_{m_i} \rangle$ and assume that $\ell \leq m_i$ since otherwise it is immediate that $a_\ell \notin \langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle$. There are two cases to consider. First assume that $\ell \leq M$. Then $z \in [F_{m_i}(a_\ell)]$ but since $z \notin W$ and $i \geq M$ it follows that $z \notin [F_{m_i}(a_\ell) \setminus F_{M,\ell}]$. Hence $z \in [F_{M,\ell}] = [F^*(a_\ell)]$. In other words, $a_\ell \notin \langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle$. If $\ell > M$ a similar argument works. \square

It follows from the claim that if $z \notin W$ and $i \geq M$ then $v^\infty(\langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle) \leq v^\infty(\langle z, F_{m_i} \rangle)$. Hence, if $i \geq M$ and $z \in S_{m_i} \setminus W$ then $v^\infty(\langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle) < j$. In other words, $S_{m_i} \setminus W \subseteq S_{m_i}^* \setminus W$. Therefore, $\square_{\Theta(j-1)-\lambda(W)}(S_{m_i}^* \setminus W) \geq \square_{\Theta(j-1)-\lambda(W)}(S_{m_i} \setminus W) \geq \Gamma(j)$. Since each of the $S_{m_i}^* \setminus W$ are compact it follows that

$$\square_{\Theta(j-1)-\lambda(W)} \left(\bigcap_{i=M}^\infty S_{m_i}^* \setminus W \right) \geq \Gamma(j).$$

Let $S^* = \bigcap_{i=M}^\infty S_i^*$. Since $\lambda(Y) < \Theta(j-1) - \lambda(W)$ it follows that $\lambda_\infty^*(S^* \setminus (W \cup Y)) \geq \Gamma(j)$.

Since $\lambda_\infty^*(S \setminus Y) < \Gamma(j)$ it is possible to select $z \in S^* \setminus (S \cup W \cup Y)$. Then $v^\infty(\langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle) < j$ for each $i \geq M$ and so the induction hypothesis guarantees that

$$v^\infty \left(\bigcup_{i=M}^\infty \langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle \right) < j.$$

But

$$\bigcup_{i=M}^\infty \langle z, F^* \upharpoonright \mathcal{A}_{m_i} \rangle = \langle z, \bigcup_{i=M}^\infty F^* \upharpoonright \mathcal{A}_{m_i} \rangle = \langle z, F^* \rangle$$

and so $v^\infty(\langle z, F^* \rangle) < j$. Since $F^*(a) \subseteq F(a)$ for each a it is immediate that $\langle z, F^* \rangle \supseteq \langle z, F \rangle$ and so $v^\infty(\langle z, F \rangle) < j$. This contradicts that $z \notin S$. \square

Corollary 5.1. *For any $j \in \mathbb{N}$ and $\mu > 0$ there is a finite set $\mathcal{X} \subseteq \mathbb{C}_\mu^1$ such that $v(\mathcal{X}) \geq j$.*

Proof. Combine Corollaries 4.1 and 4.4 to conclude that $v^\infty(\mathbb{C}_\mu^1) = \infty$. Then use Lemma 5.2 to find a finite $\mathcal{X} \subseteq \mathbb{C}_\mu^1$ such that $v^\infty(\mathcal{X}) \geq j$. Finally apply Lemma 5.1. \square

6. The forcing partial order

Using Corollary 5.1 let $\mathfrak{C}(n)$ be a finite subset of $\mathbb{C}_{2^{-n}}^1$ such that $v(\mathfrak{C}(n)) \geq n$ for each $n \in \mathbb{N}$. Recalling the notation concerning trees in Section 2, let \mathbb{P} consist of all trees T such that:

$$(\forall t \in T)(\forall i < |t|) t(i) \in \mathfrak{C}(i), \tag{6.1}$$

$$(\forall k \in \omega)(\forall t \in T)(\exists s \in T) t \subseteq s \text{ and } v(\mathcal{S}_T(s)) > k \tag{6.2}$$

and let this be ordered under inclusion. The methods of [6] can be used to establish that \mathbb{P} is an ω^ω -bounding proper partial order.

Lemma 6.1. *Let $V \subseteq W$ be models of set theory and suppose that $G \subseteq \mathbb{P} \cap V$ is generic over W . Then $W[G] \models \lambda(W \cap [0, 1]) = 0$.*

Proof. If $G \subseteq \mathbb{P} \cap V$ is generic over W then let $B_G \in \prod_{n=0}^\infty \mathfrak{C}(n)$ be the generic branch determined by G . Note that $\lambda([B_G(j)]) < 2^{-j}$ for each j and so $\lambda(\bigcup_{j=0}^\infty [B_G(j)]) < \infty$. Also, for every $T \in \mathbb{P}$ there is some $t \in T$ such that $V \models v(\mathcal{S}_T(t)) \geq 1$. Since $v(\mathcal{S}_T(t)) \geq 1$ is equivalent to $\bigcup \mathcal{S}_T(t) = \mathbb{C}$, and this is absolute, it follows from genericity that $\mathbb{C} \cap W \subseteq \bigcup_{j=m}^\infty [B_G(j)]$ for every m . \square

Definition 6.1. Let \mathbb{P}^κ be the countable support product of κ copies of \mathbb{P} .

Corollary 6.1. *If $\kappa \geq \aleph_1$ and $G \subseteq \mathbb{P}^\kappa$ is a filter generic over V then in $V[G]$ every set of reals of size less than κ has Lebesgue measure 0.*

In light of Corollary 6.1 the goal now is to establish that if \mathbb{P}^κ is the countable support product of κ copies of \mathbb{P} and $G \subseteq \mathbb{P}^\kappa$ is a filter generic over V then $\lambda_\infty^r(\mathbb{C} \cap V) = 1$ in $V[G]$.

Lemma 6.2. *Suppose that $p \Vdash_{\mathbb{P}^\kappa} \text{“} C \in \mathbb{C}_\delta^\infty \text{”}$ and $\delta < 1$. Then there are $A \subseteq \mathbb{C}$ and $B \subseteq \mathbb{C}$ such that $\lambda_\infty^r(A) + \lambda(B) < 1$ such that $p \not\Vdash_{\mathbb{P}^\kappa} \text{“} z \in [C] \text{”}$ for each $z \in \mathbb{C} \setminus (A \cup B)$.*

Proof. Let $1 > \delta_1 > \delta$ and choose a monotonically increasing function $\delta_2 : \mathbb{N} \rightarrow \mathbb{R}$ such that $\delta_1 + \lim_{n \rightarrow \infty} \delta_2(n) < 1$. For later use, define $k(i) = \prod_{j=0}^i |\mathfrak{C}(j)|$. The proof is

based on a standard fusion argument. Induction will be used to construct $p_i, S_i, \varepsilon_i, a_i, b_i$ and L_i satisfying the following conditions:

- (1) $p_i \in \mathbb{P}^\kappa$ and $p_0 \leq p_i$;
- (2) $p_{i+1} \leq p_i$;
- (3) $S_i : \kappa \rightarrow \mathbb{N}$ is 0 at all but finitely many ordinals—these will be denoted by $D(S_i)$;
- (4) $S_i(\sigma) = S_{i+1}(\sigma)$ for all but one $\sigma \in \kappa$;
- (5) $S_{i+1}(\sigma) \leq S_i(\sigma) + 1$ for all $\sigma \in \kappa$;
- (6) $L_i \in \mathbb{N}$ and $L_i < L_{i+1}$;
- (7) $p_{i+1}(\sigma)[L_i] = p_i(\sigma)[L_i]$ for each $\sigma \in D(S_i)$;
- (8) for each $\tau \in \prod_{\sigma \in D(S_i)} p_i(\sigma)[L_i]$ there is some finite $C_{i,\tau}$ such that

$$p_i \langle \tau \rangle \Vdash_{\mathbb{P}^\kappa} \text{“} \check{C}_{i,\tau} \subseteq C \text{ and } C \setminus \check{C}_{i,\tau} \in \mathbb{C}_{\varepsilon_i}^\infty \text{”}$$

where $p_i \langle \tau \rangle$ is defined by $p_i \langle \tau \rangle (\alpha) = p_i(\alpha) \langle \tau(\alpha) \rangle$;

- (9) $a_i \in \mathbb{C}_{\alpha_i}$ for some $\alpha_i < \delta_1$ and $a_i \subseteq a_{i+1}$;
- (10) $\lambda([b_i]) < \delta_2(i)$ and $b_i \subseteq b_{i+1}$;
- (11) for each $z \in \mathbb{C} \setminus [a_i \cup b_i]$ there are finite trees $T_{\sigma,z,i}$ for each $\sigma \in D(S_i)$ such that
 - (a) $T_{\sigma,z,i} \subseteq p_i(\sigma)[\leq L_i]$,
 - (b) there is a maximal antichain of $s \in T_{\sigma,z,i}[\leq L_i]$ such that $v(\mathcal{S}_{T_{\sigma,z,i}}(s)) \geq S_i(\sigma)$,
 - (c) $z \notin [C_{i,\tau}]$ for any $\tau \in \prod_{\sigma \in D(S_i)} T_{\sigma,z,i}[L_i]$,
 - (d) $T_{\sigma,z,i}[L_i] = T_{\sigma,z,i+1}[L_i]$,
- (12) $2\varepsilon_i k(L_i)^{2i} < \delta_1 - \alpha_i$;
- (13) $\lim_{n \rightarrow \infty} S_n(\sigma) = \infty$ for each element σ of the domain of some p_i .

Assuming this can be done, let $A = \bigcup_{i=0}^\infty [a_i]$ and $B = \bigcup_{i=0}^\infty [b_i]$. Then if $z \in \mathbb{C} \setminus [A \cup B]$ let

$$q_z(\sigma) = \bigcup_{j=0}^\infty T_{\sigma,z,j}$$

and note that it follows from Condition 11b that $q_z \in \mathbb{P}^\kappa$ and $q_z \leq p$. From Conditions 8 and 11c it follows that $q_z \Vdash_{\mathbb{P}^\kappa} \text{“} z \notin [C] \text{”}$.

To see that the induction can be carried out, let i be given and suppose that $p_i, S_i, \varepsilon_i, a_i, b_i$ and L_i satisfying the induction hypothesis have been chosen. Choose $\bar{\sigma}$ according to some scheme which will satisfy Condition 13 and define

$$S_{i+1}(\sigma) = \begin{cases} S_i(\sigma) & \text{if } \sigma \neq \bar{\sigma}, \\ S_i(\sigma) + 1 & \text{if } \sigma = \bar{\sigma}. \end{cases}$$

Let $\bar{p}_0 \leq p_i$ be such that $\bar{p}_0 = p_i$ if $S_i(\bar{\sigma}) > 0$ and, otherwise, $\bar{p}_0(\sigma) = p_i(\sigma)$ for $\sigma \neq \bar{\sigma}$ and $|\bar{p}_0(\bar{\sigma})[L_i]| = 1$. Let $m \geq S_{i+1}(\bar{\sigma})$ be such that

$$\frac{\Gamma(m)}{2} > \varepsilon_i k(L_i)^i, \tag{6.3}$$

$$\Gamma(m)k(L_i) < \delta_1 - \alpha_i, \tag{6.4}$$

$$\Theta(m - 1) < \frac{\delta_2(i + 1) - \delta_2(i)}{k(L_i)} \tag{6.5}$$

and for each $t \in \bar{p}_0(\bar{\sigma})[L_i]$ there is some $t^* \in \bar{p}_0(\bar{\sigma})$ such that $t \subseteq t^*$ and such that

$$v(\mathcal{S}_{\bar{p}_0(\bar{\sigma})}(t^*)) \geq m + 1. \tag{6.6}$$

Let $L_{i+1} > L_i$ be so large that $|t^*| < L_{i+1}$ for each $t \in \bar{p}_0(\bar{\sigma})[L_i]$. Then let $\bar{p}_1 \leq \bar{p}_0$ be such that

- $\{t^* : t \in \bar{p}_0(\bar{\sigma})[L_i]\}$ is a maximal antichain in $\bar{p}_1(\bar{\sigma})$;
- if $t \in \bar{p}_1(\bar{\sigma})$ and $L_i \leq |t| \leq L_{i+1}$ then
 - $|\mathcal{S}_{\bar{p}_1(\bar{\sigma})}(t)| = 1$ if $t \neq t^*$ for any $t \in \bar{p}_0(\bar{\sigma})[L_i]$;
 - $\mathcal{S}_{\bar{p}_1(\bar{\sigma})}(t^*) = \mathcal{S}_{\bar{p}_0(\bar{\sigma})}(t^*)$ if $t \in \bar{p}_0(\bar{\sigma})[L_i]$;
- $\bar{p}_1(\sigma) = \bar{p}_0(\sigma)$ for $\sigma \neq \bar{\sigma}$.

Let $\bar{p}_2 \leq \bar{p}_1$ be such that $\bar{p}_2(\sigma) = \bar{p}_1(\sigma)$ for $\sigma \notin D(S_i) \setminus \{\bar{\sigma}\}$ and if $\sigma \in D(S_i) \setminus \{\bar{\sigma}\}$ choose $\bar{p}_2(\sigma)$ such that $\bar{p}_2(\sigma) \upharpoonright L_i = \bar{p}_1(\sigma) \upharpoonright L_i$ and such that if $t \in \bar{p}_2(\sigma)$ and $L_i \leq |t| \leq L_{i+1}$ then $|\mathcal{S}_{\bar{p}_2(\sigma)}(t)| = 1$. Let $\varepsilon_{i+1} > 0$ be so small that $2\varepsilon_{i+1}k(L_{i+1})^{2(i+1)} < \delta_1 - \alpha_i - \varepsilon_i k(L_i)$.

Choose $p_{i+1} \leq \bar{p}_2$ such that $p_{i+1}(\sigma)[L_{i+1}] = \bar{p}_2(\sigma)[L_{i+1}]$ for each $\sigma \in D(S_{i+1})$ —this implies that Condition 7 holds—and for each $\tau \in \prod_{\sigma \in D(S_{i+1})} p_{i+1}(\sigma)[L_{i+1}]$ there is some finite $C_{i+1,\tau}$ such that

$$p_{i+1} \langle \tau \rangle \Vdash_{\mathbb{P}^*} \check{C}_{i+1,\tau} \subseteq C \text{ and } C \setminus \check{C}_{i+1,\tau} \in \mathbb{C}_{\varepsilon_{i+1}}^\infty.$$

For each $t \in p_i(\bar{\sigma})[L_i]$ and $x \in \mathcal{S}_{p_i(\bar{\sigma})}(t^*)$ and $\tau \in \prod_{\sigma \in D(S_i) \setminus \{\bar{\sigma}\}} p_i(\sigma)[L_i]$ let $\rho(t, x, \tau)$ be the unique element of $\prod_{\sigma \in D(S_{i+1})} p_{i+1}(\sigma)[L_{i+1}]$ such that $\rho(t, x, \tau)(\sigma) \supseteq \tau(\sigma)$ for each $\sigma \in D(S_i) \setminus \{\bar{\sigma}\}$ and $\rho(t, x, \tau)(\bar{\sigma}) \supseteq t^*$ and $\rho(t, x, \tau)(\bar{\sigma})(|t^*|) = x$. Let $\rho^*(t, \tau)$ be the unique element of $\prod_{\sigma \in D(S_i)} p_i(\sigma)[L_i]$ such that $\rho^*(t, \tau) = \tau$ if $S_i(\bar{\sigma}) = 0$ and, if $S_i(\bar{\sigma}) > 0$ then $\rho^*(t, \tau)(\sigma) \supseteq \tau(\sigma)$ for each $\sigma \in D(S_i) \setminus \{\bar{\sigma}\}$ and $\rho^*(t, \tau)(\bar{\sigma}) = t$.

Note that $C_{i+1,\rho(t,x,\tau)} \setminus C_{i,\rho^*(t,\tau)} \in \mathbb{C}_{\varepsilon_i}$ by Condition 8. Define $F_t : \mathcal{S}_{p_{i+1}(\bar{\sigma})}(t^*) \rightarrow \mathbb{C}_{\varepsilon_i k(L_i)^i}$ by

$$F_t(x) = \bigcup_{\tau \in \prod_{\sigma \in D(S_i) \setminus \{\bar{\sigma}\}} p_i(\sigma)[L_i]} C_{i+1,\rho(t,x,\tau)} \setminus C_{i,\rho^*(t,\tau)}$$

and observe that $F_t(x) \in \mathbb{C}_{\Gamma(m)/2}$ by 6.3 since

$$\left| \prod_{\sigma \in D(S_i)} p_i(\sigma)[L_i] \right| \leq k(L_i)^{|D(S_i)|} \leq k(L_i)^i.$$

By 6.6 it follows that

$$\square_{\Theta(m-1)}(\{z \in \mathbb{C} : v(\langle z, F_t \rangle) < m\}) < \Gamma(m)$$

for each $t \in p_i(\bar{\sigma})[L_i]$. Let b'_{i+1} be such that $\lambda(b'_{i+1}) < \Theta(m - 1)$ and such that there is $a'_{i+1} \in \mathbb{C}_{\Gamma(m)}$ such that

$$[a'_{i+1}] \supseteq \{z \in \mathbb{C} : v(\langle z, F_t \rangle) < m\} \setminus [b'_{i+1}]. \tag{6.7}$$

It follows from 6.5 and Induction Hypothesis 10 that if b_{i+1} is defined to be $b_i \cup \bigcup_{t \in p_i(\bar{\sigma})[L_i]} b'_{i+1}$ then Condition 10 is satisfied by b_{i+1} . Similarly, if a_{i+1} is defined to be $a_i \cup \bigcup_{t \in p_i(\bar{\sigma})[L_i]} a'_{i+1}$ then $a_{i+1} \in \mathbb{C}_{\alpha_{i+1}}$ where $\alpha_{i+1} = \alpha_i + \Gamma(m)k(L_i) < \delta_1$ by 6.4 and Induction Hypothesis 9.

In order to verify that Condition 11 holds let $z \in \mathbb{C} \setminus [a_{i+1} \cup b_{i+1}]$. Let $T_{\sigma,z,i+1} = T_{\sigma,z,i}$ if $\sigma \neq \bar{\sigma}$ and let $T_{\bar{\sigma},z,i+1}$ be the set of all $s \in p_{i+1}(\bar{\sigma})[\leq L_{i+1}]$ such that $s \upharpoonright L_i \in T_{\sigma,z,i}$ and, if $t^* \subseteq s$ then $s(|t^*|) \in \langle z, F_t \rangle$. In order to show that Condition 11b holds it suffices to show that

$$v(\mathcal{S}_{T_{\bar{\sigma},z,i+1}}(t^*)) \geq m > S_{i+1}(\bar{\sigma})$$

for any $t \in p_i(\bar{\sigma})[L_i]$. For any given t this follows from the fact that $z \notin [a'_{i+1} \cup b'_{i+1}]$ and 6.7.

In order to show that Condition 11c holds let $\tau \in \prod_{\sigma \in D(S_{i+1})} T_{\sigma,z,i}[L_{i+1}]$. Let $\bar{\tau} = \tau \upharpoonright (D(S_i) \setminus \{\bar{\sigma}\})$. Let $t \in p_i(\bar{\sigma})[L_i]$ be such that $t \subseteq \tau(\bar{\sigma})$ and note that the definition of $T_{\bar{\sigma},z,i}$ guarantees that $\tau(\bar{\sigma})(|t^*|) \in \langle z, F_t \rangle$. In other words, $z \notin [F_t(\tau(\bar{\sigma})(|t^*|))] \supseteq [C_{i+1,\rho(t,\tau(\bar{\sigma})(|t^*|),\bar{\tau})} \setminus C_{i,\rho^*(t,\bar{\tau})}]$. Moreover, since $\rho^*(t, \bar{\tau}) \in \prod_{\sigma \in D(S_i)} T_{\sigma,z,i}$ it follows that $z \notin [C_{i,\rho^*(t,\bar{\tau})}]$ by the Induction Hypothesis 11c. Hence $z \notin [C_{i+1,\tau}]$. \square

Corollary 6.2. *If $G \subseteq \mathbb{P}$ is generic over V then $V[G] \models \lambda'_\infty(V \cap \mathbb{C}) = 1$.*

Proof. Suppose not and that

$$p \Vdash_{\mathbb{P}^\kappa} \text{“} C \in \mathbb{C}_\delta^\infty \text{ and } V \cap \mathbb{C} \subseteq [C]\text{”}$$

and $\delta < 1$. Using Lemma 6.2 find $A \subseteq \mathbb{C}$ and $B \subset \mathbb{C}$ such that $\lambda'_\infty(A) + \lambda(B) < 1$ and $q \leq p$ such that $q \not\Vdash_{\mathbb{P}^\kappa} \text{“} z \in [C]\text{”}$ for each $z \in \mathbb{C} \setminus (A \cup B)$. Choose any $z \in \mathbb{C} \setminus (A \cup B)$ and $q' \leq q$ such that $q' \Vdash_{\mathbb{P}^\kappa} \text{“} z \notin [C]\text{”}$. \square

7. Remarks and open questions

One might expect that the methods developed here could be used to prove that for any two reals s and t such that $0 < t < s < 1$ it is consistent that sets of size \aleph_1 are null with respect to s -dimensional Hausdorff measure but that this is not so for

t -dimensional Hausdorff measure. While this is true of most of the argument there are some slippery spots. For example, the use of Y_F in the proof of Lemma 3.5 assumes the σ -finiteness of Lebesgue measure. Lemma 3.2 might also pose some challenges to generalization. Let \mathfrak{n}_s denote the least cardinal of a set which is not null with respect to s -dimensional Hausdorff measure. Hence the following questions remain open:

Question 7.1. Is it consistent that $0 < t < s < 1$ and $\mathfrak{n}_t < \mathfrak{n}_s$?

Question 7.2. Is it consistent that $0 < u < v < w < 1$ and $\mathfrak{n}_u < \mathfrak{n}_v < \mathfrak{n}_w$?

Question 7.3. How big can the cardinality of $\{\mathfrak{n}_s\}_{s \in (0,1)}$ be?

However, the main open problem in this area still remains Question 1.1. It would be interesting to know what the answer to this question is in the model described in Section 6.

References

- [1] T. Bartoszyński, H. Judah, *On the structure of the real line*, Set Theory, A K Peters Ltd., Wellesley, MA, 1995.
- [2] D. Fremlin, Problem sheet. <http://www.essex.ac.uk/maths/staff/fremlin/problems.htm>, 2002.
- [3] M. Goldstern, S. Shelah, Many simple cardinal invariants, *Arch. Math. Logic* 32 (1993) 203–221.
- [4] A.S. Kechris, *Classical descriptive set theory*, in: Graduate Texts in Mathematics, Vol. 156, Springer, New York, 1995.
- [5] P. Mattila, *Geometry of sets and measures in Euclidean spaces*, in: Cambridge Studies in Advanced Mathematics, Vol. 44, Cambridge University Press, Cambridge, 1995.
- [6] A. Roslanowski, S. Shelah, Norms on possibilities. I. Forcing with trees and creatures, *Mem. Amer. Math. Soc.* 141 (1999) xii–167.
- [7] T. Salisbury, J. Steprāns, Hausdorff measure and Lebesgue measure, *Real Anal. Exchange* 22 (1996/1997) 265–278.
- [8] J. Steprāns, Cardinal invariants associated with Hausdorff capacities, in: *Proceeding of the BEST Conferences 1–3, Contemporary Mathematics*, Vol. 192, American Mathematical Society, Providence, RI, 1995, pp. 157–184.
- [9] J. Steprāns, Unions of rectifiable curves and the dimension of banach spaces, *J. Symbolic Logic* 64 (2) (1999) 701–726.