

The Journal of Symbolic Logic

<http://journals.cambridge.org/JSL>

Additional services for *The Journal of Symbolic Logic*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Identities on cardinals less than ω

M. Gilchrist and S. Shelah

The Journal of Symbolic Logic / Volume 61 / Issue 03 / September 1996, pp 780 - 787
DOI: 10.2307/2275784, Published online: 12 March 2014

Link to this article: http://journals.cambridge.org/abstract_S0022481200017114

How to cite this article:

M. Gilchrist and S. Shelah (1996). Identities on cardinals less than ω . The Journal of Symbolic Logic, 61, pp 780-787 doi:10.2307/2275784

Request Permissions : [Click here](#)

IDENTITIES ON CARDINALS LESS THAN \aleph_ω

M. GILCHRIST AND S. SHELAH

§1. Introduction. Let κ be an uncountable cardinal and the edges of a complete graph with κ vertices be colored with \aleph_0 colors. For $\kappa > 2^{\aleph_0}$ the Erdős-Rado theorem implies that there is an infinite monochromatic subgraph. However, if $\kappa \leq 2^{\aleph_0}$, then it may be impossible to find a monochromatic triangle. This paper is concerned with the latter situation. We consider the types of colorings of finite subgraphs that must occur when the edges of the complete graph on $\kappa \leq 2^{\aleph_0}$ vertices are colored with \aleph_0 colors. In particular, we are concerned with the case $\aleph_1 \leq \kappa \leq \aleph_\omega$.

The study of these color patterns (known as identities) has a history that involves the existence of compactness theorems for two cardinal models [2]. When the graph being colored has size \aleph_1 , the identities that must occur have been classified by Shelah [4]. If the graph has size greater than or equal to \aleph_ω the identities have also been classified in [3]. The number of colors is fixed at \aleph_0 as it is the natural place to start and the results here can be generalized to situations where more colors are used.

There is one difference that we now make explicit. When countably many colors are used we can define the following coloring of the complete graph on 2^{\aleph_0} vertices. First consider the branches in the complete binary tree of height ω to be vertices of a complete graph. The edge $\{\eta, \nu\} \in [{}^\omega 2]^2$ is given the color $\eta \cap \nu$. (In the notation that follows this coloring induces the identities in IDM and no others.) When $\kappa > \aleph_0$ colors are used we may not be able to define an analogous coloring of the complete graph on 2^κ vertices since it is known that there exist models of set theory in which no tree has \aleph_1 nodes and 2^{\aleph_1} branches [1].

The first section of this paper deals with the definitions that are necessary to describe identities. In the remainder of the paper we use forcing to construct a model V of set theory in which the set of identities which must occur when $\kappa = \aleph_{m+1}$, strictly includes the set of identities which must occur when $\kappa = \aleph_m$. This is done by showing that a certain identity is omitted when $\kappa = \aleph_m$ but is induced when $\kappa = \aleph_{m+1}$.

§2. Preliminaries. An ω -coloring is function $f : [B]^2 \rightarrow \omega$ where B is a set of ordinals ordered in the usual way. The set B is the *field* of f and is denoted $\text{fld}(f)$.

Received May 10, 1993; revised April 10, 1995.

S. Shelah partially supported by a research grant from the basic research fund of the Israel Academy of Science; Pul. Nu. 491.

DEFINITION 1. Let f, g be ω -colorings. We say that f realizes the coloring g if there is a one-one map $k: \text{fld}(g) \rightarrow \text{fld}(f)$ such that for all $\{x, y\}, \{u, v\} \in \text{dom}(g)$

$$f(\{k(x), k(y)\}) \neq f(\{k(u), k(v)\}) \Rightarrow g(\{x, y\}) \neq g(\{u, v\}).$$

We write $f \simeq g$ if f realizes g and g realizes f . It should be clear that \simeq induces an equivalence relation on the class of ω -colorings. We call the equivalence classes *identities*. The collection of all identities is denoted ID.

DEFINITION 2. Let f, g be ω -colorings. We say that f *V-realizes* the coloring g if there is an order-preserving map $k: \text{fld}(g) \rightarrow \text{fld}(f)$ such that for all $\{x, y\}, \{u, v\} \in \text{dom}(g)$

$$f(\{k(x), k(y)\}) \neq f(\{k(u), k(v)\}) \Rightarrow g(\{x, y\}) \neq g(\{u, v\}).$$

We write $f \simeq_V g$ if f V-realizes g and g V-realizes f . Note that \simeq_V induces an equivalence relation on the class of ω -colorings. We call the equivalence classes *V-identities*. The collection of all V-identities is denoted ID_V.

For both types of realization we will call the map $k: \text{fld}(g) \rightarrow \text{fld}(f)$ an *embedding*. In the above definition the V refers to vertices. In the situations that follow, B will be a cardinal less than or equal to \aleph_ω ordered in the usual way as a set of ordinals. In the following we will speak of ω -colorings realizing (rather than V-realizing) other ω -colorings whenever the context makes the type of realization clear. If f, g, h, l are ω -colorings, with $f \simeq g$ and $h \simeq l$, then f realizes h if and only if g realizes l . Thus without risk of confusion we may speak of identities realizing colorings and of identities realizing other identities. The same is true of V-identities. If I and J are identities we call J a *subidentity* of I if I realizes J . The notion of sub-V-identity is similarly defined. We say that an identity I is of *size* r if $|\text{fld}(f)| = r$ for some (all) $f \in I$. In the following we will consider only identities of finite size.

An identity can be regarded as a finite structure $\langle A, E \rangle$, where E is an equivalence relation on $[A]^2$, see [4]. The correspondence is given by: $A = \text{fld}(f)$, and $\{x, y\} \simeq_E \{u, v\}$ if and only if $f(\{x, y\}) = f(\{u, v\})$.

A V-identity can be regarded as a finite structure $\langle A, E, <_A \rangle$, where E is an equivalence relation on $[A]^2$ and $<_A$ is a linear ordering of A . The correspondence is given by: $A = \text{fld}(f)$, $\{x, y\} \simeq_E \{u, v\}$ if and only if $f(\{x, y\}) = f(\{u, v\})$, and $x < y$ if and only if $x <_A y$.

We remark that the notion of subidentity does not correspond to that of substructure. To simplify the language in what follows we find it convenient to abuse terminology by referring to ω -colorings and structures $\langle A, E \rangle$ as identities rather than as representatives of identities. Similarly for V-identities. In each case the intended meaning should be clear.

DEFINITION 3. Let $f: [B]^2 \rightarrow \omega$ be an ω -coloring and $I = \langle A, E, <_A \rangle$ be a structure corresponding to a V-identity. Let $k: A \rightarrow B$ be an order-preserving map such that for all $\{x, y\}, \{v, w\} \in [A]^2$,

$$\{k(x), k(y)\} \not\simeq_E \{k(v), k(w)\} \Rightarrow f(\{x, y\}) \neq f(\{v, w\}).$$

Then k is called an *embedding* of I into f . A similar definition is given when I is an identity.

DEFINITION 4. Let D be a set of ordinals and $f: [B]^2 \rightarrow \omega$ be an ω -coloring.

- (1) $\mathcal{S}(f)$ ($\mathcal{S}_V(f)$) is the collection of (V-)identities realized by f .
- (2) $\mathcal{S}(D) = \bigcap \{ \mathcal{S}(g) \mid g: [D]^2 \rightarrow \omega \}$, $\mathcal{S}_V(D) = \bigcap \{ \mathcal{S}_V(g) \mid g: [D]^2 \rightarrow \omega \}$.

If $J = \langle B, F \rangle$ is an identity and $A \subset B$ we define the *restriction* of J to A to be the identity $I = \langle A, F \cap ([A]^2 \times [A]^2) \rangle$. This will be written: $I = J \upharpoonright A$. Similarly for V-identities.

DEFINITION 5. Let $n < \omega$, $I = \langle A, E \rangle$, $J = \langle B, F \rangle$ be identities, $\bar{a} = \langle a_1, \dots, a_n \rangle \in {}^n A$ be a sequence of distinct elements from A , and $\bar{b} = \langle b_1, \dots, b_n \rangle \in {}^n B$ be a sequence of distinct elements from B . J is obtained from I by *duplication* of \bar{a} to \bar{b} if

- (1) $B = A \sqcup \bar{b}$
- (2) $I = J \upharpoonright A$
- (3) the mapping which is the identity on $A \setminus \bar{a}$ and which maps \bar{a} to \bar{b} is an embedding of I into J as structures
- (4) F is the least equivalence relation on $[B]^2$ consistent with (i)–(iii).

When $n = 1$ we say that J is a *one-point* duplication of I .

DEFINITION 6. Let $I = \langle A, E, <_A \rangle$ and $J = \langle B, F, <_B \rangle$ be V-identities. J is obtained from I by *end-duplication* if there exist final segments \bar{a} , \bar{b} of $\langle A, <_A \rangle$, $\langle B, <_B \rangle$ respectively such that the structure $\langle B, F \rangle$ is obtained from the structure $\langle A, E \rangle$ by duplicating \bar{a} to \bar{b} .

Let IDE_V denote the minimal collection of V-identities which is closed under end-duplication, the taking of subidentities, and which contains the trivial V-identity of size one. Let $\text{IDE} = \{ \langle A, E, < \rangle \in \text{IDE}_V \}$.

Let IDM denote the least class of identities which is closed under duplication, the taking of subidentities and which contains the identity of size one. We quote the following results of Shelah:

THEOREM 1 ([4]). $\mathcal{S}(\aleph_1) \supset \text{IDE}$.

THEOREM 2 ([5]). *There exists $f: [\aleph_1]^2 \rightarrow \aleph_0$ such that $\mathcal{S}(f) \subseteq \text{IDE}$.*

COROLLARY 1. $\mathcal{S}(\aleph_1) = \text{IDE}$.

THEOREM 3 ([3]). *If $\kappa \geq \aleph_\omega$ then $\mathcal{S}(\kappa) \supseteq \text{IDM}$.*

§3. The partial order. We are going to define a partial order that will be used as a forcing notion to allow us to omit an identity from $\mathcal{S}(\aleph_m)$. The construction of the partial order uses historical forcing [6]. In this method conditions are allowed into the partial order if they can be constructed from the amalgamation of simpler conditions satisfying certain properties. As a preliminary we construct a model which has many definable subsets.

In the following, first-order languages will be denoted by L , possibly with subscripts. The variables of each first-order language are the same and have a canonical

well-ordering of order type ω . We denote by $F_n(L)$ the set of all formulas of L in which at most the first n variables occur free. We let $F(L) = \bigcup\{F_n(L) : n < \omega\}$. Let L_0 denote the language $\{\langle\}\}$. Let $<_m$ denote the natural ordering of \aleph_m . We regard $\langle\aleph_m; <_m\rangle$ as an L_0 -structure. From now on fix m to be an integer ≥ 1 .

LEMMA 1. *There exists a countable relational language $L \supset L_0$ and an expansion \mathcal{M} of $\langle\aleph_m; <_m\rangle$ to L such that for each $k < \omega$ and $\phi \in F_{k+1}(L)$ there exists $\psi \in F_{k+2}(L)$ such that for every k -tuple $\bar{a} \in \aleph_m$, the relation defined by $\psi(x, y, \bar{a})$ in \mathcal{M} well-orders $\{b \in \mathcal{M} : \mathcal{M} \models \phi(b, \bar{a})\}$ with order type $|\{b \in \mathcal{M} : \mathcal{M} \models \phi(b, \bar{a})\}|$.*

PROOF. We define a sequence of models and languages $\langle\langle\mathcal{M}_i, L_i\rangle : i < \omega\rangle$ with the property:

For each $k < \omega$, $\varphi \in F_{k+1}(L_n)$ and k -tuple $\bar{a} \in \aleph_m^k$ there exists $\psi \in F_{k+2}(L_{n+1})$ such that $\psi(x, y, \bar{a})$ well-orders $\{b : \mathcal{M} \models \varphi(b, \bar{a})\}$ in order type $|\{b : \mathcal{M} \models \varphi(b, \bar{a})\}|$.

Assume that $\langle\mathcal{M}_n, L_n\rangle$ has been defined. For every $k < \omega$ and $\varphi \in F_{k+1}(L_n)$ we add a $(k+2)$ -ary relation symbol W_φ to L_{n+1} and expand the structure \mathcal{M}_n to \mathcal{M}_{n+1} so that for all $\bar{a} \in \aleph_m^k$, in \mathcal{M}_{n+1} the formula $W_\varphi(x, y, \bar{a})$ well-orders $\{b : \mathcal{M}_n \models \varphi(b, \bar{a})\}$ in order type $|\{b : \mathcal{M}_n \models \varphi(b, \bar{a})\}|$. We let $L = \bigcup\{L_n : n < \omega\}$ and $\mathcal{M} = \bigcup\{\mathcal{M}_n : n < \omega\}$. \dashv

DEFINITION 7. Let $b \in \aleph_m$, $k, n < \omega$, and $\bar{a} \in \aleph_m^k$. We say $bR_n\bar{a}$ if there exists $\varphi \in F(L)$ such that $\mathcal{M} \models (\exists \leq \aleph_n x)\varphi(x, \bar{a})$ and $\mathcal{M} \models \varphi(b, \bar{a})$.

Let $\mathbb{R} = \{\langle u, c \rangle : u \in [\aleph_m]^{<\aleph_0}, c : [u]^2 \rightarrow \omega\}$.

DEFINITION 8. $p = \langle u, c \rangle \in \mathbb{R}$ is the *amalgam* of $p^0 = \langle u^0, c^0 \rangle$ and $p^1 = \langle u^1, c^1 \rangle \in \mathbb{R}$ if there exist $h < \omega$ and increasing sequences i_0^0, \dots, i_h^0 and i_0^1, \dots, i_h^1 in \aleph_m such that for all s, t with $(0 \leq s < t \leq h)$, and all $i, j, k, l < \aleph_m$:

- (1) $u^0 = \{i_0^0 \dots i_h^0\}$ and $u^1 = \{i_0^1 \dots i_h^1\}$
- (2) $c^0(\{i_s^0, i_t^0\}) = c^1(\{i_s^1, i_t^1\})$
- (3) $i_t^0 = i_t^1 \vee i_t^0 < i_t^1$
- (4) $i_t^0 \neq i_t^1$ implies $\neg i_t^1 R_0 u^0$
- (5) $u = u^0 \cup u^1$
- (6) $c \supset (c^0 \cup c^1)$
- (7) $\{\{i, j\} \notin [u^0]^2 \cup [u^1]^2\}$ implies $c(\{i, j\}) \notin \text{rng}(c^0) \cup \text{rng}(c^1)$
- (8) $c(\{i, j\}) = c(\{k, l\})$ implies $(\{i, j\} = \{k, l\} \vee \{i, j\}, \{k, l\} \in [u^0]^2 \cup [u^1]^2)$

DEFINITION 9. $q = \langle u^q, c^q \rangle \in \mathbb{R}$ is a *one-point extension* of $p = \langle u^p, c^p \rangle \in \mathbb{R}$ if $u^q = u^p \cup \{r\}$ for some $r > u^p$, $c^p \subset c^q$, and for all $i, j, k, l \in u^q$

- (1) $\{i, j\} \notin \text{dom}(c^p)$ implies $c^q(\{i, j\}) \notin \text{rng}(c^p)$
- (2) $c^q(\{i, j\}) = c^q(\{k, l\})$ implies $(\{i, j\}, \{k, l\} \in \text{dom}(p) \vee \{i, j\} = \{k, l\})$.

DEFINITION 10. We now define a sequence of subsets of \mathbb{R} . Let $\mathbb{P}_0 = \{\langle u, c \rangle \in \mathbb{R} : |u| = 1\}$. Given \mathbb{P}_n we let \mathbb{P}_{n+1} be the subset of \mathbb{R} which contains \mathbb{P}_n , all amalgam of pairs of elements from \mathbb{P}_n and all one-point extensions of elements of \mathbb{P}_n . Let $\mathbb{P} = \bigcup\{\mathbb{P}_n : n < \omega\}$. Given $p = \langle u^p, c^p \rangle$ and $q = \langle u^q, c^q \rangle$ we let $p \leq q$ if and only if $u^p \supseteq u^q$ and $c^p \supseteq c^q$.

It should be noted that the order of p^0 and p^1 in the definition of amalgamation is important because of the asymmetry in the properties of p^0 and p^1 required by

the definition. It is also worth observing that in terms of the notion of duplication, the amalgam of p^0 and $p^1 \in \mathbb{R}$ may be regarded as being obtained from p^0 by simultaneous duplication of all the elements in $u^0 \setminus u^1$. Of course, the amalgamation also requires that none of the elements of $u^0 \setminus u^1$ belong to the countable set definable over u^0 . The closure of \mathbb{P} under one-point extensions is necessary to show that our forcing produces a function whose domain is of size \aleph_m .

Let $p = \langle u, c \rangle \in \mathbb{P}$ and $I = \langle A, E \rangle$ be an identity. We say that p realizes I if the ω -coloring $c: [u]^2 \rightarrow \omega$ realizes I . The mapping $h: A \rightarrow u$ demonstrating the realization is called an *embedding* of I into p .

LEMMA 2. \mathbb{P} is c.c.c.

PROOF. Let $\langle p_\alpha : \alpha < \omega_1 \rangle$ be a sequence of conditions. By thinning we can suppose that there are $n, l < \omega$ and i_j^α ($\alpha < \omega_1, 0 \leq j \leq n$) such that for all $\alpha, \beta < \omega_1$ and all j, k with $0 \leq j < k \leq n$.

- (1) $u^{p_\alpha} = \{i_0^\alpha, \dots, i_n^\alpha\}$
- (2) $i_j^\alpha < i_k^\alpha$
- (3) $c^{p_\alpha}(\{i_j^\alpha, i_k^\alpha\}) = c^{p_\beta}(\{i_j^\beta, i_k^\beta\})$
- (4) $p_\alpha \in \mathbb{P}_l$.

Applying the Δ -system argument allows us to thin the sequence of conditions further so that

$$\forall t (0 \leq t \leq n \Rightarrow [\forall \alpha \forall \beta (i_t^\alpha = i_t^\beta) \vee (\forall \beta < \omega_1) (\forall \alpha < \beta) (i_t^\alpha < i_t^\beta)]).$$

Let $T = \{t \leq n : i_t^\alpha \neq i_t^\beta \text{ some } \alpha, \beta < \omega_1\}$. To prove that \mathbb{P} is c.c.c. it is sufficient to show that p_0 and p_α are compatible for some $\alpha < \omega_1$. For p_0 and p_α to have a common extension by definition we need only that $t \in T$ implies $\neg i_t^\alpha R_0(i_0^0 \dots i_n^0)$. Since the language is countable $|\{i \in \aleph_m : i R_0(i_0^0 \dots i_n^0)\}| = \aleph_0$. For each $t \in T$, i_t^α is strictly increasing in α . Hence $\neg i_t^\alpha R_0(i_0^0 \dots i_n^0)$ for all sufficiently large $\alpha < \omega_1$. Since T is finite, the condition above is satisfied for all sufficiently large α . This proves the lemma. \dashv

LEMMA 3. For each $\alpha < \aleph_m$

$$\{\langle u, c \rangle \in \mathbb{P} : \exists \beta (\beta \in u \wedge \alpha < \beta)\}$$

is dense in $(\mathbb{P}, <)$.

PROOF. Clear from the definition of one-point extension. \dashv

LEMMA 4. Let M be any model and G be \mathbb{P} generic. In $M[G]$ there is a function $f: [\aleph_m]^2 \rightarrow \omega$ such that every identity realized by f is a subidentity of an identity realized by some $p \in \mathbb{P}$.

PROOF. Standard forcing technique. \dashv

\S 4. Omitting I_m . Throughout this section let m be an integer ≥ 1 . We denote by I_m the identity $\langle A, E \rangle$ where $A = {}^{m+1}2$ and $\{\eta, \nu\} \simeq_E \{\alpha, \beta\}$ if and only if $\eta \cap \nu = \alpha \cap \beta$. We show that the model of ZFC produced by forcing with the partial order of the previous section contains a function $f: [\aleph_{m+1}]^2 \rightarrow \omega$ that

does not realize I_m . It should be noted that $I_m \in \text{IDM}$. Since Theorem 3 says that $\mathcal{F}(\aleph_\omega) \supseteq \text{IDM}$, the model produced by forcing clearly satisfies $\mathcal{F}(\aleph_{m+1}) \not\subseteq \mathcal{F}(\aleph_\omega)$.

DEFINITION 11. Let $I = \langle A, E \rangle$ be an identity and $p = \langle u, c \rangle \in \mathbb{R}$. Then $h: A \rightarrow u$ is an *embedding of I into p* if h is an embedding of I into the ω -coloring c .

DEFINITION 12. $\langle \eta_0, \dots, \eta_m, \eta_{m+1} \rangle$ is a *special sequence* if

- (1) $\eta_i \in {}^{m+1}2$
- (2) $|\eta_i \cap \eta_{i+1}| = i$ for all $i \leq m$.

LEMMA 5. Let $p \in \mathbb{R}$ and h be an embedding of I_m into p . There exists a special sequence $\langle \eta_0, \dots, \eta_m, \eta_{m+1} \rangle$ so that $h(\eta_i)R_0(h(\eta_0) \dots h(\eta_{m-1}))$ for $i \in \{m, m+1\}$.

PROOF. We define $\eta_k \in {}^{m+1}2$, $0 \leq k \leq m-1$, by induction on k such that

- (1) $|\eta_i \cap \eta_{i+1}| = i$ for all $i < m-1$
- (2) $h(\gamma)R_{m-(i+1)}(h(\eta_0), \dots, h(\eta_i))$ for all $i < m$ and all $\gamma \in {}^{m+1}2$ such that $|\gamma \cap \eta_i| = i$.

Let η_0 be the unique $v \in {}^{m+1}2$ such that $h(v) = \max(\text{rng}(h))$. Suppose that η_k has been suitably defined for all $k \leq j$ where $j < m-1$. Let C denote $\{v \in {}^{m+1}2 : |v \cap \eta_j| = j\}$. From the induction hypothesis there exists $D \subset \aleph_m$ such that $\text{rng}(h \upharpoonright C) \subset D$ and D definable in \mathcal{M} over $\{h(\eta_0), \dots, h(\eta_j)\}$. From the choice of \mathcal{M} there is a relation $<_D$ definable over $\{h(\eta_0), \dots, h(\eta_j)\}$ which well-orders D in the order type less than or equal to $\aleph_{m-(j+1)}$. Let η_{j+1} be the unique $v \in C$ such that $h(v)$ is the $<_D$ maximal element of $\text{rng}(h \upharpoonright C)$. Clearly, $|\eta_{j+1} \cap \eta_j| = j$. Consider $\gamma \in {}^{m+1}2$ such that $|\gamma \cap \eta_{j+1}| = j+1$. Clearly, $\gamma \in C$ and $\gamma \neq \eta_{j+1}$. Hence $h(\gamma) \in D$ and $h(\gamma) <_D h(\eta_{j+1})$. It follows that $h(\gamma)R_{m-(j+2)}(h(\eta_0), \dots, h(\eta_j), h(\eta_{j+1}))$. This completes the induction step and the definition of $\eta_0, \dots, \eta_{m-1}$. Letting η_m, η_{m+1} be the two elements of $\{v \in {}^{m+1}2 : |v \cap \eta_{m-1}| = m-1\}$ completes the proof. \dashv

LEMMA 6. Let $\langle \eta_0, \dots, \eta_{m+1} \rangle$ be a special sequence, $p, q \in \mathbb{R}$, p be a one-point extension of q , and h be an embedding of $J = I_m \upharpoonright \{\eta_0, \dots, \eta_{m+1}\}$ in p . Then h is an embedding of J in q .

PROOF. Let $p = \langle u, c \rangle$ and $q = \langle v, d \rangle$. Towards a contradiction suppose that $u \setminus v = \{h(\eta_i)\}$. If $i < m$, then $\{\eta_m, \eta_i\}, \{\eta_{m+1}, \eta_i\}$ are equivalent in I_m , but $c(\{h(\eta_m), h(\eta_i)\}) \neq c(\{h(\eta_{m+1}), h(\eta_i)\})$ since p is a one-point extension of q . This contradicts h being an embedding. If $i \in \{m, m+1\}$, the consideration of the pairs $\{\eta_0, \eta_m\}, \{\eta_0, \eta_{m+1}\}$ leads to a similar contradiction. \dashv

LEMMA 7. Let $p \in \mathbb{P}$. Then there does not exist an embedding h of I_m into p .

PROOF. Towards a contradiction suppose the theorem fails. From Lemma 5 there exist a special sequence $\langle \eta_0, \dots, \eta_{m+1} \rangle$, $p = \langle u, c \rangle \in \mathbb{P}$, and h embedding $J = I_m \upharpoonright \{\eta_0, \dots, \eta_{m+1}\}$ in p such that $h(\eta_i)R_0(h(\eta_0), \dots, h(\eta_{m-1}))$ for $i \in \{m, m+1\}$. Fixing $\langle \eta_0, \dots, \eta_{m+1} \rangle$ and h choose p to minimize $|u|$. From Lemma 6, p is not a one-point extension of $q \in \mathbb{P}$. Therefore there are $p^0 = \langle u^0, c^0 \rangle, p^1 = \langle u^1, c^1 \rangle \in \mathbb{P}$ such that p is the amalgam of p^0 and p^1 . Since neither p^0 nor p^1 can replace p , there exist $i, j \leq m+1$ and $a, b \in u$ such that $h(\eta_i) = a \in u^0 \setminus u^1, h(\eta_j) = b \in u^1 \setminus u^0$.

From the definition of amalgamation, $\{a, b\}$ is the only pair in $[u]^2$ which is assigned the color $c(\{a, b\})$ by p . The only pair in $[\{\eta_0, \dots, \eta_{m+1}\}]^2$ which is in an

equivalence class of size one in I_m is $\{\eta_m, \eta_{m+1}\}$. Without loss of generality $i = m$ and $j = m + 1$. Also, it is clear that $h(\eta_0), \dots, h(\eta_{m-1})$ are all in $u^0 \cap u^1$.

We have that $b \in u^1 \setminus u^0$ belongs to a countable set definable in \mathcal{M} over $u^0 \cap u^1$. This contradicts the definition of amalgamation and completes the proof of the lemma. \dashv

THEOREM 4. *Let M be any model and G be \mathbb{P} generic. Then $M[G]$ satisfies, $I_m \notin \mathcal{F}(\aleph_m)$.*

PROOF. The result follows from Lemma 4 and Lemma 7. \dashv

§5. Inducing I_m . To show that the identities induced at different cardinals can distinguish the cardinals themselves, as promised in the introduction, we will show that $I_m \in \mathcal{F}(\aleph_{m+1})$ for all m . This will be done by showing that if a given collection of identities is induced at \aleph_m we can extend the collection in a nontrivial way and be assured that this new collection is induced at \aleph_{m+1} .

DEFINITION 13. Let $\langle J_i : 1 \leq i \leq n \rangle$ be a finite sequence of identities. We define the *end-homogeneous amalgam* of the sequence as follows. Choose a sequence of ω -colorings $c_i: [G_i]^2 \rightarrow \omega$ such that $c_i \in J_i$, $G_i \cap G_j = \emptyset$ for $1 \leq i < j \leq n$, and $\text{rng}(c_i) \cap \text{rng}(c_j) = \emptyset$ for all $1 \leq i < j \leq n$. Let $G = \bigcup \{G_i : 1 \leq i \leq n\}$. Now choose a new ω -coloring $c: [G]^2 \rightarrow \omega$ such that for all $\{r, s\}, \{t, v\} \in [G]^2$ and all i , $1 \leq i \leq n$,

- (1) $c \supset c_i$
- (2) $c(\{r, s\}) \in \text{rng}(c_i)$ if and only if $\{r, s\} \in \text{dom}(c_i)$
- (3) if $\{r, s\}, \{t, v\}$ are not in $\bigcup \{\text{dom}(c_j) : 1 \leq j \leq n\}$, then

$$c(\{r, s\}) = c(\{t, v\}) \iff \min\{j : r \in G_j \vee s \in G_j\} = \min\{j : t \in G_j \vee v \in G_j\}.$$

The end-homogeneous amalgam of $\langle J_i : 1 \leq i \leq n \rangle$ is the identity realized by c .

Let \mathcal{F} be a collection of identities. Define the *closure* of \mathcal{F} (denoted $\text{cl}(\mathcal{F})$) to be the collection of identities produced by forming all end-homogeneous amalgam of all finite sequences of identities in \mathcal{F} .

THEOREM 5. *Let $0 < m < \omega$. If $\mathcal{F} \subseteq \mathcal{F}(\aleph_m)$ then $\text{cl}(\mathcal{F}) \subseteq \mathcal{F}(\aleph_{m+1})$.*

PROOF. Let $f: [\aleph_{m+1}]^2 \rightarrow \omega$ and $\langle J_i : 1 \leq i \leq n \rangle$ be a sequence of identities in \mathcal{F} . We will produce by recursion a sequence $\langle \langle A_k, B_k \rangle : 0 \leq k \leq n \rangle$ such that:

- (1) f induces J_i on the set A_i for $1 \leq i \leq n$
- (2) $B_i \supset B_{i+1}$ for $0 \leq i < n$
- (3) $|B_i| = \aleph_{m+1}$ for $0 \leq i \leq n$
- (4) $A_{i+1} \subset B_i \setminus B_{i+1}$
- (5) $f(\{a_1, b_1\}) = f(\{a_2, b_2\})$ whenever there exist i, j ($1 \leq i \leq j \leq n$) such that $\{a_1, a_2\} \subset A_i$ and $\{b_1, b_2\} \subset B_j$.

Define B_0 to be \aleph_{m+1} and A_0 to be empty. By induction suppose that $\langle A_i, B_i \rangle$ have been defined for $i \leq k < n$. Let C_k be the first \aleph_m elements of B_k . For each $b \in B_k \setminus C_k$ there exists a subset D_k of C_k and $c_{b,k} < \omega$ such that $|D_k| = \aleph_m$ and $f(\{b, x\}) = c_{b,k}$ for all $x \in D_k$. Now choose a finite set $A_b \subset D_k$ such that

f induces I_{k+1} on A_b . There are only \aleph_m finite subsets of C_k and a countable collection of possible values for $c_{b,k}$. Thus we can choose $B_{k+1} \subset B_k \setminus C_k$ of cardinality \aleph_{m+1} and $c_k < \omega$ such that $A_{b_1} = A_{b_2}$ for all $\{b_1, b_2\} \subset B_{k+1}$ and $c_{b,k} = c_k$ for all $b \in B_{k+1}$. We let $A_{k+1} = A_b$ for $b \in B_{k+1}$. It is easy to see that f induces the desired identity on the set $\bigcup \{A_i : 1 \leq i \leq n\}$. \dashv

THEOREM 6. *For all m such that $1 \leq m < \omega$ we have $I_m \in \mathcal{F}(\aleph_{m+1})$.*

PROOF. The proof is by induction on $k < \omega$. To show that $I_1 \in \mathcal{F}(\aleph_2)$ we let $J_1 = J_2$ be the trivial identity on two points and produce the identity J , the end-homogeneous amalgam of the sequence $\langle J_1, J_2 \rangle$. An analysis of J shows it to be I_1 . By induction we assume that $I_k \in \mathcal{F}(\aleph_{k+1})$. We then let $J_1 = J_2 = I_k$ and produce an identity K , the end-homogeneous amalgam of the sequence $\langle J_1, J_2 \rangle$. By the theorem we have that $K \in \mathcal{F}(\aleph_{k+2})$. An analysis of K shows it to be I_{k+1} . \dashv

The following theorem follow from Theorem 4 and Theorem 6.

THEOREM 7. *The consistency of ZFC implies the consistency of*

$$\mathcal{F}(\aleph_m) \subsetneq \mathcal{F}(\aleph_{m+1})$$

for $m \geq 1$.

The question that now remains to be answered is whether or not it is consistent that $\mathcal{F}(\aleph_2) = \mathcal{F}(\aleph_\omega)$. Since this is trivially true when CH holds, the question is only valid when a large continuum is also demanded. We have been successful in getting the consistency of the above statement when the continuum is large and the result will appear in the future.

REFERENCES

- [1] J. M. BAUMGARTNER, *Almost-disjoint sets*, *Annals of Mathematical Logic*, vol. 10 (1976), pp. 401–439.
- [2] J. SCHMERL, *Transfer theorems and their applications to logics*, *Model theoretic logics*, 1985, pp. 177–209.
- [3] S. SHELAH, *A two cardinal theorem and a combinatorial theorem*, *Proceedings of the American Mathematical Society*, vol. 62 (1977), no. 1, pp. 134–136.
- [4] ———, *Appendix to models with second order properties II*, *Annals of Mathematical Logic*, vol. 14 (1978), pp. 223–226.
- [5] ———, *Models with second order properties II*, *Annals of Mathematical Logic*, vol. 14 (1978), pp. 73–87.
- [6] S. SHELAH and L. STANLEY, *A theorem and some consistency results in partition calculus*, *Annals of Pure and Applied Logic*, vol. 36 (1987), pp. 119–152.

MATHEMATICS DEPARTMENT
SIMON FRASER UNIVERSITY
BURNABY, B. C., CANADA

E-mail: gilchris@cs.sfu.ca

INSTITUTE OF MATHEMATICS
THE HEBREW UNIVERSITY
JERUSALEM, ISRAEL

E-mail: shelah@math.huji.ac.il